PAPER DETAILS

TITLE: Factors Affecting Tolerance Development in Children With Food Allergies

AUTHORS: Yasemin ERSÖZLÜ, Hulya ANIL, Koray HARMANCI

PAGES: 79-87

ORIGINAL PDF URL: https://dergipark.org.tr/tr/download/article-file/2515862

Factors Affecting Tolerance Development in Children with Food Allergies

Besin Alerjisi Tanılı Hastalarda Tolerans Gelişimi ve Toleransa Etki Eden Faktörler

Yasemin Ersözlü, Hülya Anıl, Koray Harmancı

¹Department of Pediatric Infection Diseases Scholl of Medicine Acibadem University, Istanbul, Turkiye

²University of Eskişehir Osmangazi, Department of Pediatric Allergy and Immunology, Eskisehir, Turkiye

Abstract

Although the prevalence of food allergy (FA) is unknown, it is estimated to have increased in recent years. In general, food allergies are more common in the pediatric age group than in adults. In our study, we aimed to determine the development of tolerance and the factors affecting it in children with food allergies. Medical records of cases followed up for FA were retrospectively reviewed between 2013 and 2016. Gender, age of first symptom, duration of breastfeeding, family history of atopy, multiple food allergies, concomitant allergic diseases and clinical evidence of the cases were evaluated. In addition, prick test results, total IgE and specific IgE levels, food challenge test results and tolerance development were evaluated. Of 319 patients 57.7% (184) were male. The average age of onset of symptoms was 3 months. The most common symptoms were related to the gastrointestinal system (69.9%), the skin (22.3%) and the respiratory system (6.3%). When the tolerance development rates were investigated by years, it was found that the tolerance developed 87.1% in the first year, 5.9 % in the second and 2.8% in the third year. The most common allergens are cow's milk (71.3%), eggs (17.6%), peanuts (2.8%), nuts (2.04%) and other foods such as strawberries, cocoa, peaches. The mother's smoking status was significantly associated with the development of tolerance. The rate of development of tolerance in the first year was significantly higher in non-Ig E-mediated allergies and single food allergies. Keywords: Food allergy, Prognosis, Tolerance

Özet

Besin alerjisinin prevelansı bilinmemekle birlikte son yıllarda giderek arttığı tahmin edilmektedir. Genel olarak besin alerjileri pediatrik yaş grubunda erişkine oranla daha sık görülür. Çalışmamızda; polikliniğimizde gıda alerjisi tanısı olan hastalarımızın, doğal seyri, tolerans gelişimi ve toleransa etki eden faktörleri belirlemeyi hedefledik. Ocak 2013-Ocak 2016 yılı arasında polikliniğimizde Besin Alerjisi tanısı alan hastaların doşyaları retrospektif olarak incelendi. Hastaların cinsiyetleri, ilk semptom yaşı, anne sütü alma süresi, ailede atopi öyküsü, çoklu besin alerjisi, ek alerjik hastalık varlığı ve klinik bulguları değerlendirildi. Hastaların prik test sonuçları, total IgE, şpesifik IgE, düzey-leri ve tolerans geliştirme durumları değerlendirildi. 319 hastanın 184'ü erkek (%57.7) idi. Hastaların 127'sinde (%39.8) ailede atopi öyküsü ve 71'inde (%22.3) çoklu besin alerjisi mevcuttu. Hastalarda görülen en sık semptom, %65.8 gastrointestinal sistem, %21 cilt bulguları, %6.3 solunum bulgularıydı. Hastaların semptom yaşı 5.8 \pm 9,7 ay; tanı yaşı 6.6 \pm 9.08 aydı. Hastaların yıllara göre tolerans geliştirme oranları; Birinci yıl 287 (%87,1) ikinci yıl 19'unda (%5,9) üçüncü yıl 9'unda (% 2,8) olarak saptandı. En sık alerjen gıda inek sütü (%71,3), yumurta (%17,6), fistık (2,8), findık (%2,04) olmakla beraber diğer gıdaları çilek, kakao, şeftali, buğday, soya, balık olarak saptandı. Çalışmamızda ilk yılda annenin sigara kullanımı ile besin alerjierinde tolerans gelişimi arasında anlamlı fark saptanmıştır. Non Ig E aracılı besin alerjisi olanlar ve tek besine karşı alerjen olan hastalar ilk yılda tolerans gelişimi anlamlı oranda faza saptanmıştır.

Correspondence: Yasemin ERSÖZLÜ Department of Pediatric Infection Diseases Scholl ofMedicinei Acibadem Universit, Istanbul, Turkiye e-mail: yaseminersozlu@gmail.com

Received 30.06.2022 Accepted 10.10.2022 Online published 28.10.2022

Ersozlu Y, Anil H, Harmanci K, Factors Affecting Tolerance Development in Children with Food Allergies, Osmangazi Journal of Medicine, 2023;45(1): 79-87 Doi: 10.20515/otd.1138002

1. Introduction

There is a significant increase in childhood allergic diseases worldwide. The prevalence of these diseases varies from country to country and even in different regions of the same country (1). Since these changes cannot be explained by genetic factors alone, environmental factors are also blamed.

The main allergenic foods are cow's milk, chicken eggs, soybeans, wheat in childhood, peanuts, shellfish, fish and nuts in adults. Other allergenic foods include legumes, vegetables, fruits, beef, lamb, pork, chicken, and turkey meat (2-7). Sensitization with food antigen can occur not only by ingestion of food but also after inhalation and skin contact. In addition, cross-reactions can occur with allergens such as birch pollen, profilins, latex, and lipid transfer protein (8,9).

The healthy immune system maintains unresponsiveness towards to potential food allergens. Various hypotheses such as integration of vitamin D deficiency, hygiene and dual allergen exposure hypothesis have been proposed to explain the increase in FA. Morst recently investigated hypothesis is dual allergen hypothesis. Randomise controlled trials recommend including possible common food allergies such as peanuts and eggs in complementary foods nutritional regimens starting at approximately 6 months but not before 4 months. National Institutes of Health recommendations promotes the early introduction of peanut for the prevention of peanut allergy(46). Oppositevly the study Enquiring about tolerane (EAT) suggest that early introduction of allergenic food in sufficient quantity from 3 months of age may be able to help prevent food allergies developing in children(47). Future advances understanding oral tolerance in human subjects will largely determine treatment.

In our study; we aimed to determine the factors that affect the natural course, development of tolerance and tolerance of our patients diagnosed with food allergy.

2. Materials and Methods

Between January 2013 and January 2016, the files of the patients diagnosed with Food Allergy in the outpatient clinic of Eskişehir Osmangazi University Faculty of Medicine, Department of Pediatrics, AllergyImmunology Department were retrospectively analyzed according to the questionnaire form prepared. Local ethical committees approved all studies.

Food allergy was diagnosed by clinical history, allergy tests, elimination of cow's milk from the diet and food challenge test. The diagnosis of non-IgE mediated food allergy was made in the presence of clinical symptoms, the disappearance of symptoms with elimination diet and the occurrence of symptoms when suspected food was given again. From the polyclinic files of the children who were followed up with a diagnosis of food allergy, gender, first symptom age, breastfeeding time, age at starting additional foods, family history of atopy, multiple food allergy, presence of additional allergic disease, inhaled allergen sensitivity, and clinical findings were obtained. The time interval between milk protein intake and reaction formation and the pattern of milk protein intake were evaluated. Prick test, prick to prick test results with pasteurized milk, total IgE, milk and casein specific IgE levels and reactions food challenge test were evaluated. Tolerance development status of the patients was evaluated according to the results of the loading test performed both at home or in clinic and information received from the families.

In statistical analysis, chi-square test was used in comparison of categorical data, T test (Student's T test) in independent groups in case of assumptions in comparison of numerical measurements, and MannWhitney U test if assumptions were not provided. Mann Whitney U test was used for comparing two independent groups, while Pearson chisquare and Fisher's chi-square tests were used for comparison of categorical variables. Two ratio tests were used in the frequency distribution test (f) and chi-square analysis was used in the test of cross tables. A value of < 0.05 considered statistically was р significant in all tests.

3. Results

Between January 2013 and December 2016, 416 children who were followed up with a diagnosis of food allergy were found. Fortythree children with a diagnosis of food allergy excluded during their follow-up and 16 patients with a diagnosis of inflammatory bowel disease were excluded from the study. 17 patients whose ICD codes were entered incorrectly and 21 patients whose follow-up data could not be accessed were excluded from the study. A total of 319 patients were included in the study. The characteristics of the patients included in the study are shown in Table 1.

According to immunological mechanisms, 220 of the patients (69.9%) had non-IgEmediated, 48 (15%) had Ig E-mediated, and 51 (16%) had mixed disease. The clinical findings of the patients are summarized in Table 2.

Table 1. Characteristics of patients with food allergies

n
184 (57,7%)
135 (42,3%)
3 (2-6)
12 (9-17,25)
6 (5-6)
127 (39,8%)
72 (22,6%)
24 (7,5%)
101 (31,3%)
9/101
18/101
74/101

Table 2. Clinical findings of patients with food allergies

Clinical finding	n (%)
Skin manifestations	71 (22,3)
Urticaria	7 (9,9)
Atopic dermatitis	60 (84,5)
Angioedema	4 (5,6)
GIS findings	223 (69,9)
Bloody stool	61 (27,35)
Mucus stool	72(32,28)
Bloody mucus stool	41(18,3)
Vomiting	12 (5,38)
Abdominal pain	10 (4,48)
Diarrhea	11 (4,93)
Constipation	14 (6,27)
Bloody vomiting	2 (0,9)
Respiratory symptoms	20 (6,3)
Rhinitis	3 (15)
Wheezing	9 (45)
Cough	8 (40)
Anaphylaxis	5 (1,6)

The most common symptoms in children with food allergy were GIS findings in 223 (69,9%) patients, skin findings in 71 (22,3%) patients, and respiratory system findings in 20 (6,3%) patients. It was determined that five (1,6%) of the patients developed anaphylaxis after consuming food which is an allergen.

When the patients were separated according to food allergens, the most common allergen was

cow's milk (71,3%). Other common allergens were eggs (17,6%), peanuts (2,8%), and hazelnuts (2,04%).

In the first year controls while tolerance development was observed in 278 (87,1%) of patients, it did not develop in 40 (12,5%) of patients (Table 3). Tolerance did not develop in 22 patients at the second year and in 13 patients at the third year follow-up.

Tolerance development	First year	Second year	Third year
No	41 (12,8%)	22 (6,8%)	13 (4%)
Yes	278 (87,1%)	19 (5,9%)	9 (2,8%)
Total	319	41 (12,8%)	22 (6,9%)

Table 3. Distribution of tolerance development by years

In tolerance development; gender, family history of atopy, social environment, breastfeeding time, prematurity, starting month of supplementary food, eosinophilia, total and specific IgEs, food prick test were not significant. The mothers of 267 (96%) of 278 (87,1%) patients with tolerance development in the first year were nonsmokers (p < 0.01). In patients who developed tolerance in the second year and in the third year, the history of smoking in the mother was lower than those who did not develop but the difference was tolerance, not statistically significant.

While 225 (91,1%) of 278 (87,1%) children with tolerance development at the end of the first year showed an allergic reaction to a single food; 53 (19,1%) of them are allergic to multiple foods. Those who are allergic to a single food develop more tolerance than patients with multiple food allergies, which is statistically significant (p <0.01). All children with tolerance development in the second year had a single food allergy. Tolerance did not

Table 4: Variables affecting tolerance development

develop in those with multiple food allergies (p < 0.001).

When allergen types are classified as cow's milk, egg and other; 209 (75,2%) of the children with tolerance development in the first year were allergic to cow milk, 15 (5,4%) to eggs, 54 (19,4%) to other foods (p < 0.01). Only 1 (5,3%) of 19 children who developed tolerance in the second year were allergic to eggs (p <0.01). 2 (22,2%) of 9 children with tolerance development in the third year were allergic to eggs (p = 0.544). The development of tolerance in cow's milk allergy was found to be significantly higher than other allergens. When the characteristics of 13 patients without tolerance development in the third year were examined, 7 (53,8%) were male and 6 (46,2%) were female. 12 (92,3%) were living in the urban area. The mothers of 10 (76,9%) were not smoking. 6 (46,1%) of them had non Ig E mediated disease. Twelve (92,3%) of them had multiple food allergies. Variablers affecting tolarence development are summarized in Table 4.

		First year tolerance development			Second year tolerance development			Tertian year tolerance development		
		No	Yes	<i>p</i> value	No	Yes	<i>p</i> value	No	Yes	<i>p</i> value
ing	No	28 %68,3	267 %96	<0,01	17 %77,3	11 %57,9	>0,05	10 %76,9	7 %77,8	>0,05
Smok	Yes	13 %31,7	11 %4		5 %22,7	8 %42,1		3 %23,1	2 %22,2	
e sm	Ig E	14 %34,1	34 %12,5	<0,01	9 %40,9	5 %26,3	>0,05	4 %30,8	5 %55,6	>0,05
İmmun mechanis	NonIgE	21 %51,2	199 %71,6		9 %40,9	12 %63,2		6 %46,2	3 %33,3	
Num ber	Mixed Multiple food	6 %14,6 19	45 %16,2 53	<0,01	4 %18,2 19	2 %10,5 0	<0,01	3 %23,2 12	1 %11,1 7	>0,05

		%46,3	%19,1		%86,4			%92,3	%77,8	
	Single food	22 %53,7	225 %80,9		3 %13,6	19 %100		1 %7,7	2 %22,2	
be	Cow milk	16 %39	209 %75,2	<0,01	0	16 %64,2	<0,01	0	0	>0,05
llergen ty	Egg	2 %4,9	15 %5,4		2 %9,1	1 %5,3		0	2 %22,7	
A	Other	23 %56,1	54 %19,4		20 %90,9	2 %10,5		13 %100	7 %77,8	

Bold entries are statistically significant p values

4. Discussion

In many Western countries, diseases such as asthma and atopic dermatitis and allergic rhinitis have been increasing in the last 50-60 years and constitute a significant burden on the society and health systems (11-15). It is thought that the frequency of food allergy has increased in the past 10-20 years with atopic diseases and this situation is due to many different risk factors (16-20).

When studies on food allergies in the literature are examined, the most common finding is skin reactions, followed by findings gastrointestinal (14-18).GIS symptoms have been reported as 30.3% (22) in France, 49.3% (23) in the UK, 20.9% (24) in Lithuania, and 27.6% (21) in the European country. Skin reactions were followed by gastrointestinal system finding(23). In our study, the most common symptoms in children with food allergy were related to the gastrointestinal system (GIS) (69.9%). This was followed by skin findings in 22.3% of patients and respiratory system symptoms in 6.3% of patients. The reason for our different findings can be attributed to the difference in the patient population referred to as our hospital, a top-level healthcare institution.

When the children were categorized according to food allergens, the most common allergen was cow's milk (71.3%). This was followed by egg (17.6%), peanut (2.8%), and hazelnut (2.04%). Strawberry, cocoa, peach, wheat, soy and fish were other responsble food allergens. In the study conducted by Orhan et al. (20) in children aged 6-9 years, the most common allergenic foods were red meat (31,8%), cow's milk (18,1%), cocoa (18,1%), chicken egg (13,1%) and kiwi (13,6%) respectively has been reported. However, foods that cause allergies change with age. In our study, we attribute the difference of the foods most frequently responsible to food allergies from other studies conducted in our country to the age group evaluated. Egg and milk are the most common causes of food allergy in all countries (1,21,23,24).

Most of the patients develop tolerance to foods in advanced ages (28,29). In our study, tolerance developed in 87.5% of the children in the first year, 6.8% in the second year, and 2.8% in the third year. Shek (28) showed that 28 of 66 children with egg allergy and 16 of 33 children with cow's milk allergy developed tolerance over time. Similarly, Dannaeus et al. (30) found that 4 out of 12 children with cow's milk allergy, 20 out of 55 children with egg allergy, 5 out of 32 children with fish allergy developed tolerance, but none of the cases with peanut and tree nut allergy developed tolerance. Dias (19) reported in a retrospective study that 44% of 79 children with cow's milk allergy were still allergic by the time they reached the age of 10. In a study conducted in Sweden, it was found that 50% of children with egg allergy passed their allergies by the age of 3 (32). In another study, 52% of children with egg allergy developed tolerance at the age of 3 and 66% at the age of 5 (33). Savage (34) found that the development of tolerance to egg allergy was slower than predicted. The results of this study suggest that, contrary to what was previously known, egg allergy is not a food allergy that usually occurs at school age, but a food allergy that is

likely to persist in later life. Sicherer (35) showed in his multicenter observational study that age of tolerance is not as high as suggested by Savage. The different results of these studies on tolerance developmental ages seem to be influenced by many different factors such as genetic and phenotypic characteristics of the studied population and inclusion criteria.

Although passive smoking is known to be a risk factor for asthma, there is no information about whether it increases the risk of food allergy (25,36,37). There are publications indicating that smoking during pregnancy increases the prevalence of allergic diseases (38,39). Avoiding smoking during pregnancy is recommended for primary prevention from allergic diseases (40). In our study, the mothers of 96% of the patients with tolerance development in the first year were not smoking. Only 4% of their mothers were smoking. A significant relationship was found between the mother's smoking and the development of tolerance in the first year. The mothers of the children with tolerance development in the second and third years smoked at a higher rate than the mothers of the children who developed tolerance in the first year. The smoking rates of the mothers of the children with and without tolerance in the second and third years were not statistically different.

The majority of patients (80.9%) who developed tolerance at the end of the first year had allergies to a single food. These patients develop a higher rate of tolerance than those who are allergic to more than one food. In a study, it was reported that 18% of patients who developed tolerance until the age of 4 were also allergic to other foods and that multiple food allergy had no effect on the development of tolerance (41). Multiple food allergies have been reported in 3.5% of children under 2 years of age and 30.5% of children over 2 years of age with persistent cow's milk allergy (42). In the same study, the presence of allergies to other foods was reported as a risk factor for persistent cow's milk allergy. In a study investigating the differences between single food allergy and multiple food allergy, although not statistically significant, food allergy remission at 2 years of age was found to be less in those with multiple food allergies. In accordance to

our study another multicenter study revealed multipl food allergy is a risk factor for late tolerance. Although different results were reported in the literature, it was observed in our study that multiple food allergies had no effect on the development of tolerance. In our study, we think that the effect of multiple food allergies on the development of tolerance may change after the patients are exposed to other food allergens in the next follow-up period.

191 (68.7%) of 278 children who had tolerance development in the first year started to consume complementary feeding after the 4th month. 87 of them (31.3%) switched to additional food before 4 months. All 19 children with tolerance development in the second year and all 9 children with tolerance development in the third year switched to additional foods after the 4th month. A statistically significant relationship was not found between the age of starting the supplementary food and the development of tolerance. It was suggested that families with a high risk of allergic diseases should not give additional food to their atopic children in the first six months. It was recommended to postpone dairy products until the 12th month, eggs until the 24th month, peanuts, hazelnuts, fish and seafood until the 36th month. However, after 2008, European Society for Paediatric Gastroenterology, Hepatology and Nutrition (ESPGHAN) and the American Pediatrics Academy of (APA) are recommended to start supplementary foods in the first 4-6 months (29,34).

When the allergen types are classified in 3 groups as cow's milk, egg and others; Of the 278 patients with tolerance development in the first year, 209 were allergic to cow's milk, 15 to eggs, and 54 to other foods. Tolerance development was higher in those with cow's milk allergy than those allergic to eggs and other foods. There have been several studies of natural hiatory of cow's milk allergy and predictors of remission. The largest multicentre European case series in food EuroPrevall allergy, study, in which standardized evaluation including DBPCFC, population with DBPCFC, %57 comfirmed cow's milk allergy developed tolerance within 1 vear(48).

199 (71.6%) of 278 (87.1%) allergy diagnosed patients who developed tolerance in the first year is non Ig E, 45 (16.2%) of the sample is

mixed. 34 (12.2%) of them are food by Ig Emediated mechanisms. Likely others we found Ig E mediated aleergies are more resistant to tolerance. Of %21 Children aged 16 in which had increased cow's milk spesific IgE were

REFERENCES

- Woods RK, Abramson M, Bailey M, Walters EH. International prevalences of reported food allergies and intolerances. Comparisons arising from the European Community Respiratory Health Survey (ECRHS) 1991-1994. *Eur J Clin Nutr.* 2001;55:298-304.
- 2. Bischoff S, Crowe SE. Food allergy and the gastrointestinal tract. *Curr Opin Gastroenterol*. 2004 Mar;20:156-61.
- Ko J, Mayer L. Oral tolerance: lessons on treatment of food allergy. Eur J Gastroenterol 21Hepatol. 2005;17:1299-303.
- Brandtzaeg PE. Current understanding of gastrointestinal immunoregulation and its relation to food allergy. *Ann N Y Acad Sci.* 2002;964: 13-45.
- Chin JJ. Revisiting the "hygiene hypothesis" in gastrointestinal allergy. *Curr Opin Gastroenterol*. 2002;18:705-10.
- 6. Crespo JF, Rodriguez J. Food allergy in adulthood. *Allergy*. 2003;58:98-113.
- van Ree R. Clinical importance of crossreactivity in food allergy. *Curr Opin Allergy Clin Immunol.* 2004;4:235-40.
- Bock SA, Sampson HA, Atkins FM, Zeiger RS, Lehrer S, Sachs M, Bush RK, Metcalfe DD. Double-blind, placebo-controlled food challenge (DBPCFC) as an office procedure: a manual. J Allergy Clin Immunol. 1988 ;82:986-97.
- Deckers IA. McLean S. Linssen S. Mommers M. van Schayck CP. Sheikh A. Investigating international time trends in the incidence and prevalence of atopic eczema 1990-2010: a systematic review of epidemiological studies. PLoS One 2012;7:e39803.
- Anandan C, Gupta R, Simpson CR, Fischbacher C, Sheikh A. Epidemiology and disease burden from allergic disease in Scotland: analyses of national databases. J R Soc Med. 2009;102:431-42.
- Anandan C, Nurmatov U, van Schayck OC, Sheikh A. Is the prevalence of asthma declining? Systematic review of epidemiological studies. *Allergy.* 2010;65:152-67.
- Gupta R, Sheikh A, Strachan DP, Anderson HR. Burden of allergic disease in the UK: secondary analyses of national databases. *Clin Exp Allergy*. 2004;34:520-6.
- 13. Schoenwetter WF, Dupclay L Jr, Appajosyula S, Botteman MF, Pashos CL. Economic impact and

notified non- tolerated (49). The strict elimination diet resulted in improvement of symptoms and tolerance to allergic food usually with in 1 year (43).

quality-of-life burden of allergic rhinitis. Curr Med Res Opin. 2004;20:305-17.

- 14. Allen KJ, Koplin JJ. The epidemiology of IgEmediated food allergy and anaphylaxis. *Immunol Allergy Clin North Am.* 2012 ;32:35-50.
- Sampson HA. Food allergy--accurately identifying clinical reactivity. *Allergy*. 2005;60 Suppl 79:19-24.
- 16. Prescott S, Allen KJ. Food allergy: riding the second wave of the allergy epidemic. *Pediatr Allergy Immunol.* 2011;22:155-60.
- 17. Lack G. Update on risk factors for food allergy. J Allergy Clin Immunol. 2012 ;129:1187-97.
- Sicherer SH. Epidemiology of food allergy. J Allergy Clin Immunol. 2011;127:594-602.
- Dias A, Santos A, Pinheiro JA. Persistence of cow's milk allergy beyond two years of age. *Allergol Immunopathol (Madr)*. 2010;38:8-12.
- Orhan F, Karakas T, Cakir M, Aksoy A, Baki A, Gedik Y. Prevalence of immunoglobulin Emediated food allergy in 6-9-year-old urban schoolchildren in the eastern Black Sea region of Turkey. Clin Exp Allergy. 2009;39:1027-35.
- Steinke M, Fiocchi A, Kirchlechner V, Ballmer-Weber B, Brockow K, Hischenhuber C, Dutta M, Ring J, Urbanek R, Terracciano L, Wezel R; REDALL study consortium. Perceived food allergy in children in 10 European nations. A randomised telephone survey. *Int Arch Allergy Immunol.* 2007;143:290-5.
- Rancé F, Grandmottet X, Grandjean H. Prevalence and main characteristics of schoolchildren diagnosed with food allergies in France. *Clin Exp Allergy*. 2005;35:167-72.
- Rona RJ, Chinn S. Parents' perceptions of food intolerance in primary school children. *Br Med J* (*Clin Res Ed*). 1987 ;294:863-6.
- 24. Dubakiene R. Kavaliünas A. Surkiene G. Zagminas K. StukasR. Food allergies amongVilnius city primary school children. *Biologija* 2011;57:103–9.
- Kristjansson I, Ardal B, Jonsson JS, Sigurdsson JA, Foldevi M, Björkstén B. Adverse reactions to food and food allergy in young children in Iceland and Sweden. *Scand J Prim Health Care*. 1999;17:30-4.
- 26. Fiocchi A, Assa'ad A, Bahna S; Adverse Reactions to Foods Committee; American College of Allergy, Asthma and Immunology. Food allergy and the introduction of solid foods to infants: a consensus document. Adverse

Reactions to Foods Committee, American College of Allergy, Asthma and Immunology. *Ann Allergy Asthma Immunol.* 2006;97:10-20; quiz 21, 77.

- 27. Liu AH, Jaramillo R, Sicherer SH, Wood RA, Bock SA, Burks AW, Massing M, Cohn RD, Zeldin DC. National prevalence and risk factors for food allergy and relationship to asthma: results from the National Health and Nutrition Examination Survey 2005-2006. J Allergy Clin Immunol. 2010;126:798-806.e13.
- Shek LP, Soderstrom L, Ahlstedt S, Beyer K, Sampson HA. Determination of food specific IgE levels over time can predict the development of tolerance in cow's milk and hen's egg allergy. J Allergy Clin Immunol. 2004;114:387-91.
- 29. Sicherer SH. Epidemiology of food allergy. J Allergy Clin Immunol. 2011;127:594-602.
- Dannaeus A, Inganäs M. A follow-up study of children with food allergy. Clinical course in relation to serum IgE- and IgG-antibody levels to milk, egg and fish. *Clin Allergy*. 1981;11:533-9.
- 31. Agostoni C, Decsi T, Fewtrell M, Goulet O, Kolacek S, Koletzko B, Michaelsen KF, Moreno L, Puntis J, Rigo J, Shamir R, Szajewska H, Turck D, van Goudoever J; ESPGHAN Committee on Nutrition:. Complementary feeding: a commentary by the ESPGHAN Committee on Nutrition. J Pediatr Gastroenterol Nutr. 2008;46:99-110.
- 32. Hattevig G, Kjellman B, Björkstén B. Clinical symptoms and IgE responses to common food proteins and inhalants in the first 7 years of life. *Clin Allergy*. 1987;17:571-8.
- Boyano-Martínez T, García-Ara C, Díaz-Pena JM, Martín-Esteban M. Prediction of tolerance on the basis of quantification of egg white-specific IgE antibodies in children with egg allergy. J Allergy Clin Immunol. 2002;110:304-9.
- Savage JH, Matsui EC, Skripak JM, Wood RA. The natural history of egg allergy. J Allergy Clin Immunol. 2007;120:1413-7.
- 35. Sicherer SH, Wood RA, Vickery BP, Jones SM, Liu AH, Fleischer DM, Dawson P, Mayer L, Burks AW, Grishin A, Stablein D, Sampson HA. The natural history of egg allergy in an observational cohort. *J Allergy Clin Immunol*. 2014 ;133:492-9.
- Cook DG, Strachan DP. Health effects of passive smoking-10: Summary of effects of parental smoking on the respiratory health of children and implications for research. *Thorax.* 1999;54:357-66.
- Strachan DP, Cook DG. Health effects of passive smoking. 6. Parental smoking and childhood asthma: longitudinal and case-control studies. *Thorax.* 1998;53:204-12.
- Tariq SM, Matthews SM, Hakim EA, Stevens M, Arshad SH, Hide DW. The prevalence of and risk factors for atopy in early childhood: a whole population birth cohort study. J Allergy Clin Immunol. 1998;101:587-93.
- 39. Ponsonby AL, Couper D, Dwyer T, Carmichael A. Cross sectional study of the relation between sibling number and asthma, hay fever, and eczema. *Arch Dis Child*. 1998;79:328-33.

- 40. Arshad SH. Primary prevention of asthma and allergy. *J Allergy Clin Immunol*. 2005;116:3-14; quiz 15.
- 41. Skripak JM, Matsui EC, Mudd K, Wood RA. The natural history of IgE-mediated cow's milk allergy. *J Allergy Clin Immunol.* 2007;120:1172-7.
- 42. Santos A, Dias A, Pinheiro JA. Predictive factors for the persistence of cow's milk allergy. *Pediatr Allergy Immunol.* 2010;21:1127-34.
- 43. Meyer R, Chebar Lozinsky A, Fleischer DM, Vieira MC, Du Toit G, Vandenplas Y, Dupont C, Knibb R, Uysal P, Cavkaytar O, Nowak-Wegrzyn A, Shah N, Venter C. Diagnosis and management of Non-IgE gastrointestinal allergies in breastfed infants-An EAACI *Position Paper. Allergy.* 2020;75:14-32.
- Koksal BT, Barıs Z, Ozcay F, Yilmaz Ozbek O. Single and multiple food allergies in infants with proctocolitis. *Allergol Immunopathol (Madr)*. 2018;46:3-8.
- 45. Buyuktiryaki B, Kulhas Celik I, Erdem SB, Capanoglu M, Civelek E, Guc BU, Guvenir H, Cakir M, Dibek Misirlioglu E, Akcal O, Volkan B, Toyran M, Sag E, Kertel AC, Ginis T, Kocabas CN, Orhan F, Can D. Risk Factors Influencing Tolerance and Clinical Features of Food Protein-induced Allergic Proctocolitis. J Pediatr Gastroenterol Nutr. 2020;70:574-79.
- 46. Du Toit G, Sampson HA, Plaut M, Burks AW, Akdis CA, Lack G. Food allergy: Update on prevention and tolerance. *J Allergy Clin Immunol*. 2018;141:30-40.
- 47. Perkin MR, Logan K, Marrs T, Radulovic S, Craven J, Flohr C, Lack G; EAT Study Team. Enquiring About Tolerance (EAT) study: Feasibility of an early allergenic food introduction regimen. J Allergy Clin Immunol. 2016;137:1477-1486.e8.
- 48. Fernández-Rivas M, Barreales L, Mackie AR, Fritsche P, Vázquez-Cortés S, Jedrzejczak-Czechowicz M, Kowalski ML, Clausen M, Gislason D, Sinaniotis A, Kompoti E, Le TM, Knulst AC, Purohit A, de Blay F, Kralimarkova T, Popov T, Asero R, Belohlavkova S, Seneviratne SL, Dubakiene R, Lidholm J, Hoffmann-Sommergruber K, Burney P, Crevel R, Brill M, Fernández-Pérez C, Vieths S, Clare Mills EN, van Ree R, Ballmer-Weber BK. The EuroPrevall outpatient clinic study on food allergy: background and methodology. *Allergy*. 2015;70:576-84.
- 49. Longo G, Berti I, Burks AW, Krauss B, Barbi E. IgE-mediated food allergy in children. *The Lancet.* 2013;382:1656-64.

Ethics

Ethics Committee Approval: The study was approved by Eskişehir Osmangazi University Noninterventional Clinical Research Ethical Committee (Number: 24, Date: 14.11.2016).

Informed Consent: The authors declared that it was not considered necessary to get consent from the patients because the study was a retrospective data analysis.

Authorship Contributions: Surgical and Medical Practices:Y.E ,H.A, K.H. Concept:Y.E ,H.A, K.H. Design:Y.E ,H.A, K.H. Data Collection or Processing: Y.E. Analysis or Interpretation: Y.E ,H.A, K.H. Literature Search:Y.E ,H.A, K.H. Writing:Y.E, H.A., K.H.

Copyright Transfer Form: Copyright Transfer Form was signed by all authors. Peer-review: Internally peer-reviewed. Conflict of Interest: No conflict of interest was declared by the authors. Financial Disclosure: The authors declared that this study received no financial support.

©Copyright 2023 by Osmangazi Tıp Dergisi - Available online at tip.ogu.edu.tr ©Telif Hakkı 2023 ESOGÜ Tıp Fakültesi - Makale metnine dergipark.org.tr/otd web sayfasından ulaşılabilir.