

## PAPER DETAILS

TITLE: DOES JASMONIC ACID PREVENT THE GERMINATION

AUTHORS: Kürsat ÇAVUSOGLU,Kudret KABAR

PAGES: 35-41

ORIGINAL PDF URL: <https://dergipark.org.tr/tr/download/article-file/116216>

## DOES JASMONIC ACID PREVENT THE GERMINATION OF BARLEY SEEDS?

Kürşat ÇAVUŞOĞLU\*, Kudret KABAR

Süleyman Demirel University, Faculty of Arts and Science, Biology Department,  
32260, Isparta, Turkey

\*Corresponding author, e-mail: kursat@fef.sdu.edu.tr, fax: 0 246 237 11 06

Received: 12 September 2006, Accepted: 10 October 2006

**Abstract:** Effect of jasmonic acid on seed germination and seedling growth of barley (*Hordeum vulgare* L. cv. Bülbül 89) was investigated in the present study. Jasmonic acid concentrations less than 1500  $\mu$ M have not inhibited the seed germination, while 1500 and 2000  $\mu$ M jasmonic acid levels caused atypical germination. The germination was completely inhibited at 3000  $\mu$ M level of jasmonic acid. However, the seedling growth clearly slowed down with increasing concentrations of jasmonic acid. Furthermore, these inhibitions on the seed germination and seedling growth were irreversible.

**Key words:** *Hordeum vulgare*, jasmonic acid, seed germination, seedling growth

## JASMONİK ASİT ARPA TOHURLARININ ÇİMLENMESİNİ ENGELLER Mİ?

**Özet:** Bu çalışmada, arpanın (*Hordeum vulgare* L. var. Bülbül 89) tohum çimlenmesi ve fide büyümesi üzerine jasmonik asitin etkisi araştırılmıştır. Jasmonik asidin 1500  $\mu$ M'dan daha düşük konsantrasyonları tohum çimlenmesini engellemezken, 1500 ve 2000  $\mu$ M jasmonik asit düzeyleri atipik çimlenmeye neden olmuştur. 3000 $\mu$ M jasmonik asit düzeyinde ise çimlenme tamamen engellenmiştir. Bununla birlikte, fide büyümesi jasmonik asit konsantrasyonlarındaki artışlar ile belirgin bir şekilde yavaşlamıştır. Dahası, tohum çimlenmesi ve fide büyümesi üzerindeki bu engellemelerin geri dönüşümsüz olduğu tespit edilmiştir.

**Anahtar kelimeler:** *Hordeum vulgare*, jasmonik asit, tohum çimlenmesi, fide büyümesi

## INTRODUCTION

Jasmonic acid (JA) and its methyl ester (JA-Me) have been proposed as naturally occurring plant growth regulators because of their wide distribution (PARTIER 1991). These compounds were isolated from fungi (CROSS & WEBSTER 1970, FERNANDEZ-MACULET & YANG 1992), algae (KRUPINA & DATHE 1991) and many higher plants (MEYER et al. 1984). Jasmonates play an important role in various physiological processes connected with plant growth and development, such as seedling and stem growth (DATHE et al. 1981, KODA et al. 1991), leaf abscission (CURTIS 1984), leaf senescence (UEDA et al. 1981), inhibition chlorophyll synthesis (ABELES

et al. 1989), reduction photosynthesis (MASLENKOVA et al. 1990), callus growth (UEDA & KATO 1982), synthesis of storage proteins (ANDERSON et al. 1989) and respiratory activity (POPOVA et al. 1988).

Exogenous jasmonates seem to have inhibitory effects on germination in many cases of nondormant seeds, for example, sunflower (CORBINEAU et al. 1988), amaranth (KEPCZYNSKI & BIALECKA 1994), lettuce (YAMANE et al. 1981), rapeseed and flax seeds (WILEN et al. 1991). But, jasmonates break the seed dormancy of apple (RANJAN & LEWAK 1992, 1995) and other woody species (DALETSKAYA & SEMBDNER 1989) and promote germination. However, the physiological roles of jasmonates in seed germination still remain unclear.

In view of the conflicting evidence referred to above, the aim of the present work was to check whether JA contributes germination of barley. For similar reason the effect of JA on seedling growth was checked.

## MATERIAL AND METHODS

Barley seeds were used in the present study. The seeds were surface sterilized with 1.0 % sodium hypochloride. 25 seeds were placed in 10 cm petri dishes lined by two sheets of Whatman No.1 filter paper moistened with sufficient amount of solutions of 0 (control, C), 1, 3, 5, 10, 20, 30, 40, 50, 100, 200, 400, 500, 1000, 1500, 2000 and 3000  $\mu$ M JA. Afterwards, they were left in an incubator to germinate at 20 °C, in continuous dark. The seeds were considered germinated when the radicles reached to 10 mm in length. The germination percentages were recorded after incubation for 7 days. At the end of 7<sup>th</sup> day, following the estimation of germination percentage, final percentage of coleoptile emergence, radicle and coleoptile lengths were determined. Furthermore, with special replicas for the experiments fresh weight of the seedlings were recorded.

Seeds kept for 7 days at 20 °C, in germination media of 1500, 2000 and 3000  $\mu$ M JA did not germinate. When these seeds were removed from JA solutions and transferred into petri dishes with distilled water, they were germinated as previously described. Each treatment was repeated 4 times. Statistical analyses were done using SPSS program and Duncan's multiple range test.

## RESULTS

Table 1 shows the time courses of the germination of barley seeds kept in the various concentrations of JA. The seeds started to germinate easily when incubating in the C medium. Germination percentage of C seeds was 96 % at the 2<sup>th</sup> day of incubation. JA concentrations under 500  $\mu$ M have not slowed the germination down. In the other words, these concentrations showed the same effect as C group. 500 and 1000  $\mu$ M JA partly delayed the germination. Conversely, the seed germination was completely inhibited at 1500  $\mu$ M JA.

**Table 1.** The time course of the germination of *Hordeum vulgare* kept in various concentrations of JA

| JA<br>concentrations<br>( $\mu\text{M}$ ) | Germination, % |     |     |     |     |     |     |
|---|----------------|-----|-----|-----|-----|-----|-----|
|   | Days           |     |     |     |     |     |     |
|   | 1              | 2   | 3   | 4   | 5   | 6   | 7   |
| C   | 0.0            | 100 | 100 | 100 | 100 | 100 | 100 |
| 1   | 0.0            | 100 | 100 | 100 | 100 | 100 | 100 |
| 3   | 0.0            | 100 | 100 | 100 | 100 | 100 | 100 |
| 5   | 0.0            | 100 | 100 | 100 | 100 | 100 | 100 |
| 10  | 0.0            | 100 | 100 | 100 | 100 | 100 | 100 |
| 20  | 0.0            | 100 | 100 | 100 | 100 | 100 | 100 |
| 30  | 0.0            | 100 | 100 | 100 | 100 | 100 | 100 |
| 40  | 0.0            | 100 | 100 | 100 | 100 | 100 | 100 |
| 50  | 0.0            | 100 | 100 | 100 | 100 | 100 | 100 |
| 100                                       | 0.0            | 100 | 100 | 100 | 100 | 100 | 100 |
| 200                                       | 0.0            | 100 | 100 | 100 | 100 | 100 | 100 |
| 400                                       | 0.0            | 0.0 | 100 | 100 | 100 | 100 | 100 |
| 500                                       | 0.0            | 0.0 | 100 | 100 | 100 | 100 | 100 |
| 1000                                      | 0.0            | 0.0 | 0.0 | 0.0 | 100 | 100 | 100 |
| 1500                                      | 0.0            | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 2000                                      | 0.0            | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 3000                                      | 0.0            | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |

In parallel with concentration rise, a negative effect of JA has increased all the growth parameters (final percentages of germination and coleoptile emergence, radicle and coleoptile elongation and fresh weight of the seedlings). For instance, germination percentage of C seeds was 100 % at 7<sup>th</sup> day in distilled water. The concentrations of JA under 1500  $\mu\text{M}$  have not inhibited final germination percentage of barley seeds. These concentrations showed the same effect as C group. But the germination was completely inhibited by JA concentrations more than 1500  $\mu\text{M}$ . Furthermore, 1500 and 2000  $\mu\text{M}$  JA levels caused atypical germination (coleoptile emerges without radicle emergence). Atypical germination percentage was 100 % and 74 % at 1500 and 2000  $\mu\text{M}$  JA, respectively. On the other hand, coleoptile emergence of barley seedlings was completely inhibited at 3000  $\mu\text{M}$  JA level only (Table 2).

Similarly, radicle and coleoptile lengths and fresh weights of the seedlings slowed down with increasing levels of JA. Although, the radicle elongation was completely inhibited by JA concentrations higher than 1000  $\mu\text{M}$ , the coleoptile elongation and fresh weight of seedlings were only inhibited by 3000  $\mu\text{M}$  JA (Table 2).

**Table 2.** Final percentages of seed germination and coleoptile emergence, radicle and coleoptile lengths, and fresh weight of the seedlings of *Hordeum vulgare* in various concentrations of JA

| JA concentrations (µM) | Germination percentage (%) | Coleoptile emergence (%) | Radicle length (mm)    | Coleoptile length (mm)  | Fresh weight (mg/seedling) |
|------------------------|----------------------------|--------------------------|------------------------|-------------------------|----------------------------|
| C                      | *100±0,0 <sup>b</sup>      | 100±0,0 <sup>c</sup>     | 121,5±0,8 <sup>j</sup> | 99,8±0,8 <sup>l</sup>   | 258,0±1,2 <sup>m</sup>     |
| 1                      | 100±0,0 <sup>b</sup>       | 100±0,0 <sup>c</sup>     | 97,9±0,2 <sup>i</sup>  | 95,5±2,1 <sup>k</sup>   | 258,0±1,6 <sup>m</sup>     |
| 3                      | 100±0,0 <sup>b</sup>       | 100±0,0 <sup>c</sup>     | 96,9±1,6 <sup>i</sup>  | 92,8±1,6 <sup>j</sup>   | 254,4±0,8 <sup>l</sup>     |
| 5                      | 100±0,0 <sup>b</sup>       | 100±0,0 <sup>c</sup>     | 92,6±1,2 <sup>i</sup>  | 91,3±0,8 <sup>j</sup>   | 245,6±2,1 <sup>k</sup>     |
| 10                     | 100±0,0 <sup>b</sup>       | 100±0,0 <sup>c</sup>     | 92,8±1,6 <sup>i</sup>  | 91,2±1,2 <sup>j</sup>   | 240,4±0,6 <sup>j</sup>     |
| 20                     | 100±0,0 <sup>b</sup>       | 100±0,0 <sup>c</sup>     | 83,9±1,4 <sup>h</sup>  | 82,9±1,2 <sup>i</sup>   | 239,0±1,6 <sup>j</sup>     |
| 30                     | 100±0,0 <sup>b</sup>       | 100±0,0 <sup>c</sup>     | 68,6±1,6 <sup>g</sup>  | 80,0±0,2 <sup>i</sup>   | 226,8±0,2 <sup>i</sup>     |
| 40                     | 100±0,0 <sup>b</sup>       | 100±0,0 <sup>c</sup>     | 68,5±0,8 <sup>g</sup>  | 77,5±0,8 <sup>h</sup>   | 221,2±0,4 <sup>i</sup>     |
| 50                     | 100±0,0 <sup>b</sup>       | 100±0,0 <sup>c</sup>     | 67,8±0,2 <sup>g</sup>  | 76,6±1,4 <sup>h</sup>   | 217,6±1,8 <sup>h</sup>     |
| 100                    | 100±0,0 <sup>b</sup>       | 100±0,0 <sup>c</sup>     | 64,0±2,1 <sup>f</sup>  | 76,5±0,6 <sup>h</sup>   | 216,8±2,1 <sup>h</sup>     |
| 200                    | 100±0,0 <sup>b</sup>       | 100±0,0 <sup>c</sup>     | 51,3±1,6 <sup>e</sup>  | 63,3±1,8 <sup>g</sup>   | 208,8±1,2 <sup>g</sup>     |
| 400                    | 100±0,0 <sup>b</sup>       | 100±0,0 <sup>c</sup>     | 34,0±1,2 <sup>d</sup>  | 51,3±0,2 <sup>f</sup>   | 194,2±0,6 <sup>f</sup>     |
| 500                    | 100±0,0 <sup>b</sup>       | 100±0,0 <sup>c</sup>     | 31,0±0,8 <sup>c</sup>  | 45,6±2,1 <sup>e</sup>   | 185,5±0,8 <sup>e</sup>     |
| 1000                   | 100±0,0 <sup>b</sup>       | 100±0,0 <sup>c</sup>     | 14,0±1,6 <sup>b</sup>  | 34,6±0,8 <sup>d</sup>   | 126,3±1,6 <sup>d</sup>     |
| 1500                   | **0,0±0,0 <sup>a</sup>     | **100±0,0 <sup>c</sup>   | **0,0±0,0 <sup>a</sup> | **19,2±0,6 <sup>c</sup> | **82,0±0,8 <sup>c</sup>    |
| 2000                   | **0,0±0,0 <sup>a</sup>     | **74±2,3 <sup>b</sup>    | **0,0±0,0 <sup>a</sup> | **13,2±0,2 <sup>c</sup> | **58,3±1,2 <sup>b</sup>    |
| 3000                   | 0,0±0,0 <sup>a</sup>       | 0,0±0,0 <sup>a</sup>     | 0,0±0,0 <sup>a</sup>   | 0,0±0,0 <sup>a</sup>    | 0,0±0,0 <sup>a</sup>       |

\* Shows values with insignificant difference (< 0.05) for each column shown with same letters.

\*\* The seedlings from atypically germinated seeds.

Additionally, these severe inhibitions on the germination and seedling growth were irreversible because they were not relieved by washing the seeds with water after the treatments with 1500, 2000 and 3000 µM JA for 7 days (Table 3).

**Table 3.** Final percentages of germination and coleoptile emergence, radicle and coleoptile lengths, and fresh weights of *Hordeum vulgare* after transfer from JA to distilled water

| JA concentrations (µM) | Germination percentage (%) | Coleoptile emergence (%) | Radicle length (mm) | Coleoptile length (mm) | Fresh weight (mg/seedling) |
|------------------------|----------------------------|--------------------------|---------------------|------------------------|----------------------------|
| C                      | 100±0,0                    | 100±0,0                  | 121,5±0,8           | 99,8±0,8               | 258,0±1,2                  |
| 1500                   | *0,0±0,0                   | *100±0,0                 | *0,0±0,0            | *38,5±1,8              | *101,0±0,8                 |
| 2000                   | *0,0±0,0                   | *74±2,3                  | *0,0±0,0            | *20,2±1,2              | *96,4±1,8                  |
| 3000                   | 0,0±0,0                    | 0,0±0,0                  | 0,0±0,0             | 0,0±0,0                | 0,0±0,0                    |

\* The seedlings from atypically germinated seeds.

## DISCUSSION

In this study, effect of JA on seed germination and seedling growth of barley was investigated. Our results showed that JA concentrations under 500 µM did not regard the time courses of the germination of barley seeds, while higher JA levels delayed the germination (Table 1). However, ASGHARI & ISHIZAWA (1998) pointed out that JA-

Me at concentrations over 1  $\mu$ M delayed the time courses of the germination of cocklebur seeds.

JAs have been reported to inhibit the seed germination of several plants such as *Amaranthus* (KEPCZYNSKI et al. 1999, BIALECKA & KEPCZYNSKI 2003), *Agrostemma* (SEMBDNER & GROSS 1986), *Lactuca* (YAMANE et al. 1981), *Helianthus* (CORBINEAU et al. 1988) and *Linum* (WILEN et al. 1991). In all of these cases, high concentrations of JA or JA-Me are necessary to inhibit germination. This results are consistent with our findings. Besides, we determined that 1500 and 2000  $\mu$ M JA levels cause atypical germination (Table 2). However, SEMBDNER & PARTHIER (1993) and ASGHARI & ISHIZAWA (1998) pointed out that a physiological effect of JAs on seed germination is doubtful. In the case of cocklebur seeds, significant inhibition of the germination was detected even at 1  $\mu$ M Me-JA.

On the other hand, we observed that the seedling growth slowed down with increasing concentrations of JA (Table 2). Similarly, KODA et al. (1991), PARTHIER (1991), STASWICK et al. (1992) and TSAI et al. (1997) reported that JAs inhibited root and shoot elongation and fresh weight at some plant species. Furthermore BAZAKANA et al. (1999) and ASGHARI & ISHIZAWA (1998) pointed out that inhibitory effect of JA on germination of *Dioscorea alata* and cocklebur seeds was reversible. Contrarily, TSAI et al. (1997) determined that inhibitory effect of JA-Me on seedling growth of rice was irreversible. Our results showed that this inhibition on both the germination and seedling growth was irreversible (Table 3). Consequently, JA may affect seed germination and seedling growth in different ways depending on plant taxon, applied concentration and treatment styles.

In summary, these results show clearly that JA has dual effect on germination of barley seeds. At low concentrations, JA does not inhibit the germination and at high concentrations JA inhibits the germination. JAs may inhibit the growth and development by reducing nucleic acid and protein synthesis (ANANIEVA & ANANIEV 1997), or by inhibiting cell division (SWIATEK et al. 2002) and hydrolytic activity (BIALECKA & KEPCZYNSKI 2003). JA may also be effective on the growth and development by causing the reduction of endogenous amounts of stimulating hormones (ASGHARI & ISHIZAWA 1998).

As seen above, effects of JA on seed germination and seedling growth are insufficiently known, hence there is not a consensus. To clarify the effect mechanism(s) dealt chemical further research, fine and detailed, is needed and we believe that our study will contribute to future studies.

## REFERENCES

- ABELES FB, HERSHBERGER WL, DUNN LJ, 1989. Hormonal regulation, and intracellular localization of a 33-kD cationic peroxidase in excised cucumber cotyledons. *Plant Physiology*, 89, 664-668.

- ANANIEVA K, ANANIEV ED, 1997. Comparative study of the effects of methyl jasmonate and abscisic acid on RNA and protein synthesis in excised cotyledons of *Cucurbita pepo* L. (Zucchini). *Bulgarian Journal of Plant Physiology*, 23, 80-90.
- ANDERSON JM, SPILATRO SR, KLAUER SF, FRANCESCHI VR, 1989. Jasmonic acid-dependent increase in the level of vegetative storage proteins in soybean. *Plant Science*, 62, 45-52.
- ASGHARI MN, ISHIZAWA K, 1998. Inhibitory effects of methyl jasmonate on the germination and ethylene production in cocklebur seeds. *Journal of Plant Growth Regulation*, 17, 13-18.
- BAZABAKANA R, FAUCONNIER ML, DIALLO B, DUPONT JP, HOMES J, JAZIRI M, 1999. Control of *Dioscorea alata* microtuber dormancy and germination by jasmonic acid. *Plant Growth Regulation*, 27, 113-117.
- BIALECKA B, KEPCZYNSKI J, 2003. Regulation of  $\alpha$ -amylase activity in *Amaranthus caudatus* seeds by methyl jasmonate, gibberellin A<sub>3</sub>, benzyladenine and ethylene. *Plant Growth Regulation*, 39, 51-56.
- CORBINEAU F, RUDNICKI RM, COME D, 1988. The effect of methyl jasmonate on sunflower (*Helianthus annuus* L.) seed germination and seedling development. *Plant Growth Regulation*, 7, 157-169.
- CROSS BE, WEBSTER GRB, 1970. New metabolites of *Gibberella fujikuroi*. Part XV, N-jasmonoyl- and N-dihydrojasmonoyl-isoleucin. *Journal of the Chemical Society Commun*, 1839-1842.
- DALETSKAYA TV, SEMBDNER G, 1989. Effect of jasmonic acid on germination of nondormant and dormant seeds. *Fiziologija Rastenij*, 36, 1118-1123.
- DATHE W, RONSCH H, PREISS A, SCHADE W, SEMBDNER G, SCHREIBER K, 1981. Endogenous plant hormones of the broad bean, *Vicia faba* L. (-) jasmonic acid, a plant growth inhibitor in pericarp. *Planta*, 153, 530-535.
- FERNANDEZ-MACULET J, YANG SF, 1992. Extraction and partial characterisation of the ethylene-forming enzyme from apple fruit. *Plant Physiology*, 99, 751-754.
- KEPCZYNSKI J, BIALECKA B, 1994. Stimulatory effect of ethephon, ACC, gibberellin A<sub>3</sub> and A<sub>4+7</sub> on germination of methyl jasmonate inhibited *Amaranthus caudatus* L. seeds. *Plant Growth Regulation*, 14, 211-216.
- KEPCZYNSKI J, BIALECKA B, KEPCZYNSKA E, 1999. Ethylene biosynthesis in *Amaranthus caudatus* seeds in response to methyl jasmonate. *Plant Growth Regulation*, 28, 59-65.
- KODA Y, KIKUTA Y, 1991. Possible involvement of jasmonic acid in tuberization of yam plants. *Plant Cell Physiology*, 32, 629-633.
- KODA Y, YOSHIDA K, KIKUTA Y, 1991. Evidence for the involvement of jasmonic acid in the control of the stem-growth habit of soybean plants. *Physiologia Plantarum*, 83, 22-26.
- KRUPINA MV, DATHE W, 1991. Occurrence of jasmonic acid in the red alga *Gelidium latifolium* Z. *Naturforsch*, 46, 1127-1129.
- MASLENKOVA LT, ZANEV Y, POPOVA LP, 1990. Oxygen-evolving activity of thylakoids from barley plants cultivated on different concentrations of jasmonic acid. *Plant Physiology*, 93, 1316-1320.
- MEYER A, MIERSCH O, BUTTNER C, DATHE W, SEMBDNER G, 1984. Occurrence of the plant growth regulator jasmonic acid in plants. *Journal of Plant Growth Regulation*, 3, 1-8.

- PARTHIER B, 1991. Jasmonates, new regulators of plant growth and development: many factors and few hypotheses on their actions. *Botanical Acta*, 104, 446-454.
- POPOVA LP, TSONEV TD, VAKLINOVA SG, 1988. Changes in some photosynthetic and photorespiratory properties in barley leaves after treatment with jasmonic acid. *Journal of Plant Physiology*, 132, 257-261.
- RANJAN R, LEWAK S, 1992. Jasmonic acid promotes germination and lipase activity in non-stratified apple embryos. *Physiologia Plantarum*, 86, 335-339.
- RANJAN R, LEWAK S, 1995. Interaction of jasmonic and abscisic acid in the control of lipases and proteases in germinating apple embryos. *Physiologia Plantarum*, 93, 421-426.
- SEMBDNER G, GROSS D, 1986. Plant Growth Substances of Plant and Microbial Origin. In: BOPP M. (Ed.) *Plant Growth Substances*. Springer-Verlag, Berlin, pp. 139-147.
- SEMBDNER G, PARTHIER B, 1993. The biochemistry and the physiological and molecular actions of jasmonates. *Annual Review Plant Physiology and Plant Molecular Biology*, 44, 569-589.
- STASWICK PE, SU W, HOWELL SH, 1992. Methyl jasmonate inhibition of root growth and induction of a leaf protein are decreased in an *Arabidopsis thaliana* mutant. *Proceedings of the National Academy Sciences USA*, 89, 6837-6840.
- SWIATEK A, LENJOU M, BOCKSTAELE DV, INZE D, ONCKELEN HV, 2002. Differential effects of jasmonic acid and abscisic acid on cell cycle progression in Tobacco BY-2 Cell. *Plant Physiology*, 128, 201-211.
- TSAI FY, LIN CC, KAO CH, 1997. A comparative study of the effects of abscisic acid and methyl jasmonate on seedling growth of rice. *Plant Growth Regulation*, 21, 37-42.
- UEDA J, KATO J, 1982. Inhibition of cytokinin-induced plant growth by jasmonic acid and its methyl ester. *Physiologia Plantarum*, 54, 249-254.
- UEDA J, KATO J, YAMANE H, TAKAHASHI N, 1981. Inhibitory effect of methyl jasmonate and its related compounds on kinetin induced retardation of oat leaf senescence. *Physiologia Plantarum*, 52, 305-309.
- WILEN RW, VAN ROOIJEN JH, PEARCE DW, PHARIS RP, HOLBROOK LA, MOLONEY MM, 1991. Effects of jasmonic acid on embryo specific processes in *Brassica* and *Linum* oilseeds. *Plant Physiology*, 399-405.
- YAMANE H, TAKAGI H, ABE T, YOKATA T, TAKAHASHI N, 1981. Identification of jasmonic acid in three species of higher plants and its biological activities. *Plant Cell Physiology*, 22, 689-697.