PAPER DETAILS

TITLE: Comparison of Biliary Stent and Nasobiliary Drain in Patients with Biliary Leak After

Cholecystectomy

AUTHORS: Ibrahim Ethem Güven

PAGES: 75-76

ORIGINAL PDF URL: https://dergipark.org.tr/tr/download/article-file/4321996



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License



Comparison of Biliary Stent and Nasobiliary Drain in Patients with Biliary Leak After Cholecystectomy

Editöre Mektup: Kolesistektomi Sonrası Biliyer Kaçak Gelişen Hastalarda Biliyer Stent ve Nazobiliyer Drenin Karşılaştırılması

İbrahim Ethem GÜVEN 🧧

Yenimahalle Training and Research Hospital, Department of Gastroenterology, Ankara, Türkiye

Dear Editor,

We are writing to express our appreciation for the recent publication in the Balıkesir Medical Journal Research titled "Comparison of Biliary Stent and Nasobiliary Drain in Patients with Biliary Leak After Cholecystectomy" by Ayte et al (Ayte and Yüksel, 2024). This study is valuable in terms of comparing the two main modalities used in the treatment of biliary leakage.

In the management of the biliary leakage, nasobiliary tubes and biliary stent placement are widely used. The goal of the both endoscopic treatment is to reduce intra-biliary pressure and allow for fistula closure (Pioche and Ponchon, 2013). Among these modalities, the advantage of nasobiliary drainage is that it allows for frequent contrast evaluation to monitor the efficiency of the treatment and to determine the exact duration of therapy required. However, the nasobiliary stent is a serious discomfort for patients, since the distal end of the nasobiliary tube is in the nasal cavity. In addition, patients may experience difficulties with feeding (Elmi and Silverman, 2005).

In this research, authors reported that both biliary stent and nasobiliary drainage were effective methods in the treatment of biliary leakage. However, in the nasobiliary drainage group, the length of hospital stay was longer. In addition, in the biliary stent group, the rate of ERCP related complication was lower. In the light of these findings, biliary stents can be safely used as an alternative to nasobiliary drainage in the treatment of biliary leakage due to their equal efficacy, less complication rate and hospitalization length. While the findings of this study are encouraging, the limited patient population from a single center indicates that further studies with larger cohorts would provide more generalizable data. Overall, we congratulate the authors for their contribution to advancing knowledge on the management of biliary leak.

Corresponding Author: İbrahim Ethem Güven e-mail: drethemg@gmail.com Received: 28 October 2024 Accepted: 1 January 2025 DOI: 10.33716/bmedj.1575211 Güven

REFERENCES

- Ayte, M. R., and Yüksel, İ. (2024). Comparison of Biliary Stent and Nasobiliary Drain in Patients with Biliary Leak After Cholecystectomy. *Balikesir Medical Journal*, 7(3), 95-103. https://doi.org/10.33716/bmedj.1403519
- Pioche, M., and Ponchon, T. (2013). Management of bile duct leaks. *Journal of Visceral Surgery*, 150(3 Suppl), 33-38. https://doi.org/10.1016/j.jviscsurg.2013.05.004
- Elmi, F., & Silverman, W. B. (2005). Nasobiliary tube management of postcholecystectomy bile leaks. *Journal of clinical gastroenterology*, 39(5), 441-444.

https://doi.org/10.1097/01.mcg.0000159215.6266 6.d6