

PAPER DETAILS

TITLE: SOME NOVEL FRACTIONAL INTEGRAL INEQUALITIES FOR DIFFERENT KINDS OF CONVEX FUNCTIONS

AUTHORS: Sinan ASLAN

PAGES: 27-32

ORIGINAL PDF URL: <https://dergipark.org.tr/tr/download/article-file/3194174>

Some Novel Fractional Integral Inequalities for Different Kinds of Convex Functions

Sinan ASLAN^{1,2*}

¹ Ağrı Türk Telekom Social Sciences High School, Ağrı Türkiye

² Ağrı İbrahim Çeçen University, Institute of Graduate Studies, Ağrı Türkiye

sinanaslan0407@gmail.com (ORCID: 0000-0001-5970-1926)

Abstract

In this paper, some novel integral inequalities for different kinds of convex functions have been proved by using Caputo-Fabrizio fractional integral operators. The findings includes several new integral inequalities h -convex functions, s -convex functions in the second sense. We have used the properties of Caputo-Fabrizio fractional operator, definitions of different kinds of convex functions and elementary analysis methods.

Keywords: Caputo-Fabrizio fractional integral operator, h -convex functions, s -convex functions.

1. Introduction

Inequality theory is a field in which many researchers work, with new findings that can be given applications in many disciplines such as mathematical analysis, statistics, approximation theory and numerical analysis together with convex functions. Although the concept of convex function is a concept intertwined with inequalities by definition, it has also formed the main motivation of many researches with its aesthetic structure, features and different types. Let's start with the definition of this important class of functions.

2. Materials and Methods

Definition 2.1. Let I be an interval in R . Then $f: I \rightarrow R$ is said to be convex, if

$$f(tx + (1-t)y) \leq tf(x) + (1-t)f(y)$$

holds for all $x, y \in I$ and $t \in [0, 1]$ (Pečarić et al. 1992).

Definition 2.2. A function $f: R^+ \rightarrow R$, where $R^+ = [0, \infty)$, is said to be s -convex in the second sense if

$$f(\alpha x + \beta y) \leq \alpha^s f(x) + \beta^s f(y)$$

for all $x, y \in [0, \infty)$, $\alpha, \beta \geq 0$ with $\alpha + \beta = 1$ and for some fixed $s \in (0, 1]$. We denote by K_s^2 the class of all s -convex functions (Breckner 1978).

Definition 2.3. (Varosanec 2007) Let $h: J \subseteq R \rightarrow R$ be a non-negative function. We say that $f: I \subseteq R \rightarrow R$ is an h -convex function or that f belongs to the class $SX(h, I)$, if f is non-negative and for all $x, y \in I$ and $\alpha \in [0, 1]$, we have

$$f(\alpha x + (1-\alpha)y) \leq h(\alpha)f(x) + h(1-\alpha)f(y).$$

Definition 2.4. Let $f \in H^1(0, b)$, $b > a$, $\alpha \in [0, 1]$ then, the definition of the left and right side of Caputo-Fabrizio fractional integral is:

$$({}^{CF}_a I^\alpha)(t) = \frac{1-\alpha}{B(\alpha)} f(t) + \frac{\alpha}{B(\alpha)} \int_a^t f(y) dy,$$

and

$$({}^{CF}_b I^\alpha)(t) = \frac{1-\alpha}{B(\alpha)} f(t) + \frac{\alpha}{B(\alpha)} \int_t^b f(y) dy$$

where $B(\alpha) > 0$ is normalization function (Abdeljawad and Baleanu 2017).

In the sequel of the paper, we will denote normalization function as $B(\alpha)$ with $B(0) = B(1) = 1$.

In (Tariq et al. 2022), the authors provided an integral inequality of Hermite-Hadamard type for preinvex functions via Caputo-Fabrizio fractional integral inequality as follows.

Theorem 2.1. Let $f: I = [k_1, k_1 + \mu(k_2, k_1)] \rightarrow (0, \infty)$ be a preinvex function on I° and $f \in$

Received: 07/06/2023

Accepted: 27/06/2023

Published: 30/06/2023

*Corresponding author: Sinan ASLAN, PhD

Ağrı İbrahim Çeçen University, Institute of Graduate Studies, Ağrı Türkiye

E-mail: sinanaslan0407@gmail.com

Cite this article as: S. ASLAN, Some Novel Fractional Integral Inequalities for Different Kinds of Convex Functions, Eastern Anatolian Journal of Science, Vol. 9, Issue 1, 27-32, 2023.

$L[k_1, k_1 + \mu(k_2, k_1)]$. If $\alpha \in [0, 1]$, then the following inequality holds

$$\begin{aligned} & f\left(\frac{2k_1 + \mu(k_2, k_1)}{2}\right) \\ & \leq \frac{B(\alpha)}{\alpha\mu(k_2, k_1)} \\ & \times \left[{}^{CF}I_{k_1}^\alpha \{f(k)\} + {}^{CF}I_{k_1+\mu(k_2, k_1)}^\alpha \{f(k)\} \right. \\ & \quad \left. - \frac{2(1-\alpha)}{B(\alpha)} f(k) \right] \\ & \leq \frac{f(k_1) + f(k_2)}{2} \end{aligned}$$

where $k \in [k_1, k_1 + \mu(k_2, k_1)]$.

Atangana and Baleanu produced a new derivative operators using Mittag-Leffler function in Caputo-Fabrizio derivative operator as following.

Definition 2.5. (Atangana and Baleanu 2016). Let $f \in H^1(0, b)$, $b > a$, $\alpha \in [0, 1]$ then, the definition of the new fractional derivative is given:

$$(1.1) \quad ({}^{ABC}_a D_t^\alpha)[f(t)] = \frac{B(\alpha)}{1-\alpha} \int_a^t f'(x) E_\alpha \left[-\alpha \frac{(t-x)^\alpha}{(1-\alpha)} \right] dx.$$

Definition 2.6. (Atangana and Baleanu 2016). Let $f \in H^1(0, b)$, $b > a$, $\alpha \in [0, 1]$ then, the definition of the new fractional derivative is given:

$$(1.2) \quad ({}^{ABR}_a D_t^\alpha)[f(t)] = \frac{B(\alpha)}{1-\alpha} \frac{d}{dt} \int_a^t f(x) E_\alpha \left[-\alpha \frac{(t-x)^\alpha}{(1-\alpha)} \right] dx.$$

Equations (1.1) and (1.2) have a non-local kernel. Also in equation (1.1) when the function is constant we get zero.

The related fractional integral operator has been defined by Atangana-Baleanu as follows.

Definition 2.7. The fractional integral associate to the new fractional derivative with non-local kernel of a function $f \in H^1(a, b)$ as defined:

$$\begin{aligned} {}^{AB}_a I^\alpha \{f(t)\} &= \frac{1-\alpha}{B(\alpha)} f(t) \\ &+ \frac{\alpha}{B(\alpha)\Gamma(\alpha)} \int_a^t f(y)(t-y)^{\alpha-1} dy \end{aligned}$$

where, $b > a$, $\alpha \in [0, 1]$ (Atangana and Baleanu 2016).

Abdeljawad and Baleanu introduced right hand side of integral operator as following; The right

fractional new integral with ML kernel of order $\alpha \in [0, 1]$ is defined by

$$\begin{aligned} {}^{AB}_b I_b^\alpha \{f(t)\} &= \frac{1-\alpha}{B(\alpha)} f(t) \\ &+ \frac{\alpha}{B(\alpha)\Gamma(\alpha)} \int_t^b f(y)(y-t)^{\alpha-1} dy. \end{aligned}$$

where, $b > a$, $\alpha \in [0, 1]$ (Abdeljawad and Baleanu 2017).

For more information related to different kinds of fractional operators, we recommend to the readers the following papers (Abdeljawad 2015, Abdeljawad and Baleanu 2016, Akdemir et al. 2021- Akdemir et al. 2017, Butt et al. 2020, Caputo and Fabrizio 2015- Gürbüz et al. 2020, Rashid et al 2020-Samko et al. 1993, Set 2012, Set et al. 2017).

3. Results

Theorem 3.1. Let $I \subseteq \mathbb{R}$. Suppose that $f: [a, b] \subseteq I \rightarrow \mathbb{R}$ is a h -convex function on $[a, b]$ such that $f \in L_1[a, b]$. Then, we have following inequality for Caputo-Fabrizio fractional integrals:

$$\begin{aligned} & ({}^{CF}_a I_a^\alpha f)(k) + ({}^{CF}_b I_b^\alpha f)(k) \\ & \leq \frac{2(1-\alpha)}{B(\alpha)} f(k) + \frac{\alpha(b-a)f(a)}{B(\alpha)} \int_0^1 h(t) dt \\ & \quad + \frac{\alpha(b-a)f(b)}{B(\alpha)} \int_0^1 h(1-t) dt \end{aligned}$$

where $B(\alpha) > 0$ is normalization function and $\alpha \in [0, 1]$.

Proof. By using the definition of h -convex function, we can write

$$f(ta + (1-t)b) \leq h(t)f(a) + h(1-t)f(b).$$

By integrating both sides of the inequality over $[0, 1]$ with respect to t , we get

$$\begin{aligned} & \int_0^1 f(ta + (1-t)b) dt \\ & \leq f(a) \int_0^1 h(t) dt \\ & \quad + f(b) \int_0^1 h(1-t) dt. \end{aligned}$$

By changing of the variable as $x = ta + (1-t)b$, we obtain

$$\begin{aligned} & \frac{1}{b-a} \int_a^b f(x) dx \leq f(a) \int_0^1 h(t) dt \\ & \quad + f(b) \int_0^1 h(1-t) dt. \end{aligned}$$

By multiplying both sides of the above inequality with $\frac{\alpha(b-a)}{B(\alpha)}$ and adding $\frac{2(1-\alpha)}{B(\alpha)} f(k)$, we have

$$\begin{aligned} & \frac{2(1-\alpha)}{B(\alpha)} f(k) + \frac{\alpha}{B(\alpha)} \int_a^b f(x) dx \\ & \leq \frac{2(1-\alpha)}{B(\alpha)} f(k) \\ & \quad + \frac{\alpha(b-a)f(a)}{B(\alpha)} \int_0^1 h(t) dt \\ & \quad + \frac{\alpha(b-a)f(b)}{B(\alpha)} \int_0^1 h(1-t) dt. \end{aligned}$$

By simplifying the inequality, we get the result

$$\begin{aligned} & \left(\frac{1-\alpha}{B(\alpha)} f(k) + \frac{\alpha}{B(\alpha)} \int_a^k f(x) dx \right) \\ & + \left(\frac{1-\alpha}{B(\alpha)} f(k) + \frac{\alpha}{B(\alpha)} \int_k^b f(x) dx \right) \\ & \leq \frac{2(1-\alpha)}{B(\alpha)} f(k) \\ & \quad + \frac{\alpha(b-a)f(a)}{B(\alpha)} \int_0^1 h(t) dt \\ & \quad + \frac{\alpha(b-a)f(b)}{B(\alpha)} \int_0^1 h(1-t) dt \end{aligned}$$

Namely,

$$\begin{aligned} & ({}^{CF}I_a^\alpha f)(k) + ({}^{CF}I_b^\alpha f)(k) \\ & \leq \frac{2(1-\alpha)}{B(\alpha)} f(k) + \frac{\alpha(b-a)f(a)}{B(\alpha)} \int_0^1 h(t) dt \\ & \quad + \frac{\alpha(b-a)f(b)}{B(\alpha)} \int_0^1 h(1-t) dt. \end{aligned}$$

This completes the proof.

Theorem 3.2. Let $I \subseteq \mathbb{R}$. Suppose that $f: [a, b] \subseteq I \rightarrow \mathbb{R}$ is a s -convex function in the second sense on $[a, b]$ such that $f \in L_1[a, b]$. Then, we have the following inequality for Caputo-Fabrizio fractional integrals:

$$\begin{aligned} & ({}^{CF}I_a^\alpha f)(k) + ({}^{CF}I_b^\alpha f)(k) \\ & \leq \frac{2(1-\alpha)f(k)(s+1) + \alpha(b-a)(f(a) + f(b))}{B(\alpha)(s+1)}. \end{aligned}$$

where $B(\alpha) > 0$ is normalization function $s \in (0, 1]$ and $\alpha \in [0, 1]$.

Proof. By using the definition of s -convex function in the second sense, we can write

$$f(ta + (1-t)b) \leq t^s f(a) + (1-t)^s f(b).$$

By integrating both sides of the inequality over $[0, 1]$ with respect to t , we get

$$\begin{aligned} & \int_0^1 f(ta + (1-t)b) dt \\ & \leq \int_0^1 t^s f(a) dt \\ & \quad + \int_0^1 (1-t)^s f(b) dt. \end{aligned}$$

By changing of the variable as $x = ta + (1-t)b$, and by calculating the right hand side, we obtain

$$\frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{s+1}.$$

By multiplying both sides of the above inequality with $\frac{\alpha(b-a)}{B(\alpha)}$ and adding $\frac{2(1-\alpha)}{B(\alpha)} f(k)$, we have

$$\begin{aligned} & \frac{2(1-\alpha)}{B(\alpha)} f(k) + \frac{\alpha}{B(\alpha)} \int_a^b f(x) dx \\ & \leq \frac{2(1-\alpha)}{B(\alpha)} f(k) \\ & \quad + \frac{\alpha(b-a)f(a) + f(b)}{B(\alpha)(s+1)}. \end{aligned}$$

By simplifying the inequality, we get the result.

$$\begin{aligned} & \left(\frac{1-\alpha}{B(\alpha)} f(k) + \frac{\alpha}{B(\alpha)} \int_a^k f(x) dx \right) \\ & + \left(\frac{1-\alpha}{B(\alpha)} f(k) + \frac{\alpha}{B(\alpha)} \int_k^b f(x) dx \right) \\ & \leq \frac{2(1-\alpha)}{B(\alpha)} f(k) \\ & \quad + \frac{\alpha(b-a)f(a) + f(b)}{B(\alpha)(s+1)} \end{aligned}$$

Then, it is easy to see

$$\begin{aligned} & ({}^{CF}I_a^\alpha f)(k) + ({}^{CF}I_b^\alpha f)(k) \\ & \leq \frac{2(1-\alpha)f(k)(s+1) + \alpha(b-a)(f(a) + f(b))}{B(\alpha)(s+1)}. \end{aligned}$$

This completes the proof

Theorem 3.3. Let $I \subseteq \mathbb{R}$. Suppose that $f: [a, b] \subseteq I \rightarrow \mathbb{R}$ is a s -convex function in the second sense on $[a, b]$ such that $f \in L_1[a, b]$. Then, we have the following inequality for Caputo-Fabrizio fractional integrals:

$$\begin{aligned} & ({}^{CF}I_a^\alpha f)(k) + ({}^{CF}I_b^\alpha f)(k) \\ & \leq \frac{2(1-\alpha)f(k)(ps+1)^{\frac{1}{p}} + \alpha(b-a)(f(a) + f(b))}{B(\alpha)(ps+1)^{\frac{1}{p}}} \end{aligned}$$

where $B(\alpha) > 0$ is normalization function $s \in (0, 1]$, $q > 1$, $\frac{1}{p} + \frac{1}{q} = 1$ and $\alpha \in [0, 1]$.

Proof : By using the definition of s -convex function in the second sense, we can write

$$f(ta + (1-t)b) \leq t^s f(a) + (1-t)^s f(b).$$

By integrating both sides of the inequality over $[0, 1]$ with respect to t , we get

$$\begin{aligned} & \int_0^1 f(ta + (1-t)b) dt \\ & \leq f(a) \int_0^1 t^s dt \\ & \quad + f(b) \int_0^1 (1-t)^s dt. \end{aligned}$$

If we apply the Hölder's inequality to the right-hand side of the inequality, we get

$$\begin{aligned} & \int_0^1 |f(ta + (1-t)b)| dt \\ & \leq f(a) \left(\left(\int_0^1 t^{ps} dt \right)^{\frac{1}{p}} \left(\int_0^1 1^q dt \right)^{\frac{1}{q}} \right) \\ & \quad + f(b) \left(\left(\int_0^1 (1-t)^{ps} dt \right)^{\frac{1}{p}} \left(\int_0^1 1^q dt \right)^{\frac{1}{q}} \right) \\ & = f(a) \left(\left(\frac{1}{ps+1} \right)^{\frac{1}{p}} (1^q)^{\frac{1}{q}} \right) \\ & \quad + f(b) \left(\left(\frac{1}{ps+1} \right)^{\frac{1}{p}} (1^q)^{\frac{1}{q}} \right). \end{aligned}$$

By changing of the variable as $x = ta + (1-t)b$ and by calculating the right hand side, we obtain

$$\frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{(ps+1)^{\frac{1}{p}}}.$$

By multiplying both sides of the above inequality with $\frac{\alpha(b-a)}{B(\alpha)}$ and adding $\frac{2(1-\alpha)}{B(\alpha)} f(k)$, we have

$$\begin{aligned} & \frac{2(1-\alpha)}{B(\alpha)} f(k) + \frac{\alpha}{B(\alpha)} \int_a^b f(x) dx \\ & \leq \frac{2(1-\alpha)}{B(\alpha)} f(k) \\ & \quad + \frac{\alpha(b-a)}{B(\alpha)} \frac{f(a) + f(b)}{(ps+1)^{\frac{1}{p}}}. \end{aligned}$$

By simplifying the inequality, we get the result

$$\begin{aligned} & \left(\frac{1-\alpha}{B(\alpha)} f(k) + \frac{\alpha}{B(\alpha)} \int_a^k f(x) dx \right) \\ & + \left(\frac{1-\alpha}{B(\alpha)} f(k) + \frac{\alpha}{B(\alpha)} \int_k^b f(x) dx \right) \\ & \leq \frac{2(1-\alpha)}{B(\alpha)} f(k) \\ & \quad + \frac{\alpha(b-a)}{B(\alpha)} \frac{f(a) + f(b)}{(ps+1)^{\frac{1}{p}}}. \end{aligned}$$

Then, we can conclude that

$$\begin{aligned} & ({}^{CF}I_a^\alpha f)(k) + ({}^{CF}I_b^\alpha f)(k) \\ & \leq \frac{2(1-\alpha)f(k)(ps+1)^{\frac{1}{p}} + \alpha(b-a)(f(a) + f(b))}{B(\alpha)(ps+1)^{\frac{1}{p}}}. \end{aligned}$$

Theorem 3.4. Let $I \subseteq \mathbb{R}$. Suppose that $f: [a, b] \subseteq I \rightarrow \mathbb{R}$ is a s -convex function in the second sense on $[a, b]$ such that $f \in L_1[a, b]$. Then, we have following inequality for Caputo-Fabrizio fractional integrals:

$$\begin{aligned} & ({}^{CF}I_a^\alpha f)(k) + ({}^{CF}I_b^\alpha f)(k) \\ & \leq \frac{2(1-\alpha)}{B(\alpha)} f(k) \\ & \quad + \frac{\alpha(b-a)(f(a) + f(b))(q + p(ps+1))}{B(\alpha)pq(ps+1)} \end{aligned}$$

where $B(\alpha) > 0$ is normalization function $s \in (0, 1]$, $q > 1$, $\frac{1}{p} + \frac{1}{q} = 1$ and $\alpha \in [0, 1]$.

Proof : By using the definition of s -convex function in the second sense, we can write

$$f(ta + (1-t)b) \leq t^s f(a) + (1-t)^s f(b).$$

By integrating both sides of the inequality over $[0, 1]$ with respect to t , we get

$$\begin{aligned} & \int_0^1 f(ta + (1-t)b) dt \\ & \leq \int_0^1 t^s f(a) dt \\ & \quad + \int_0^1 (1-t)^s f(b) dt. \end{aligned}$$

If we apply the Young's inequality to the right-hand side of the inequality, we get

$$\begin{aligned} & \int_0^1 |f(ta + (1-t)b)| dt \\ & \leq f(a) \left(\frac{1}{p} \left(\int_0^1 t^{ps} dt \right) + \frac{1}{q} \left(\int_0^1 1^q dt \right) \right) \\ & \quad + f(b) \left(\frac{1}{p} \left(\int_0^1 (1-t)^{ps} dt \right) + \frac{1}{q} \left(\int_0^1 1^q dt \right) \right) \\ & \leq f(a) \left(\frac{1}{p(ps+1)} + \frac{1}{q} \right) + f(b) \left(\frac{1}{p(ps+1)} + \frac{1}{q} \right). \end{aligned}$$

By changing of the variable as $x = ta + (1-t)b$ and by calculating the right hand side, we obtain

$$\frac{1}{b-a} \int_a^b f(x) dx \leq \frac{(f(a) + f(b))(q + p(ps+1))}{pq(ps+1)}.$$

By multiplying both sides of the above inequality with $\frac{\alpha(b-a)}{B(\alpha)}$ and adding $\frac{2(1-\alpha)}{B(\alpha)} f(k)$, we have

$$\begin{aligned} & \frac{2(1-\alpha)}{B(\alpha)} f(k) + \frac{\alpha}{B(\alpha)} \int_a^b f(x) dx \\ & \leq \frac{2(1-\alpha)}{B(\alpha)} f(k) \\ & + \frac{\alpha(b-a)(f(a)+f(b))(q+p(ps+1))}{B(\alpha)pq(ps+1)}. \end{aligned}$$

By simplifying the inequality, we get the result

$$\begin{aligned} & \left(\frac{1-\alpha}{B(\alpha)} f(k) + \frac{\alpha}{B(\alpha)} \int_a^k f(x) dx \right) \\ & + \left(\frac{1-\alpha}{B(\alpha)} f(k) + \frac{\alpha}{B(\alpha)} \int_k^b f(x) dx \right) \\ & \leq \frac{2(1-\alpha)}{B(\alpha)} f(k) \\ & + \frac{\alpha(b-a)(f(a)+f(b))(q+p(ps+1))}{B(\alpha)pq(ps+1)} \end{aligned}$$

Namely,

$$\begin{aligned} & ({}^{CF}I_a^\alpha f)(k) + ({}^{CF}I_b^\alpha f)(k) \\ & \leq \frac{2(1-\alpha)}{B(\alpha)} f(k) \\ & + \frac{\alpha(b-a)(f(a)+f(b))(q+p(ps+1))}{B(\alpha)pq(ps+1)}. \end{aligned}$$

Remark 1. If we set $s = 1$ in the main findings, we obtain new estimations for convex functions via Caputo-Fabrizio fractional integrals.

References

- ABDELJAWAD, THABET. On conformable fractional calculus. *Journal of computational and Applied Mathematics*, 2015, 279: 57-66.
- ABDELJAWAD, THABET; BALEANU, DUMITRU. Integration by parts and its applications of a new nonlocal fractional derivative with Mittag-Leffler nonsingular kernel. *arXiv preprint arXiv:1607.00262*, 2016.
- ABDELJAWAD, THABET; BALEANU, DUMITRU. On fractional derivatives with exponential kernel and their discrete versions. *Reports on Mathematical Physics*, 2017, 80.1: 11-27.
- AKDEMİR, A. O.; ASLAN, S.; ÇAKALOĞLU, M.N. EKİNCİ, A. Some New Results for Different Kinds of Convex Functions CaputoFabrizio Fractional Operators. 4th International Conference on Mathematical and Related Sciences. Page (92) ICMRS 2021.
- AKDEMİR, A. O.; ASLAN, S.; ÇAKALOĞLU, M.N. SET, E. New Hadamard Type Integral Inequalities via Caputo-Fabrizio Fractional Operators. 4th International Conference on Mathematical and Related Sciences. Page (91) ICMRS 2021.
- AKDEMİR, A. O.; ASLAN, S.; EKİNCİ, A. Novel Approaches for s-Convex Functions via Caputo-Fabrizio Fractional Integrals. *Proceedings of IAM*, 2022, 11.1: 3-16.
- AKDEMİR, AHMET OCAK, et al. New general variants of Chebyshev type inequalities via generalized fractional integral operators. *Mathematics*, 2021, 9.2: 122.
- AKDEMİR, AHMET OCAK; EKİNCİ, ALPER; SET, ERHAN. Conformable fractional integrals and related new integral inequalities. 2017.
- ATANGANA, ABDON; BALEANU, DUMITRU. New fractional derivatives with nonlocal and non-singular kernel: theory and application to heat transfer model. *arXiv preprint arXiv:1602.03408*, 2016.
- BRECKNER, WOLFGANG W. Stetigkeitsaussagen für eine Klasse verallgemeinerter konvexer Funktionen in topologischen linearen Räumen. *Publ. Inst. Math.(Beograd)(NS)*, 1978, 23.37: 13-20.
- BUTT, SAAD IHSAN, et al. New Hermite-Jensen-Mercer-type inequalities via k-fractional integrals. *Advances in Difference Equations*, 2020, 2020: 1-24.
- CAPUTO, MICHELE; FABRIZIO, MAURO. A new definition of fractional derivative without singular kernel. *Progress in Fractional Differentiation & Applications*, 2015, 1.2: 73-85.
- EKİNCİ, ALPER; OZDEMİR, MUHAMET. Some new integral inequalities via Riemann-Liouville integral operators. *Applied and computational mathematics*, 2019, 18.3.
- GÜRBÜZ, MUSTAFA, et al. Hermite-Hadamard inequality for fractional integrals of Caputo-Fabrizio type and related inequalities. *Journal of Inequalities and Applications*, 2020, 2020: 1-10.
- PEAJCARIAAC, JOSIP E.; TONG, YUNG LIANG. *Convex functions, partial orderings, and statistical applications*. Academic Press, 1992.
- RASHID, SAIMA, et al. New investigation on the generalized K-fractional integral operators. *Frontiers in Physics*, 2020, 8: 25.
- RASHID, SAIMA, et al. New multi-parametrized estimates having pth-order differentiability in fractional calculus for predominating h-convex functions in Hilbert space. *Symmetry*, 2020, 12.2: 222.

- SAMKO, STEFAN G., et al. *Fractional integrals and derivatives*. Yverdon-les-Bains, Switzerland: Gordon and Breach Science Publishers, Yverdon, 1993.
- SET, ERHAN. New inequalities of Ostrowski type for mappings whose derivatives are s -convex in the second sense via fractional integrals. *Computers & Mathematics with Applications*, 2012, 63.7: 1147-1154.
- SET, ERHAN; AKDEMİR, AHMET OCAK; ÖZDEMİR, EMİN. M. Simpson type integral inequalities for convex functions via Riemann-Liouville integrals. *Filomat*, 2017, 31.14: 4415-4420.
- TARIQ, MUHAMMAD, et al. New fractional integral inequalities for preinvex functions involving Caputo-Fabrizio operator. 2022.
- VAROŠANEC, SANJA. On h -convexity. *Journal of Mathematical Analysis and Applications*, 2007, 326.1: 303-311.