## PAPER DETAILS

TITLE: Short-term monitoring of the winching and skidding effects on soil microbial biomass in

Turkish red pine in the Mediterranean Region

AUTHORS: Emre BABUR

PAGES: 107-121

ORIGINAL PDF URL: https://dergipark.org.tr/tr/download/article-file/1924909



Emre BABUR\*

\*Department of Forest Engineering, Faculty of Forestry, Kahramanmaraş Sütçü Imam University, 46050, Kahramanmaraş, Turkey. Corresponding Author: <u>emrebabur@ksu.edu.tr</u>

### Abstract

Harvesting activities in forests can seriously damage soils and cause short and long-term changes in some of their properties. The aim of this study is to determine the effects on the microbial properties of the failure of the soil by the whole tree method using a farm tractor in the short term in the Turkish red-pine forests of Kahramanmaraş Forestry Operations Directorate in Başkonuş Forest Enterprise Chief. In total, 72 soil samples were collected on two soil layers (0-10 and 10-20 cm) and three seasons (spring, summer, and autumn) for identifying some physicochemical and microbial properties of soil. Mean values of the soil organic carbon and nitrogen were statistically different in the skidding (2.15% - 0.13%) and control (2.90% - 0.16%) areas, respectively. Also, It was determined that the skidding activities had a statistically significant effect on the microbial biomass carbon, nitrogen, and microbial soil respiration. A significant reduction in organic carbon and microbial biomass was observed in the soils in the skidding line. According to the seasonal patterns, the microbial biomass of the samples was found the lowest in summer (657.17  $\mu$ g g<sup>-1</sup>) and the highest (763.76  $\mu$ g g-1) in autumn. In the control areas, the lowest was 773.99  $\mu$ g g<sup>-1</sup> in the spring season and the highest was 886  $\mu$ g g-1 in the autumn season. It is predicted that the decomposition rate may have increased in parallel with the soil temperature, which increases as a result of the removal of the litter layer from the soil surface to any other places in the harvesting application. Consequently, it is important for forest and soil health to monitor the changes in the microbiological characteristics of soils for long periods and to produce in a way that causes minimum damage to the soil in harvesting activities of forests. Therefore, harvesting activities should be carried out in periods when the soil is hard. In seasons, when soils are soft and sensitive, there is a need to develop alternative harvesting methods such as aerial yarding systems from stump to landing instead of skidding.

Keywords: harvesting activities, carbon, soil properties, microbial biomass, seasonal pattern

## Akdeniz Bölgesinde Kızılçam meşcerelerinde taşıma ve sürütme ile bölmeden çıkartma faaliyeti sonucu toprakların mikrobiyal özelliklerinin değişiminin mevsimsel olarak izlenmesi

### Özet

Ormanlarda yapılan hasat (üretim) faaliyetleri topraklara ciddi şekilde zarar verebilmekte ve bazı özelliklerinde kısa ve uzun vadeli değişikliklere neden olabilmektedir. Bu çalışmanın amacı, tarım traktörü kullanılarak bütün ağaç yöntemi ile hasat yapıldıktan sonra yırtılan (bozulan) toprakların kısa vadede bazı fizikokimyasal ve mikrobiyal özellikleri üzerindeki etkilerini belirlemektir. Toprağın bazı fizikokimyasal ve mikrobiyal özellikleri üzerindeki etkilerini belirlemektir. Toprağın bazı fizikokimyasal ve mikrobiyal özellikleri toprak katmanında (0-10 ve 10-20 cm) ve üç mevsimde (ilkbahar, yaz ve sonbahar) toplam 72 toprak örneği toplanmıştır. Toprak organik karbon ve azot ortalama değerleri sırasıyla örneklerde (%2.15 ve %0.13) ve kontrol (%2.90 ve %0.16) alanlarında istatistiksel olarak farklı bulunmuştur. Ayrıca, üretim uygulamasının mikrobiyal biyokütle karbon, azot ve mikrobiyal toprak solunumu üzerinde istatistiksel olarak önemli etkilere sahip olduğu belirlenmiştir. Özellikle sürütme hattındaki topraklarda organik karbon ve mikrobiyal biyokütlede önemli miktarda azalma gözlenmiştir. Mevsimsel desene göre, toprak örneklerinin mikrobiyal

biyokütlesi en düşük (657.17 µg g<sup>-1</sup>) yaz ve en yüksek (763.76 µg g<sup>-1</sup>) sonbahar mevsiminde bulunmuştur. Kontrol alanlarında en düşük 773.99 µg g-1 ilkbahar mevsiminde, en yüksek 886 µg g-1 sonbahar mevsiminde olmuştur. Üretim faaliyetleri sonucunda ölü örtü tabakasının toprak yüzeyinden uzaklaştırılması sonucu artan toprak sıcaklığı ve azalan toprak nemine paralel olarak toprakların mikrobiyal özelliklerinde azalmalar olduğu gözlemlenmiştir. Sonuç olarak, ormanlarda üretimden sonra toprakların mikrobiyolojik özelliklerindeki değişimlerin uzun vadede izlenmesi, toprağa en az zarar verecek şekilde üretim yapılması orman ve toprak sağlığı için önemlidir. Bu nedenle üretim faaliyetleri toprağın sert olduğu dönemlerde yapılmalıdır. Toprakların yumuşak ve hassas olduğu mevsimlerde bütün ağaç veya sürütme yöntemi yerine alternatif hasat yöntemlerinin geliştirilmesine ihtiyaç vardır.

Anahtar Kelimeler: hasat faaliyetleri, karbon, toprak özellikleri, mikrobiyal biyokütle, mevsimsel farklar

## 1. Introduction

Some physicochemical properties and especially microbiological properties of soils show considerable sensitivity to changes in environmental factors and forestry practices such as harvesting, afforestation, restoration (Foote et al. 2015; Kara et al. 2016; Babur 2019). A significant amount of carbon is stored in the litter and topsoil of forest ecosystems. The amount of stored carbon under the influence of climate change and human affects these pools negatively (Chen et al. 2013; Dangal et al. 2014). Timber harvesting, which is one of the main forestry activities, causes loss a huge amount of organic matter in the forest ecosystem by soil compaction, soil rupture, and erosion. So, this decreases the productivity of forest ecosystems. Unless an appropriate and sustainable harvest planning is made, the fertility of the soils is endangered due to the decrease in the C and N resources of the ecosystem (Henderson 1995; Carter et al. 2002; Turner and Lambert 2011). However, the forest ecosystems can only be able to renew itself in about 100 years (Chen et al. 2013; Kellman et al. 2014). In addition, soil compaction and degradation with rupture caused by harvesting activities increase soil temperature and bulk density, changing the soil structure and porosity, which directly affects the soil aeration and water capacity which extremely related with infiltration capacity (Labelle and Jaeger 2011; Cambi et al. 2015).

Considering that the microbiological population and indices are affected the most against all these changes in forest ecosystems, and microbiological soil properties can be used as indicators about changes in an ecosystem (Wardle 1992; Jordan et al. 2003; Frey et al. 2009).

Harvesting processes made according to different methods cause different effects on forest soils and litter (Jones et al. 2011; Grand and Lavkulich 2012; Kellman et al. 2014; Scott et al. 2014). However, it has not been clearly demonstrated how this method affects carbon and nitrogen amounts in soils and litter, microbial decomposition rates and soil aggregation. Six et al. (2004) stated in their study that there is degradation in the microaggregates of the top soils compaction as a result of harvesting activities. In addition, it has been determined that soil compaction decreases root biomass in the upper soils and N uptake from the soil, N washing and denitrification in the ecosystem (Torbert and Wood 1992; Jordan et al. 2003).

Soil microbial organisms play an important role in C and N storage and nutrient cycle (Haubensak et al. 2002; Allen and Schlesinger 2004). There are positive relationships between soil microbial biomass and C and N stocks of soils (Li et al. 2004; Kara et al. 2016; Babur 2019). LeDuc and Rothstein (2007) reported that the post-production reductions in C and N may result in a reduction in microbial biomass. In addition, it has been stated that microbial activity in the top soils significantly decreased due to the effect of soil compaction on the air and water economy of the soils (Li et al. 2004; Mariani et al. 2006). In this study, the effects of the deteriorated soil structure on some physicochemical and microbial properties in the short term were investigated after the whole tree method was removed from the partition using a farm tractor.

## 2. Materials and Methods

## 2.1. Material

This study was located at 50 km far from northwest of Kahramanmaras province in the Northeast Taurus Mountain region of Turkey (37° 35' 19" N; 36° 36' 13"E) in the Turkish red pine forests of Kahramanmaras Forestry Operations Directorate in Baskonus Forest Enterprise Chief and was carried out in the spring, summer and autumn seasons throughout 2018 (Figure 1). Pure Turkish red pine stand in section 33 to be sampled has been put into production and the trees cut with the whole tree method in this area were transported by sliding uphill using a single drum crane mounted in front of an agricultural tractor in April 2018. During transportation, especially the parts of the log that were separated from the bottom log, destroyed the soil by tearing and removed the litter on it, revealing the mineral soil. In particular, samples were taken along this destroyed line and control samples were taken from undisturbed areas on both sides of this line (Figure 2). According to Köppen-Geiger climate classification (www.koeppen-geiger.vu-wien.ac.at/), the study site is characterized as a "hot and dry summer" environment (Csa). Weather data, including the long-term (1990-2017) normal averages for monthly air temperature (°C) and monthly precipitation (in mm) at Başkonuş in Kahramanmaraş province, Turkey, were obtained from the Turkish State Meteorological Service (www.mgm.gov.tr/). The mean annual temperature is 13.3 °C and annual accumulated average precipitation is 700 mm, with peaks slightly exceeding the 110 mm in December and the minimum is below 10 mm in July and August. Sandstone bedrock is dominant in the area and topography of the study site is highly variable, with an average slope of 30% and elevation 1000 m. Soil particle fractions such as average sand, silt, and clay ratios were found the higher coarse fractions in soil samples. Study site soils were found in sandy clay loam texture class.

Some characteristics for the sites under study, including sand, silt and clay content (%), slope, stand canopy, and others, are presented in Table 1.



Figure 1. Geographical location of the study site in Turkey.



**Figure 2.** Turkish red pine timber extraction by using farm tractor: uphill winching (a), crumbs of soil left on the log (b) and damage to the soil by skidding whole tree (c)

**Table 1.** Location and selected topographic features, soil texture, and pH classes of the study site in the Kahramanmaraş province, Turkey.

Properties	Sample Area	Control Area	
Longitude	36°36'35''	36°36'50''	
Latitude	37°35'25''	37 °35'50''	
Altitude (m)	1000	1000	
Slope (%)	30	30	
Vegetation Types	Çzcd3	Çzcd3	
Canopy (%)	70-100	70-100	
Soil texture	Sandy Clay loam	Sandy clay loam	
pH classes	Slightly alkaline	Moderately alkaline	

## 2.2. Method

## 2.2.1. Experimental design and soil sampling

Disturbed, undisturbed and microbial soil samples were taken from two levels (0-10 cm and 10-20 cm) from the middle of the scuffing pit and the closest undisturbed control areas every 20 m along two

different skidding lines (60 m). In order to monitor the seasonal changes in the skidding line, soil samples were taken in three different seasons (spring, summer and autumn), with regarding to the average temperature and precipitation values of the seasons. Thus, a total of 72 soil samples, 2 lines x 6 points x 2 depth levels x 3 seasons, were collected from the study areas. Disturebed soil samples were taken for some physical and chemical analyses, and undisturbed soil samples (using 8 cm x 10 cm steel cylinders) were taken to determine the bulk density. In addition, microbial soil samples, which were passed through a 2 mm sieve and moistened 50-60% with distilled water, were kept in a +4 °C refrigerator in the laboratory until their analysis.

## 2.2.2 Physical and Chemical Properties of Soils

Following field study, soil samples were oven-dried until constant mass weight in the laboratory (Karaöz 1992). Prior to analysis of soil parameters, air-dried soils were sieved to pass a 2 mm screen. The bulk denisity of the soil samples was calculated as the dry weight (105 °C at 24h) of soil divided by its volume. Soil moisture content was determened by gravimetically. Soil pH and electrical conductivity (EC) were measured in distilled water by using a glass electrode (1:2.5 w/v and 1:5 w/v soil/ water suspension for soil pH and EC, respectively) (Rowell, 1994). Soil particle size for texture determination was performed by mechanical analysis of the samples following the Bouyoucos Hydrometer Method (Ashworth et al., 2001; Bouyoucos, 1962). Organic matter values were determined by using the wet-oxidation method of Walkley-Black (Walkley and Black 1934) and total nitrogen was analyzed by usin midified-kjeldahl digestion method (Karaöz, 1992; Rowell, 1994). After saturating soil samples with taps water for 24 hours, soil water content at the field capacity (FC) was measured equilibrating soil moisture for 96 hours at 1500 kPa by using a pressure plate apparatus (Gülser and Candemir, 2014).

## 2.2.3. Soil microbial biomass and microbial respiration

Soil samples were removed from the refrigerator and equilibrated at room temperature for the soil microbial biomass analysis. Then, soil  $C_{mic}$  was analyzed in the incubation beakers by following the chloroform-fumigation-extraction method (Ladd and Amato, 1989). The  $C_{mic}$  of soils were determined by using the modified Walkley-Black method (Brookes et al., 1985; Vance et al., 1987). Briefly, 30 g (oven-dry basis) moist soil was fumigated for 24 h at 25 °C with ethanol-free chloroform (CHCl<sub>3</sub>). Following fumigant removal, the sample was extracted with 40 mL 0.5 M K<sub>2</sub>SO<sub>4</sub> by 30 min horizontal shaking at 90 rpm min<sup>-1</sup> and filtered through a folded filter paper. Another 30 g portion of the same soil sample was also weighed for the nonfumigation analysis. The nonfumigated samples was similarly extracted with 40 mL 0.5 M K<sub>2</sub>SO<sub>4</sub> by 30 min horizontal shaking at 90 rpm min<sup>-1</sup> and filtered through a folded filter paper. Finally, the  $C_{mic}$  was calculated as folded filter paper without utilizing the chloroform-fumigation process. Finally, the  $C_{mic}$  was calculated as follows in Eq. (1) (Vance et al., 1987):

Microbial Biomass C (
$$C_{mic}$$
) =  $K_{EC} \times 2.64$ 

(1)

(2)

where  $K_{EC}$  refers to the difference in extractable organic C between the fumigated and unfumigated samples and 2.64 is the proportionality factor that accounts for the biomass C released by the <u>fumigation</u> <u>extraction</u> process.

The  $N_{mic}$  was analyzed according to the Modified Kjeldahl digestion method by using the following Eq. (2) (Brookes et al., 1985; Anderson and Ingram, 1996):

Microbial Biomass N ( $N_{mic}$ ) =  $F_N/0.54$ 

where  $F_N$  is the difference between N extracted from fumigated and unfumigated samples and 0.54 is the fraction of microbial biomass N released by the fumigation extraction process.

Microbial respiration (MR) was estimated with the <u>sodium hydroxide</u> (NaOH) trap method (<u>Alef, 1995</u>). Briefly, 30 g oven-dry equivalent weight was transferred separately into 250 mL capacity glass beakers.

Small beakers filled with 10 mL of 1 M NaOH were placed at the bottom of the jar to trap the evolved CO<sub>2</sub>. The jars were fastened airtight and incubated for 7 d at 25 °C. After a week of incubation, the small beaker with the CO<sub>2</sub> absorbed in NaOH solution was removed and titrated with 0.1 M HCl after the addition of BaCl<sub>2</sub>. The NaOH solution without soil, incubated as above, was also titrated. Then, MR was calculated as the amount of CO<sub>2</sub> evolution over the incubation period divided by the dry mass of soil. The *q*CO<sub>2</sub> was calculated as the ratio between MR rates (mg CO<sub>2</sub>-C h<sup>-1</sup> g<sup>-1</sup>) to C<sub>mic</sub> (Anderson and Domsch, 1993).

## 2.3. Statistical analysis

All of the soil samples taken from the field in different periods were analyzed in paralel. The SPSS 16.0 software package program was used for all statistical analyses. The normality of data was assessed with the UNIVARIATE procedure. Prior to analysis, assumptions of equal variances for each group were visually checked by plotting the studentized residuals against predicted values at P > 0.05. Across response variables and tree species, assumptions of homoscedasticity and approximation to <u>normal distribution</u> were met for all comparisons. By using the independent samples t test, the differences between some physical, chemical and microbiological properties of the soils in the control areas and the skidding areas were determined at the sgnificant level (P<0.05).

## 3. Results and Discussion

## 3.1. Physical and chemical properties of soil

The average values of some physical, chemical and moisture properties of soil samples taken from the rubbing and control points are shown in Table 2. According to these average values, sandy clay loam soil texture was noticed in both areas. The amount of sand, silt and clay of the top soils in the skidding area was found 62.8%, 12.9% and 24.3%, and in control areas 64.86%, 11.64% and 23.50%, respectively. There was no statistical difference in the mean values of the sand, silt and clay percentages of the top soils in the skidding and control areas (P>0.05). In the subsoils, it was determined that there was a decrease in the percentage of sand and an increase in the percentage of clay in the friction areas. The reason for this is thought to be due to the fact that 3-5 cm of the topsoil was torn away from the area and the topsoil of the skidding area was taken more deeply than the control area. The topsoils of both areas are lime-free and slightly alkaline and subsoils are less calcareous and medium alkaline soils. It was determined that soils were statistically significantly affected by harvesting in terms of lime and pH values (P<0.05). In terms of electrical conductivity average values, the EC values of the upper soils were higher than the lower soils. The mean values of soil organic carbon (SOC) and toral nitrogen (TN) were found to be statistically different from each other in the skidding (2.15%-0.13%) and control (2.90%-0.16%) areas (P<0.05). Also, It was determined that there was a significant decrease in the amount of carbon and nitrogen stock (26.17 C kg/ha-1.88 N kg/ha) in the topsoil of the skidding areas compared to the control areas (34.27 C kg/ha-2.01 N kg/ha) (Table 2). So, the carbon loss or displacements in the area as a result of harvesting activities are an effective factor in this.

Statistically significant differences were found in the  $C_{org}/N_{total}$  ratio (P<0.05). It is predicted that the rate of decomposition may have increased in parallel with the increasing soil temperature as a result of the removal of the litter layers from the soil surface in the skidding areas. Therefore, the  $C_{org}/N_{total}$  ratio was found to be lower (25%) in the skidding areas than in the control areas. As it is known, the increase in the amount of nitrogen in the soil is an indicator of increased mineralization and accelerated decomposition (Babur and Dindaroglu, 2020). Zeller et al. (2000) noticed that the high  $C_{org}/N_{total}$  (>20) ratio in forest ecosystems causes low carbon and nitrogen mineralization, while low  $C_{org}/N_{total}$  ratio (<20) ratio is an indicator of intense carbon and nitrogen mineralization.

As a result of the examination of the water potential of the soils, it was determined that the average

values of the FC, PWP and plant available water capacity (PAWC) were higher in the control areas and that the skidding negatively affected the water holding capacity of the soils. The mean particle density and bulk density values of the sample and control area soils were found to be statistically insignificant (P>0.05). In addition, the porosity of the soils was found to be almost close to each other.

Table 2. Comparison of some general characteristics of the soils belonging to the control and destroyed areas by harvesting activities in the production area (n=36).

Soil properties	Soil s	amples	Cor	itrol
	Topsoil	Subsoil	Topsoil	Subsoil
Sand %	62, 78±4,62a	60, 77±4,71a	64,86±4,60a	62, 53±5,13a
Silt %	12,93±1,53a	12,03±2,48a	11,60±0,58a	12,97±3,79a
Clay %	24,20±6,06a	27,20±4,05a	23,50±5,19a	24,50±3,98a
pH (H <sub>2</sub> O)	7,68±0,10a	7,55±0,08a	7,92±0,27a	7,66±0,26a
EC ds m <sup>-1</sup>	0,11±0,01a	0,04±0,01a	0,19±0,01b	$0,05{\pm}0,00c$
Corg %	2,15±0,15a	1,63±0,27a	2,90±0,32b	1,85±0,10a
Ntotal %	0,13±0,03a	0,08±0,01a	0,16±0,03a	$0,08{\pm}0,00a$
Corg/Ntotal	13,94±0,22a	15,83±1,41a	17,18±1,17b	20,60±0,51a
Cstock kg/ha	26,17±1,45a	20,48±3,01a	34,27±2,92b	23,41±1,24a
Nstok kg/ha	1,88±0,12a	1,30±0,23a	2,01±0,30b	1,14±0,04a
FC	27,97±1,52a	30,27±0,59a	29,96±2,04a	33,05±1,30a
PWP	16,41±1,53a	15,92±1,11a	16,51±1,71a	18,49±0,94a
AWHC	11,55±2,53a	14,36±1,24a	13,45±0,95a	14,53±0,72a
Bulk density	1,22±0,03a	1,26±0,03a	1,19±0,29a	1,27±0,02a
Particle density	2,67±0,04a	2,77±0,05a	2,58±0,25a	2,78±0,04a
Porosity	54,37±1,22a	54,64±1,73a	54,04±1,48a	54,43±1,41a

Abbreviations: EC: Electrical conductivity, Corg: Organic carbon, Ntotal: Total nitrogen, FC: field capacity, PWP; parmenent wilting point, AWHC: Available water holding capacity

## 3.2. Seasonal changes of microbial soil properties

As a result of seasonal variations of the temperature and moisture values of the soils belonging to the sampling and control areas, it was determined that the soil surface was significantly affected by the climatic conditions as a result of the destruction of the litter covering the surface of the soil in the skidding areas and the breaking of the stand cover as a result of production (P<0.05; Table 3). The increase in temperature and the low rainfall in summer caused a significant increase in soil temperature. Although the temperature of the sample area soils is higher than that of the control areas in summer season, this is the opposite in terms of soil moisture (Table 3). The destruction of the litter layers, which protects the soil against external influences and acts as a buffer, along the skidding line has caused a significant decrease in soil moisture. As a result of the seasonal changes of the pH and EC average values of the topsoil, the highest pH average value of 7.92 was found in the control areas in the spring, and the lowest pH value of 7.45 in the summer season in the plowing areas. Also, in the subsoils, the highest average pH values were observed in the spring and autumn seasons. The highest mean EC values of the topsoils were found in the summer and autumn seasons of the control areas soils, as well. There was no statistical difference in the mean EC values in the subsoils.

When the changes in the OC and TN amounts of the topsoils according to the seasons were examined, it was determined that the highest carbon content (3.01%) was in the control areas in autumn, and the lowest OC amount was in the summer months. In order to meet the nutrient requirements of plants and microbial organisms in the summer season, the organic material in the soil must be synthesized and organic matter consumption is higher than the amount of organic matter entering the soil, causing a decrease in the amount of organic carbon (Kara et al., 2016; Babur et al., 2021). The highest total nitrogen content of the soil (0.18%) was found in the autumn season in the control area, and the lowest total nitrogen average value (0.13%) was found in the spring season in the skidding areas. The total nitrogen average values of the soils in the subsoil level were found to be close to each other and had no significant difference (Table 4).

It was determined that the harvesting activities had a statistically significant effect on the  $C_{\text{mic}}$ ,  $N_{\text{mic}}$  and

MR average values of the soils. The  $C_{mic}$  values of the topsoil in the sampling areas were lowest in the summer with 657.17 µg•g<sup>-1</sup> and the highest in the autumn with 763.76 µg•g<sup>-1</sup>. In the control areas, the lowest  $C_{mic}$  value was 773.99 µg•g<sup>-1</sup> in spring and the highest 886.15 µg•g<sup>-1</sup> in autumn (Figure 3 and 4). Significant reductions in microbial biomass were observed with loss of organic carbon in the soils in the skidding areas. This is thought to be due to the loss of organic matter and moisture in the soil which are the most important parameters for the life of soil microorganisms (Patel et al., 2010; Babur and Dindaroğlu, 2020). In addition, statistically significant differences were found in microbial properties in parallel with seasonal variations in changing the temperature and humidity values in the soil (*P*<0.05; Table 3 and 4). It was also determined that the amount of  $N_{mic}$  in the both soil layers of the control areas of the research areas was higher than the amount of  $N_{mic}$  in the sample areas (Table 3 and 4; Figure 4).

From the results obtained, it was determined that the chemical and microbial properties of the topsoil in the skidding areas decreased significantly compared to the control areas. Soil macro and micro organisms are very sensitive to even the smallest changes in the soil (Babur ve Dindaroglu, 2020). Microbial biomass and microbial activities of soils significantly affect the amount of soil organic matter (Jenkinson, 1988). In addition, some other characteristics such as temperature, humidity, pH, lime content, soil structure and texture, parent material, plant species and stand type ...etc. affect the populations and contents of soil microbial organisms (Patel et al., 2010; Babur, 2019). The results found in this study ara similar to soil  $C_{mic}$  (61–2,000 lg g-1) found in temperate forests by Vance et al., (1987b); (102–2,073 lg g-1) in tropical forests by Hernot and Robertson (1994) and (386–1,050 lg g-1) in larch forests by Kara and Bolat (2007).

From the data obtained in the study, it was determined that the microbial biomass carbon (r=0.709<sup>\*</sup>) and nitrogen (r=0.915<sup>\*\*</sup>) values of the topsoil were positively and strongly related to soil moisture and therefore negatively related to temperature (r=-0.317<sup>\*</sup> ve r=-0.676<sup>\*</sup>, respectively). In addition, soil C<sub>mi</sub> and N<sub>mi</sub> values were found to have positive relationships with soil carbon (r=0.719<sup>\*</sup> and r=0.725<sup>\*</sup>) and nitrogen values (r=0.885<sup>\*\*</sup> and r=0.629<sup>\*</sup>, respectively).

Microbial quotient ( $C_{mic}/C_{org}$ ) and metabolic quotient (MR/ $C_{mic}$ ) ratios of microbial indexes of topsoils in the study areas have changed significantly (*P*<0.05). It was also found that the  $C_{mic}/N_{mic}$  ratio in the skidding areas were higher than the control areas. The  $C_{mic}/N_{mic}$  ratio is used as an indicator of the microorganism type. A ratio of 3-5 indicates that the bacterial population is dominant, while a ratio of 10-12 indicates that the fungal population is dominant (Devi and Yadava, 2006). According to our results, it was determined that both bacterial and fungal populations are present in the topsoils of the research area, but the bacterial population is more dominant into the deeper of soil (Table 3 and 4).  $N_{mic}/N_{total}$  percentages of the topsoil of the skidding areas were higher than the control area in all seasons except summer. Statistically significant differences were found in the  $N_{mic}/N_{total}$  percentages in different seasons (P<0.05). It was determined that the  $N_{mic}/N_{total}$  amount of control areas was higher in subsoils. A high rate indicates that the amount of available nitrogen in the organic matter composition is good (Khan and Joergensen 2006). If this value is low, it is an indication of poor soil substrate quality (Bauhus et al., 1998).

Microbial soil respiration is the amount of  $CO_2$  released into the atmosphere as a result of the oxidation of organic matter by aerobic microorganisms. Soil microbial respiration is used to predict the microbial activity of soils (Winding et al., 2005). According to Sparling (1997) and Alvarez et al., (2009), soil microbial respiration is a very important parameter in monitoring the rate of decomposition. However, soil respiration is affected by ecological and edaphic factors such as moisture content, temperature, substrate quality...etc. (Tufekcioglu et al., 2006; Güner et al., 2010; Babur, 2019). Nilsen and Strand (2008) determined that thinning affects soil respiration by directly affecting soil moisture, temperature, litter biomass and stand cover.

The mean values of soil microbial respiration in the skidding areas were found to be higher than the

soils of the control areas, in contrast to the  $C_{mic}$  and  $N_{mic}$  values. The reason for this is that microbial organisms tend to continue their lives by breathing more due to moisture loss in the soil that is directly exposed to external factors such as sunlight after skidding. In this study, microbial respiration have average values varying between 1.24 and 2.08 lg CO<sub>2</sub>–C g<sup>-1</sup> h<sup>-1</sup> in the topsoils and 0.68–0.99 lg CO<sub>2</sub>–C g<sup>-1</sup> h<sup>-1</sup> in the subsoils (Figure 5). Microbial respiration directly affects the carbon storage capacity in terrestrial ecosystems as it affects all of the soil organic matter and other microbial parameters (Alef, 1995; Winding et al., 2005).

It was determined that the metabolic quotient ( $qCO_2$ ) average values of the topsoils were highest in the skidding area soils in summer months and the lowest values in the control areas in the spring. According to the seasons, the  $qCO_2$  average values of the skidding soils were found to be statistically higher than the control (P<0.05). Similarly, Anderson (2003) emphasized that neutral microbial respiration ranged between 0.5 and 2.0 mg CO<sub>2</sub>–C g<sup>-1</sup> C<sub>mic</sub> h<sup>-1</sup>. Bolat (20014) found a negative relationship between  $qCO_2$  and SOC, pH and lime. High  $qCO_2$  amount is one of the stress indicators of terrestrial ecosystem. Odum (1985), and Anderson and Domsch (1993) also stated that the amount of  $qCO_2$  is higher in unhealthy soils. However, it was stated that low  $qCO_2$  amount was associated with plant succession (Insam and Haselwandter, 1989). In addition,  $qCO_2$  provides information about the ecophysiological (Anderson and Domsch 1985; Insam et al., 1996), substrate quality and availability of soil microbial communities (Dilly et al., 1997; Dilly and Munch, 1998).

The percentage of  $C_{mic}/C_{org}$  (microbial quotient) also correlates with substrate quality. This rate is also sensitive to changes in the ecosystem and is highly affected (Bauhus et al. 1998; Anderson, 2003). In addition, this ratio can help us to predict changes in soil carbon (Anderson and Domsch 1989). Jenkinson and Ladd (1981) stated that the equilibrium threshold of  $C_{mic}/C_{org}$  percentage in soil is 2.2. Also, the  $C_{mic}/C_{org}$  percentage has been reported to have a wide range of values between 0.27 and 7.0% (Anderson and Domsch, 1989). In this study, it was determined that the  $C_{mic}/C_{org}$  average values of the skidding soils were higher than those of the control soils. According to data, the highest  $C_{mic}/C_{org}$  ratio was found in the topsoils of the skidding areas in autumn (3.64%) and the lowest (2.17%) in the subsoils of control in the spring (Tables 3 and 4). Smilarly, Kara et al., (2008) found the ratio of  $C_{mic}/C_{org}$  in the soils of temperate forests to be between 2.26–3.17%.

These results show us that changes in some factors such as soil, plant biodiversity, forestry practices and sampling time have a significant effect on the change in the  $C_{mic}/C_{org}$  ratio (Anderson and Domsch, 1989; Vesterdal et al., 1995).

Soil	Soil sample			Control		
properties	Spring	Summer	Autumn	Spring	Summer	Autumn
Temp. C <sup>o</sup>	15,30±0,27a	25,93±0,60c	18,13±0,15e	13,97±0,15b	22,33±0,21d	19,57±0,12f
SM %	30,53±1,02a	11,88±0,84c	42,98±1,23e	38,88±2,73b	27,99±1,25d	53,27±0,32f
pH(H <sub>2</sub> O)	7,68±0,10a	7,45±0,11	7,63±0,12e	7,92±0,27a	$7,80\pm0,22$	7,91±0,24e
EC ds m <sup>-1</sup>	0,11±0,01a	0,13±0,02c	0,13±0,01e	0,19±0,01b	0,20±0,00d	0,20±0,02f
OC %	2,15±0,15a	2,02±0,11c	2,10±0,15e	2,90±0,32b	2,85±0,30d	3,01±0,36f
TN%	0,13±0,03a	$0,16\pm0,01$	0,14±0,02e	0,16±0,03a	$0,17\pm0,02$	0,18±0,04e
C <sub>mic</sub>	700,9±71,6a	657,17±65,4c	763,57±76,3e	773,99±70,1b	855,17±77,6d	886,15±75,5e
N <sub>mic</sub>	97,65±1,95a	93,62±2,18c	105,19±4,07e	112,20±9,43a	118,49±5,05d	124,29±6,03f
MR	1,24±0,11a	2,08±0,30c	1,55±0,32e	1,30±0,17a	1,69±0,10d	1,67±0,11e
$C_{mic}/N_{mic}$	7,17±0,63a	7,01±0,55	7,27±0,79e	6,90±0,10a	7,21±0,38	7,13±0,41e
$C_{mic}/C_{org}$	3,26±0,20a	$3,25\pm0,27$	3,64±0,23e	2,68±0,24b	3,01±0,29	3,00±0,35f
N <sub>mic</sub> /N <sub>total</sub>	7,72±1,60a	6,09±0,32	7,87±1,62e	7,18±1,03a	$7,08\pm0,96$	7,66±0,94e
MR/C <sub>mic</sub>	1,77±0,22a	3,17±0,10c	2,03±0,11e	1,68±0,19a	1,98±0,19d	1,88±0,15f

Table 3. Seasonal changes in some physicochemical and microbial properties of topsoils (0-10 cm) (n=12).

Abbreviations: SM; soil moisture, EC; electrical conductivity, OC; organic carbon, TN; total nitrogen, MBC; microbial biomass carbon, MBN; microbial biyomass nitrogen, MR; microbial respiration,

EJFS-Short-term monitoring of the winching and skidding effects on soil microbial biomass in Turkish red pine in the Mediterranean Region – by Babur 2021

Soil	Soil sample			Control		
properties	Spring	Summer	Autumn	Spring	Summer	Autumn
Temp. C <sup>o</sup>	14,50±0,17a	23,57±0,40c	19,03±0,25e	13,33±0,2b	21,17±0,21d	20,40±0,44f
SM%	26,15±1,59a	17,70±1,14c	39,04±0,80e	42,39±1,5b	30,04±1,29d	50,23±2,75f
$pH(H_2O)$	7,55±0,08a	7,40±0,04c	7,57±0,06e	7,66±0,26a	7,59±0,21c	7,66±0,27e
EC ds m <sup>-1</sup>	0,04±0,01a	0,04±0,06c	0,05±0,01e	0,04±0,00a	0,05±0,00c	0,04±0,00e
OC %	1,63±0,27a	1,58±0,28c	1,67±0,26e	1,85±0,10a	1,82±0,10c	1,93±0,07e
TN%	0,08±0,01a	0,10±0,01c	0,09±0,01e	0,08±0,00a	0,09±0,00c	0,09±0,00e
MBC	382,37±12,2a	401,2±12,3c	432,6±21,1e	400,9±9,8a	438,5±25,8c	457,3±25,7e
MBN	65,11±4,1a	68,60±3,1c	72,41±2,2e	93,74±2,5b	97,91±1,0d	103,01±1,8f
MR	0,68±0,03a	0,99±0,05c	0,87±0,06e	0,75±0,07a	0,88±0,09c	0,85±0,10e
$C_{mic}/N_{mic}$	5,89±0,42a	5,86±0,37c	5,99±0,66e	4,28±0,21b	4,48±0,25d	4,44±0,25f
$C_{mic}/C_{org}$	2,38±0,36a	2,59±0,44c	2,62±0,30e	2,17±0,10a	2,41±0,03c	2,37±0,06e
Nmic/Ntoplam	8,82±1,55a	6,77±1,25c	8,64±1,77e	11,92±1,54a	10,93±0,36d	11,28±0,62e
MR/C <sub>mic</sub>	1,77±0,26a	2,46±0,25c	2,01±0,22e	1,87±0,23a	2,01±0,14c	1,86±0,23e

Table 4. Seasonal changes in some physicochemical and microbial properties of subsoils (10-20 cm) (n=12).



**Figure 3.** Seasonal changes in microbial biomass C content of topsoil (a) and subsoil (b) samples. Values showed by different numbers and lowercase letters differ significantly between seasons (p < 0.05).



**Figure 4.** Seasonal changes in microbial biomass N content of topsoil (a) and subsoil (b) samples. Values showed by different numbers and lowercase letters differ significantly between seasons (p < 0.05).

EJFS-Short-term monitoring of the winching and skidding effects on soil microbial biomass in Turkish red pine in the Mediterranean Region – by Babur 2021



**Figure 5.** Seasonal changes in microbial respiration of topsoil (a) and subsoil (b) samples. Values showed by different numbers and lowercase letters differ significantly between seasons (p < 0.05).

## 4. Conclusion

In this study, the effects of the harvesting application on some physicochemical and microbial properties in the short term were investigated after the whole tree method was removed from the partition using a farm tractor. As a result of short-term monitoring in the research area, it has been observed that the skidding and production processes of wood raw materials cause significant changes in the soil properties such as the organic carbon, total nitrogen, temperature, moisture, microbial biomass C, N, and microbial respiration. Significant decreases were observed in the carbon and nitrogen stoks and microbiological properties of the soils compared to the control areas, especially in the summer months when drought is observed in the soil destroyed as a result of harvesting. The most negative effect of skidding can be seen as degradation and destroying the topsoil (5-10 cm deep-Ah horizon) with the dead cover on it. The microbial respiration and metabolic quotient values in the control areas are less than the sample areas. This is an indication that the microbial population in the sampling areas consumes more energy in parallel with the increase in temperature and that stress conditions begin to occur in the soil. In forestry activities, it is important for forest and soil health to monitor the changes in the microbiological properties of soils over long periods and to produce in a way that will cause minimum damage to the soil. For this reason, skidding is required during the periods of production activities when the soils are harder. In addition, there is a need to develop alternative extraction methods without skidding and destruction of the soil surface.

#### Acknowldgemend

I am grateful to Dr. Sercan GÜLCİ at the Kahramanmaraş Sütçü Imam University and unit chief Nihat NURDOĞAN at the Kahramanmaraş Forestry General Diroctorate for their valuable Assisting in the supply of the research area and the equipment used.

#### References

Alef, K. (1995). Soil respiration. In: Alef K, Nannipieri P (eds) Methods in applied soil microbiology and biochemistry. Academic Press, London, pp 214–218.

Allen, A.S., Schlesinger, W.H. (2004). Nutrient limitations to soil microbial biomass and activity in loblolly pine forests. Soil Biol. Biochem. 36, 581–589.

Alvarez, E., Torrado, V.M., Fernandez Marcos, M.L., Diaz-Ravia, M. (2009). Microbial biomass and activity in forest soil under different tree species. Electron J Environ Agric Food Chem 8(9):878–887

Anderson, T.H. (2003). Microbial eco-physiological indicators to assess soil quality. Agric Ecosyst Environ 98:285–293.

Anderson, T.H., Domsch, K.H. (1985). Maintenance requirements of actively metabolizing microbial populations under in situ conditions. Soil Biol Biochem 17:197–203.

Anderson, J.P.E., Domsch, K.H. (1989). Ratios of microbial biomass carbon to total organic carbon in arable soils. Soil Biol Biochem 21:471–479.

Anderson, T.H., Domsch, K.H. (1990). Application of eco-physiological quotients (qCO2, and qD) on microbial biomasses from soils of different cropping histories. Soil Biol Biochem 22:251–255.

Anderson, T.H., Domsch, K.H. (1993). The metabolic quotient for CO2 (qCO2) as a specific activity parameter to assess the effects of environmental conditions, such as pH, on the microbial biomass of forest soils. Soil Biol Biochem 25:393–395.

Anderson, J.M., Ingram, J.S.I. (1996). Tropical soil biology and fertility a handbook of methods, 2nd edn. Cab International Wallingford, UK.

Ashworth, J., Keyes, D., Kirk, R., Lessard, R. (2001). Standard procedure in the hydrometer method for particle size analysis. Commun. Soil Sci. Plant Anal. 32 (5–6), 633–642.

Babur, E. (2019). Effects of parent material on soil microbial biomass carbon and basal respiration within young afforested areas, Scandinavian Journal of Forest Research, 34:2, 94-101, DOI: 10.1080/02827581.2018.1561936.

Babur E, Dindaroglu T. (2020). Seasonal Changes of Soil Organic Carbon and Microbial Biomass Carbon in Different Forest Ecosystems, Environmental Factors Affecting Human Health, Ivan Uher, IntechOpen, DOI: 10.5772/intechopen.90656.

Babur, E., Dindaroğlu, T., Solaiman, Z., Battaglia M.L., (2021). Microbial respiration, microbial biomass and activity are highly sensitive to forest tree species and seasonal patterns in the Eastern Mediterranean Karst Ecosystems. Science of the Total Environment. 775, 145868.

Bauhus, J.D., Pare, D., Cote, L. (1998). Effects of tree species, stand age, and soil type on soil microbial biomass and its activity in a southern boreal forest. Soil Biol Biochem 30:1077–1089.

Bolat, I. (2014). The effect of thinning on microbial biomass C, N and basal respiration in black pine forest soils in Mudurnu, Turkey. Eur J Forest Res (2014) 133:131–139.

Bouyoucos, G.J. (1962). Hydrometer method improved for making particle size analyses of soils. Agron J 54:464-465.

Brookes, P.C., Landman, A., Pruden, G., Jenkinson, D.S. (1985). Chloroform fumigation and the release of soil nitrogen: a rapid extraction method to measure microbial biomass nitrogen in soil. Soil Biol Biochem 17:837–842.

Cambi, M., Certini, G., Neri, F., Marchi, E. (2015). The impact of heavy traffic on forest soils: a review. For. Ecol. Manage. 338, 124–138.

Carter, M.C., Dean, T.J., Zhou, M., Messina, M.G., Wang, Z. (2002). Short-term changes in soil C, N, and biota following harvesting and regeneration of loblolly pine (Pinus taeda L.). For. Ecol. Manage. 164, 67–88.

Chen, G., Tian, H., Huang, C., Prior, S.A., Pan, S. (2013). Integrating a process-based ecosystem model with Landsat imagery to assess impacts of forest disturbance on terrestrial carbon dynamics: case studies in Alabama and Mississippi. J. Geophys. Res.: Biogeosci. 118, 1208–1224.

Dangal, S.R.S., Felzer, B.S., Hurteau, M.D. (2014). Effects of agriculture and timber harvest on carbon sequestration in eastern US forests. J. Geophys. Res.: Biogeosci. 119, 35–54.

Devi, N.B., Yadava, P.S. (2006). Seasonal dynamics in soil microbial biomass C, N and P in a mixed-oak forest ecosystem of Manipur, Northeast India. Appl Soil Ecol 31:220–227.

Dilly, O., Munch, J.C. (1998). Ratios between estimates of microbial biomass content and microbial activity in soils. Biol Fertil Soils 27:374–379.

Dilly, O., Bernhard, M., Kutsch, W.L., Kappen, L., Munch, J.C. (1997). Aspects of carbon and nitrogen cycling in soils of the Bornhoved Lake district I. Microbial characteristics and emission of carbon dioxide and nitrous oxide of arable and grassland soils. Biogeochemistry 39:189–205.

Foote, J.A., Boutton, T.W., Scott, D.A. (2015). Soil C and N storage and microbial biomass in US southern pine forests: Influence of forest management. Forest Ecology and Management 355 48–57.

Frey, B., Kremer, J., Rudt, A., Sciacca, S., Matthies, D., Luscher, P. (2009). Compaction of forest soils with heavy logging machinery affects soil bacterial community structure. Eur. J. Soil Biol. 45, 312–320.

Grand, S., Lavkulich, L.M. (2012). Effects of forest harvest on soil carbon and related variables in Canadian Spodosols. Soil Sci. Soc. Am. J. 76, 1816–1827.

Gülser. C., Candemir, F. (2014). Using soil moisture constants and physical properties to predict saturated hydraulic conductivity. Eurasian Journal of Soil Science 3 (2014) 77 – 81.

Güner, S., Tüfekcioğlu, A., Gülenay, S., Küçük, M. (2010). Land-use type and slope position effects on soil respiration in black locust plantations in Artvin, Turkey. African Journal of Agricultural Research 5(8):719-724.

Haubensak, K.A., Hart, S.C., Stark, J.M. (2002). Influences of chloroform exposure time and soil water content on C and N release in forest soils. Soil Biol. Biochem. 34, 1549–1562.

Henderson, G.S. (1995). Soil organic matter: a link between forest management and productivity. In: McFee, W.W., Kelly, J.M. (Eds.), Carbon Forms and Functions in Forest Soils. Soil Science Society of America Inc., Madison, pp. 419–435.

Hernot, J., Robertson, G.P. (1994). Vegetation removal in two soils of the humid tropics: effect on microbial biomass. Soil Biol Biochem 26:111–116.

Insam, H., Haselwandter, K. (1989). Metabolic quotient of the soil microflora in relation to plant succession. Oecologia 79:174–178.

Insam, H., Hutchinson, T.C., Reber, H.H. (1996). Effects of heavy metal stress on the metabolic quotient of soil microflora. Soil Biol Biochem 28:691–694.

Jenkinson, D.S. (1988). The determination of microbial biomass carbon and nitrogen in soil. In: Wilson JR (ed) Advances in nitrogen cycling in agricultural ecosystems. CAB, Wallingford, pp 368–386.

Jenkinson, D.S., Ladd, J.N. (1981). Microbial biomass in soil measurement and turnover. In: Paul EA, Ladd JN (eds) Soil biochemistry, vol 5. Marcel Dekker Inc, New York and Basel, pp 415–471.

Jordan, D., Ponder Jr., F., Hubbard, V.C. (2003). Effects of soil compaction, forest leaf litter, and nitrogen fertilizer on two oak species and microbial activity. Appl. Soil Ecol. 23, 33–41.

Jones, H.S., Beets, P.N., Kimberley, M.O., Garrett, L.G. (2011). Harvest residue management and fertilisation effects on soil carbon and nitrogen in a 15- year-old Pinus radiata plantation forest. For. Ecol. Manage. 262, 339–347.

Joergensen, R.G., Wu, J., Brookes, P.C. (2011). Measuring soil microbial biomass using an automated procedure. Soil Biol. Biochem. 43, 873–876.

Kara, O., Bolat, I. (2007). The effect of wildfire on the microbial biomass C of black pine plantation soils. In: Proceedings of international symposium, bottlenecks, solutions, and priorities in the context of functions of forest resources, October 17–19, Istanbul University, Faculty of Forestry, Istanbul, Turkey, pp 1021–1030.

Kara, O., Bolat, I., Cakıroglu, K., Ozturk, M. (2008). Plant canopy effects on litter accumulation and soil microbial biomass in two temperate forests. Biol Fertil Soils 45(2):193–198.

Kara, O., Babur, E., Altun, L., Seyis, M. (2016). Effects of afforestation on microbial biomass C and respiration in eroded soils of Turkey. J Sustain For 35(6):385–396.

Karaöz M.Ö. (1992). Leaf and litter analysis methods (in Turkish). J. Fac. For. Istanbul U. Series B, 42 (1–2), pp. 57-71

Kellman, L., Kumar, S., Diochon, A. (2014). Soil nitrogen dynamics within soil profiles of a managed moist temperate forest chronosequence consistent with longterm harvesting-induced losses. J. Geophys. Res.: Biogeosci. 119, 1309–1321.

Khan, K.S., Joergensen, R.G. (2006). Microbial C, N and P relationships in moisture stressed soils of Potohar, Pakistan. J Plant Nutr Soil Sci 169:494–500.

Labelle, E.R., Jaeger, D. (2011). Soil compaction caused by cut-to-length forest operations and possible short-term natural rehabilitation of soil density. Soil Sci. Soc. Am. J. 75, 2314–2329.

LeDuc, S.D., Rothstein, D.E. (2007). Initial recovery of soil carbon and nitrogen pools and dynamics following disturbance in jack pine forests: a comparison of wildfire and clearcut harvesting. Soil Biol. Biochem. 39, 2865–2876.

Li, Q., Allen, H.L., Wollum II, A.G. (2004). Microbial biomass and bacterial functional diversity in forest soils: effects of organic matter removal, compaction, and vegetation control. Soil Biol. Biochem. 36, 571–579.

Mariani, L., Chang, S.X., Kabzems, R. (2006). Effects of tree harvesting, forest floor removal, and compaction on soil microbial biomass, microbial respiration, and N availability in a boreal aspen forest in British Columbia. Soil Biol. Biochem. 38, 1734–1744.

Nilsen, P., Strand, L.T. (2008). Thinning intensity effects on carbon and nitrogen stores and fluxes in a Norway spruce (Picea abies (L.) Karst.) stand after 33 years. For Ecol Manage 256:201–208.

Odum, E.P. (1985). Trends expected in stressed ecosystems. Bioscience 35:419-422.

Patel, K., Nirmal Kumar, J.I.N., Kumar, R., Kumar Bhoi, R. (2010). Seasonal and temporal variation in soil microbial biomass C, N and P in different types land uses of dry deciduous forest ecosystem of Udaipur, Rajasthan, Western India. Appl Ecol Environ Res 8(4):377–390.

Paul, E.A., Harris, D., Klug, M.J., Ruess, R.W. (1999). The determination of microbial biomass. In: Robertson, G.P., Coleman, D.C., Bledsoe, C.S., Sollins, P. (Eds.), Standard Soil Methods for Long-Term Ecological Research. Oxford University Press, New York, pp. 291–317.

Rowell, D.L. (1994). Soil science; methods and applications. Longman Publishers (Pte) Ltd, Singapore.

Six, J., Bossuyt, H., Degryze, S., Denef, K. (2004). A history of research on the link between (micro) aggregates, soil biota, and soil organic matter dynamics. Soil Tillage Res. 79, 7–31.

Sparling, G.P. (1997). Soil microbial biomass, activity and nutrient cycling as indicators of soil health. In: Pankhurst CE, Doube BM, Gupta VVSR (eds) Biological indicators of soil health. CAB International, Wallingford, pp 97–119.

Torbert, H.A., Wood, C.W. (1992). Effects of soil compaction and water-filled pore space on soil microbial activity and N losses. Commun. Soil Sci. Plant Anal. 23 (11&12), 1321–1331.

Turner, J., Lambert, M.J. (2011). Analysis of nutrient depletion in a radiata pine plantation. For. Ecol. Manage. 262, 1327–1336.

Tüfekçioğlu, A., Küçük, M., Sağlam, B., Bilgili, E., Altun, L., Küçük, Ö. (2006). Influence of fire on root biomass dynamics and soil respiration rates in young corsican pine (Pinus nigra) stands in Turkey. Forest Ecology and Management, 234,: S195, https://doi.org/10.1016/j.foreco.2006.08.314.

Vance, E.D., Brookes, P.C., Jenkinson, D.S. (1987a). An extraction method for measuring soil microbial biomass C. Soil Biol Biochem 19:703–707.

Vance, E.D., Brookes, P.C., Jenkinson, D.S. (1987b). Microbial biomass measurements in forest soils: the use of chloroform fumigationincubations methods for strongly acid soils. Soil Biol Biochem 19:697–702.

Walkley, A., Black, I.A. (1934). An examination of the degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. *Soil Sci.* 37, 29–38.

Wardle, D.A. (1992). A comparative assessment of factors which influence microbial biomass carbon and nitrogen levels in soil. Biol. Rev. 67, 321–358.

Vesterdal, L., Dalsgaard, M., Felby, C., Raulund-Rasmussen, K., Jorgensen, B.B. (1995). Effects of thinning and soil properties on accumulation of carbon, nitrogen and phosphorus in the forest floor of Norway spruce stands. For Ecol Manage 77:1–10.

Winding, A., Hund-Rinke, K., Rutgers, M. (2005). The use of microorganisms in ecological soil classification and assessment concepts. Ecotoxicol Environ Saf 62:230–248.

Zeller, B., Colin-Belgrand, M., Dambrine, E., Martin, F., Bottner, P. (2000). Decomposition of 15 N-labelled beech litter and fate of nitrogen derived from litter in a beech forest. Oecologia 123:550–559.

Dergiye başvuru tarihi: 09.10.2021

Yayınlanmaya kabul edilme tarihi: 29.09.2021