

PAPER DETAILS

TITLE: Some Refinements and Reverses of Callebaut's Inequality for Isotonic Functionals via a Result Due to Cartwright and Field

AUTHORS: Sever Dragomir

PAGES: 1-11

ORIGINAL PDF URL: <https://dergipark.org.tr/tr/download/article-file/3417847>

Some Refinements and Reverses of Callebaut's Inequality for Isotonic Functionals via a Result Due to Cartwright and Field

Sever Silvestru Dragomir^{1,2,†}, 

¹Mathematics, School of Engineering & Science Victoria University, PO Box 14428, Melbourne City, MC 8001, Australia

²School of Computer Science & Applied Mathematics, University of the Witwatersrand, Private Bag 3, Johannesburg 2050, South Africa

†sever.dragomir@vu.edu.au

Article Information

Keywords: Isotonic functionals; Schwarz's inequality; Callebaut's inequality; Integral inequalities; Discrete inequalities

AMS 2020 Classification: 26D15; 26D10

Abstract

In this paper we obtain some refinements and reverses of Callebaut's inequality for isotonic functionals via a result of Young's inequality due to Cartwright and Field.

1. Introduction

Let L be a linear class of real-valued functions $g : E \rightarrow \mathbb{R}$ having the properties

- (L1) $f, g \in L$ imply $(\alpha f + \beta g) \in L$ for all $\alpha, \beta \in \mathbb{R}$;
 (L2) $\mathbf{1} \in L$, i.e., if $f_0(t) = 1, t \in E$ then $f_0 \in L$.

An isotonic linear functional $A : L \rightarrow \mathbb{R}$ is a functional satisfying

- (A1) $A(\alpha f + \beta g) = \alpha A(f) + \beta A(g)$ for all $f, g \in L$ and $\alpha, \beta \in \mathbb{R}$.
 (A2) If $f \in L$ and $f \geq 0$, then $A(f) \geq 0$.

The mapping A is said to be *normalized* if

- (A3) $A(\mathbf{1}) = 1$.

Isotonic, that is, order-preserving, linear functionals are natural objects in analysis that enjoy a number of convenient properties. Thus, they provide, for example, Jessen's inequality, which is a functional form of Jensen's inequality (see [1], [2] and [3]). For other inequalities for isotonic functionals see [4], and [5]-[15].

We note that common examples of such isotonic linear functionals A are given by

$$A(g) = \int_E g d\mu \text{ or } A(g) = \sum_{k \in E} p_k g_k,$$

where μ is a positive measure on E in the first case and E is a subset of the natural numbers $\mathbb{N}$, in the second, with $g = \{g_k\}_{k \in E}$ and $p_k \geq 0, k \in E$.

We have the following inequality that provides a refinement and a reverse for the celebrated *Young's inequality*

$$\frac{1}{2} v(1-v) \frac{(b-a)^2}{\max\{a, b\}} \leq (1-v)a + vb - a^{1-v}b^v \leq \frac{1}{2} v(1-v) \frac{(b-a)^2}{\min\{a, b\}} \quad (1.1)$$

for any $a, b > 0$ and $\nu \in [0, 1]$.

This result was obtained in 1978 by Cartwright and Field [16] who established a more general result for n variables and gave an application for a probability measure supported on a finite interval.

The functional version of *Callebaut's inequality* states that

$$A^2(fg) \leq A(f^{2-\nu}g^\nu)A(f^\nu g^{2-\nu}) \leq A(f^2)A(g^2) \quad (1.2)$$

provided that $f^2, g^2, f^{2-\nu}g^\nu, f^\nu g^{2-\nu}, fg \in L$ for some $\nu \in [0, 2]$. For the discrete and integral of one real variable versions see [17].

In this paper we obtain some inequalities for isotonic functionals via the reverse and refinement of Young's inequality (1.1) that are related to the second part of Callebaut's inequality (1.2). Applications for integrals and n -tuples of real numbers are also provided.

2. On Callebaut's Inequality

We have the following result that provides a refinement and reverse of Callebaut's second inequality:

Theorem 2.1. Let $A, B : L \rightarrow \mathbb{R}$ be two normalized isotonic functionals. If $f, g : E \rightarrow \mathbb{R}$ are such that, $f^2, g^2, \frac{g^4}{f^2}, f^{2(1-\nu)}g^{2\nu}, f^{2\nu}g^{2(1-\nu)} \in L$ for some $\nu \in [0, 1]$, and

$$0 < m \leq \frac{f}{g} \leq M < \infty \quad (2.1)$$

for real numbers $M > m > 0$, then

$$\begin{aligned} \frac{1}{2}\nu(1-\nu)m^2 \left(A\left(\frac{g^4}{f^2}\right)B(f^2) + A(f^2)B\left(\frac{g^4}{f^2}\right) - 2 \right) &\leq (1-\nu)A(f^2)B(g^2) + \nu A(g^2)B(f^2) \\ &\quad - A(f^{2(1-\nu)}g^{2\nu})B(f^{2\nu}g^{2(1-\nu)}) \\ &\leq \frac{1}{2}\nu(1-\nu)M^2 \left(A\left(\frac{g^4}{f^2}\right)B(f^2) + A(f^2)B\left(\frac{g^4}{f^2}\right) - 2 \right). \end{aligned} \quad (2.2)$$

Proof. Since $ab = \min\{a, b\} \max\{a, b\}$ for any $a, b > 0$, then from (1.1) we have

$$\frac{1}{2}\nu(1-\nu)\min\{a, b\} \frac{(b-a)^2}{ab} \leq (1-\nu)a + \nu b - a^{1-\nu}b^\nu \leq \frac{1}{2}\nu(1-\nu)\max\{a, b\} \frac{(b-a)^2}{ab},$$

where $\nu \in [0, 1]$. This can be written as

$$\frac{1}{2}\nu(1-\nu)\min\{a, b\} \left(\frac{b}{a} + \frac{a}{b} - 2 \right) \leq (1-\nu)a + \nu b - a^{1-\nu}b^\nu \leq \frac{1}{2}\nu(1-\nu)\max\{a, b\} \left(\frac{b}{a} + \frac{a}{b} - 2 \right), \quad (2.3)$$

for any $a, b > 0$.

Let $x, y \in E$ such that $g(x), g(y) \neq 0$. If we use the inequalities (2.3) for

$$a = \frac{f^2(x)}{g^2(x)}, \quad b = \frac{f^2(y)}{g^2(y)} \in [m^2, M^2]$$

then we get

$$\begin{aligned} \frac{1}{2}\nu(1-\nu)m^2 \left(\frac{f^2(y)}{g^2(y)} \frac{g^2(x)}{f^2(x)} + \frac{f^2(x)}{g^2(x)} \frac{g^2(y)}{f^2(y)} - 2 \right) &\leq (1-\nu) \frac{f^2(x)}{g^2(x)} + \nu \frac{f^2(y)}{g^2(y)} - \left(\frac{f^2(x)}{g^2(x)} \right)^{1-\nu} \left(\frac{f^2(y)}{g^2(y)} \right)^\nu \\ &\leq \frac{1}{2}\nu(1-\nu)M^2 \left(\frac{f^2(y)}{g^2(y)} \frac{g^2(x)}{f^2(x)} + \frac{f^2(x)}{g^2(x)} \frac{g^2(y)}{f^2(y)} - 2 \right), \end{aligned} \quad (2.4)$$

where $\nu \in [0, 1]$.

If we multiply (2.4) by $g^2(x)g^2(y)$, then we get

$$\begin{aligned} \frac{1}{2}\nu(1-\nu)m^2 \left(f^2(y) \frac{g^4(x)}{f^2(x)} + f^2(x) \frac{g^4(y)}{f^2(y)} - 2 \right) &\leq (1-\nu)g^2(y)f^2(x) + \nu f^2(y)g^2(x) \\ &\quad - f^{2\nu}(y)g^{2(1-\nu)}(y)f^{2(1-\nu)}(x)g^{2\nu}(x) \\ &\leq \frac{1}{2}\nu(1-\nu)M^2 \left(f^2(y) \frac{g^4(x)}{f^2(x)} + f^2(x) \frac{g^4(y)}{f^2(y)} - 2 \right), \end{aligned} \quad (2.5)$$

which holds for any $x, y \in E$.

Fix $y \in E$. Then by (2.5) we have in the order of L that

$$\begin{aligned} \frac{1}{2} \nu(1-\nu) m^2 \left(f^2(y) \frac{g^4}{f^2} + \frac{g^4(y)}{f^2(y)} f^2 - 2 \right) &\leq (1-\nu) g^2(y) f^2 + \nu f^2(y) g^2 - f^{2\nu}(y) g^{2(1-\nu)}(y) f^{2(1-\nu)} g^{2\nu} \quad (2.6) \\ &\leq \frac{1}{2} \nu(1-\nu) M^2 \left(f^2(y) \frac{g^4}{f^2} + \frac{g^4(y)}{f^2(y)} f^2 - 2 \right). \end{aligned}$$

If we take the functional A in (2.6), then we get

$$\begin{aligned} \frac{1}{2} \nu(1-\nu) m^2 \left(f^2(y) A \left(\frac{g^4}{f^2} \right) + \frac{g^4(y)}{f^2(y)} A(f^2) - 2 \right) &\leq (1-\nu) g^2(y) A(f^2) + \nu f^2(y) A(g^2) - f^{2\nu}(y) g^{2(1-\nu)}(y) A(f^{2(1-\nu)} g^{2\nu}) \\ &\leq \frac{1}{2} \nu(1-\nu) M^2 \left(f^2(y) A \left(\frac{g^4}{f^2} \right) + \frac{g^4(y)}{f^2(y)} A(f^2) - 2 \right), \end{aligned}$$

for any $y \in E$.

If we write this inequality in the order of L , then we have

$$\begin{aligned} \frac{1}{2} \nu(1-\nu) m^2 \left(A \left(\frac{g^4}{f^2} \right) f^2 + A(f^2) \frac{g^4}{f^2} - 2 \right) &\leq (1-\nu) A(f^2) g^2 + \nu A(g^2) f^2 - A(f^{2(1-\nu)} g^{2\nu}) f^{2\nu} g^{2(1-\nu)} \\ &\leq \frac{1}{2} \nu(1-\nu) M^2 \left(A \left(\frac{g^4}{f^2} \right) f^2 + A(f^2) \frac{g^4}{f^2} - 2 \right), \end{aligned}$$

and by taking the functional B we deduce the desired result (2.2). $\square$

Corollary 2.2. Let $A : L \rightarrow \mathbb{R}$ be a normalized isotonic functional. If $f, g : E \rightarrow \mathbb{R}$ are such that $f^2, g^2, \frac{g^4}{f^2}, f^{2(1-\nu)} g^{2\nu}, f^{2\nu} g^{2(1-\nu)} \in L$ for some $\nu \in [0, 1]$ and the condition (2.1) holds, then

$$\begin{aligned} \nu(1-\nu) m^2 \left(A \left(\frac{g^4}{f^2} \right) A(f^2) - 1 \right) &\leq A(f^2) A(g^2) - A(f^{2(1-\nu)} g^{2\nu}) A(f^{2\nu} g^{2(1-\nu)}) \quad (2.7) \\ &\leq \nu(1-\nu) M^2 \left(A \left(\frac{g^4}{f^2} \right) A(f^2) - 1 \right). \end{aligned}$$

In particular, if $f^2, g^2, \frac{g^4}{f^2}, fg \in L$ and the condition (2.1) holds, then

$$\begin{aligned} \frac{1}{4} m^2 \left(A \left(\frac{g^4}{f^2} \right) A(f^2) - 1 \right) &\leq A(f^2) A(g^2) - A^2(fg) \quad (2.8) \\ &\leq \frac{1}{4} M^2 \left(A \left(\frac{g^4}{f^2} \right) A(f^2) - 1 \right). \end{aligned}$$

The following result also holds:

Theorem 2.3. Let $A, B : L \rightarrow \mathbb{R}$ be two normalized isotonic functionals. If $f, g : E \rightarrow \mathbb{R}$ are such that $f \geq 0, g > 0, f^2, g^2, \frac{f^4}{g^2}, f^{2(1-\nu)} g^{2\nu}, f^{2\nu} g^{2(1-\nu)} \in L$ for some $\nu \in [0, 1]$ and the condition (2.1) holds, then

$$\begin{aligned} \frac{1}{2M^2} \nu(1-\nu) \left(A(g^2) B \left(\frac{f^4}{g^2} \right) - 2A(f^2) B(f^2) + A \left(\frac{f^4}{g^2} \right) B(g^2) \right) &\leq (1-\nu) A(f^2) B(g^2) + \nu A(g^2) B(f^2) \quad (2.9) \\ &\quad - A(f^{2(1-\nu)} g^{2\nu}) B(f^{2\nu} g^{2(1-\nu)}) \\ &\leq \frac{1}{2m^2} \nu(1-\nu) \\ &\quad \times \left(A(g^2) B \left(\frac{f^4}{g^2} \right) - 2A(f^2) B(f^2) + A \left(\frac{f^4}{g^2} \right) B(g^2) \right). \end{aligned}$$

Proof. For any $x, y \in E$ we have

$$m^2 \leq \frac{f^2(x)}{g^2(x)}, \frac{f^2(y)}{g^2(y)} \leq M^2.$$

If we use the inequality (1.1) for

$$a = \frac{f^2(x)}{g^2(x)}, b = \frac{f^2(y)}{g^2(y)},$$

then we get

$$\begin{aligned} \frac{1}{2M^2} \nu(1-\nu) \left(\frac{f^2(y)}{g^2(y)} - \frac{f^2(x)}{g^2(x)} \right)^2 &\leq (1-\nu) \frac{f^2(x)}{g^2(x)} + \nu \frac{f^2(y)}{g^2(y)} - \left(\frac{f^2(x)}{g^2(x)} \right)^{1-\nu} \left(\frac{f^2(y)}{g^2(y)} \right)^\nu \\ &\leq \frac{1}{2m^2} \nu(1-\nu) \left(\frac{f^2(y)}{g^2(y)} - \frac{f^2(x)}{g^2(x)} \right)^2 \end{aligned}$$

for any $x, y \in E$.

This can be written as

$$\begin{aligned} \frac{1}{2M^2} \nu(1-\nu) \left(\frac{f^4(y)}{g^4(y)} - 2 \frac{f^2(y)}{g^2(y)} \frac{f^2(x)}{g^2(x)} + \frac{f^4(x)}{g^4(x)} \right) &\leq (1-\nu) \frac{f^2(x)}{g^2(x)} + \nu \frac{f^2(y)}{g^2(y)} - \left(\frac{f^2(x)}{g^2(x)} \right)^{1-\nu} \left(\frac{f^2(y)}{g^2(y)} \right)^\nu \quad (2.10) \\ &\leq \frac{1}{2m^2} \nu(1-\nu) \left(\frac{f^4(y)}{g^4(y)} - 2 \frac{f^2(y)}{g^2(y)} \frac{f^2(x)}{g^2(x)} + \frac{f^4(x)}{g^4(x)} \right). \end{aligned}$$

Now, if we multiply (2.10) by $g^2(x)g^2(y) > 0$ then we get

$$\begin{aligned} \frac{1}{2M^2} \nu(1-\nu) \left(\frac{f^4(y)}{g^2(y)} g^2(x) - 2f^2(y)f^2(x) + \frac{f^4(x)}{g^2(x)} g^2(y) \right) &\leq (1-\nu) g^2(y) f^2(x) + \nu f^2(y) g^2(x) \quad (2.11) \\ &\quad - f^{2\nu}(y) g^{2(1-\nu)}(y) f^{2(1-\nu)}(x) g^{2\nu}(x) \\ &\leq \frac{1}{2m^2} \nu(1-\nu) \left(\frac{f^4(y)}{g^2(y)} g^2(x) - 2f^2(y)f^2(x) + \frac{f^4(x)}{g^2(x)} g^2(y) \right) \end{aligned}$$

for any $x, y \in E$.

Fix $y \in E$. Then by (2.11) we have in the order of L that

$$\begin{aligned} \frac{1}{2M^2} \nu(1-\nu) \left(\frac{f^4(y)}{g^2(y)} g^2 - 2f^2(y)f^2 + g^2(y) \frac{f^4}{g^2} \right) &\leq (1-\nu) g^2(y) f^2 + \nu f^2(y) g^2 - f^{2\nu}(y) g^{2(1-\nu)}(y) f^{2(1-\nu)} g^{2\nu} \quad (2.12) \\ &\leq \frac{1}{2m^2} \nu(1-\nu) \left(\frac{f^4(y)}{g^2(y)} g^2 - 2f^2(y)f^2 + g^2(y) \frac{f^4}{g^2} \right). \end{aligned}$$

If we take the functional A in (2.12), then we get

$$\begin{aligned} \frac{1}{2M^2} \nu(1-\nu) \left(\frac{f^4(y)}{g^2(y)} A(g^2) - 2f^2(y)A(f^2) + g^2(y)A\left(\frac{f^4}{g^2}\right) \right) &\leq (1-\nu) g^2(y)A(f^2) + \nu f^2(y)A(g^2) \quad (2.13) \\ &\quad - f^{2\nu}(y) g^{2(1-\nu)}(y)A\left(f^{2(1-\nu)} g^{2\nu}\right) \\ &\leq \frac{1}{2m^2} \nu(1-\nu) \\ &\quad \times \left(\frac{f^4(y)}{g^2(y)} A(g^2) - 2f^2(y)A(f^2) + g^2(y)A\left(\frac{f^4}{g^2}\right) \right) \end{aligned}$$

for any $y \in E$.

This inequality can be written in the order of L as

$$\begin{aligned} \frac{1}{2M^2} \nu(1-\nu) \left(A(g^2) \frac{f^4}{g^2} - 2A(f^2)f^2 + A\left(\frac{f^4}{g^2}\right) g^2 \right) &\leq (1-\nu) A(f^2) g^2 + \nu A(g^2) f^2 - A\left(f^{2(1-\nu)} g^{2\nu}\right) f^{2\nu} g^{2(1-\nu)} \quad (2.14) \\ &\leq \frac{1}{2m^2} \nu(1-\nu) \left(A(g^2) \frac{f^4}{g^2} - 2A(f^2)f^2 + A\left(\frac{f^4}{g^2}\right) g^2 \right). \end{aligned}$$

Now, if we take the functional B in (2.14), then we get the desired result (2.9). $\square$

Corollary 2.4. Let $A : L \rightarrow \mathbb{R}$ be a normalized isotonic functional. If $f, g : E \rightarrow \mathbb{R}$ are such that $f \geq 0, g > 0, f^2, g^2, \frac{f^4}{g^2}, f^{2(1-\nu)} g^{2\nu}, f^{2\nu} g^{2(1-\nu)} \in L$ for some $\nu \in [0, 1]$ and the condition (2.1) is valid, then

$$\begin{aligned} \frac{1}{M^2} \nu(1-\nu) \left(A(g^2) A\left(\frac{f^4}{g^2}\right) - A^2(f^2) \right) &\leq A(f^2) A(g^2) - A\left(f^{2(1-\nu)} g^{2\nu}\right) A\left(f^{2\nu} g^{2(1-\nu)}\right) \quad (2.15) \\ &\leq \frac{1}{m^2} \nu(1-\nu) \left(A(g^2) A\left(\frac{f^4}{g^2}\right) - A^2(f^2) \right). \end{aligned}$$

In particular, if $f^2, g^2, \frac{f^4}{g^2}, fg \in L$ and the condition (2.1) is valid, then we have

$$\begin{aligned} \frac{1}{4M^2} \left(A(g^2) A\left(\frac{f^4}{g^2}\right) - A^2(fg) \right) &\leq A(f^2) A(g^2) - A^2(fg) \\ &\leq \frac{1}{4m^2} \left(A(g^2) A\left(\frac{f^4}{g^2}\right) - A^2(fg) \right). \end{aligned} \quad (2.16)$$

3. Other Related Results

If we write the inequality (1.1) for $a = 1$ and $b = x$ we get

$$\frac{1}{2} \nu(1-\nu) \frac{(x-1)^2}{\max\{x, 1\}} \leq 1 - \nu + \nu x - x^\nu \leq \frac{1}{2} \nu(1-\nu) \frac{(x-1)^2}{\min\{x, 1\}} \quad (3.1)$$

for any $x > 0$ and for any $\nu \in [0, 1]$.

If $x \in [t, T] \subset (0, \infty)$, then $\max\{x, 1\} \leq \max\{T, 1\}$ and $\min\{x, 1\} \leq \min\{t, 1\}$ and by (3.1) we get

$$\begin{aligned} \frac{1}{2} \nu(1-\nu) \frac{\min_{x \in [t, T]} (x-1)^2}{\max\{T, 1\}} &\leq \frac{1}{2} \nu(1-\nu) \frac{(x-1)^2}{\max\{T, 1\}} \\ &\leq 1 - \nu + \nu x - x^\nu \\ &\leq \frac{1}{2} \nu(1-\nu) \frac{(x-1)^2}{\min\{t, 1\}} \\ &\leq \frac{1}{2} \nu(1-\nu) \frac{\max_{x \in [t, T]} (x-1)^2}{\min\{t, 1\}} \end{aligned} \quad (3.2)$$

for any $x \in [t, T]$ and for any $\nu \in [0, 1]$.

Observe that

$$\min_{x \in [t, T]} (x-1)^2 = \begin{cases} (T-1)^2 & \text{if } T < 1, \\ 0 & \text{if } t \leq 1 \leq T, \\ (t-1)^2 & \text{if } 1 < t \end{cases}$$

and

$$\max_{x \in [t, T]} (x-1)^2 = \begin{cases} (t-1)^2 & \text{if } T < 1, \\ \max\{(t-1)^2, (T-1)^2\} & \text{if } t \leq 1 \leq T, \\ (T-1)^2 & \text{if } 1 < t. \end{cases}$$

Then

$$c(t, T) := \frac{\min_{x \in [t, T]} (x-1)^2}{\max\{T, 1\}} = \begin{cases} (T-1)^2 & \text{if } T < 1, \\ 0 & \text{if } t \leq 1 \leq T, \\ \frac{(t-1)^2}{T} & \text{if } 1 < t \end{cases} \quad (3.3)$$

and

$$C(t, T) := \frac{\max_{x \in [t, T]} (x-1)^2}{\min\{t, 1\}} = \begin{cases} \frac{(t-1)^2}{t} & \text{if } T < 1, \\ \frac{1}{t} \max\{(t-1)^2, (T-1)^2\} & \text{if } t \leq 1 \leq T, \\ (T-1)^2 & \text{if } 1 < t. \end{cases} \quad (3.4)$$

Using the inequality (3.2) we have

$$\begin{aligned} \frac{1}{2} \nu(1-\nu) c(t, T) &\leq \frac{1}{2} \nu(1-\nu) \frac{(x-1)^2}{\max\{T, 1\}} \leq 1 - \nu + \nu x - x^\nu \\ &\leq \frac{1}{2} \nu(1-\nu) \frac{(x-1)^2}{\min\{t, 1\}} \leq \frac{1}{2} \nu(1-\nu) C(t, T) \end{aligned} \quad (3.5)$$

for any $x \in [t, T]$ and for any $\nu \in [0, 1]$.

Now, if $a, b > 0$ and assume that $\frac{b}{a} \in [t, T]$, then by (3.5) we get

$$\begin{aligned} \frac{1}{2} \nu(1-\nu) c(t, T) a &\leq \frac{1}{2} \nu(1-\nu) \frac{(b-a)^2}{\max\{T, 1\} a} \\ &\leq (1-\nu) a + \nu b - b^\nu a^{1-\nu} \\ &\leq \frac{1}{2} \nu(1-\nu) \frac{(b-a)^2}{\min\{t, 1\} a} \leq \frac{1}{2} \nu(1-\nu) C(t, T) a \end{aligned} \quad (3.6)$$

for any $\nu \in [0, 1]$, where $c(t, T)$ and $C(t, T)$ are defined by (3.3) and (3.4), respectively.

Theorem 3.1. Let $A, B : L \rightarrow \mathbb{R}$ be two normalized isotonic functionals. If $f, g : E \rightarrow \mathbb{R}$ are such that, $f^2, g^2, \frac{g^4}{f^2}, \frac{f^4}{g^2}, f^{2(1-\nu)}g^{2\nu}, f^{2\nu}g^{2(1-\nu)} \in L$ for some $\nu \in [0, 1]$ and the condition (2.1) holds, then

$$\begin{aligned} 0 &\leq \frac{1}{2}\nu(1-\nu)\frac{m^2}{M^2}\left(A\left(\frac{g^4}{f^2}\right)B\left(\frac{f^4}{g^2}\right)-2A(g^2)B(f^2)+A(f^2)B(g^2)\right) \\ &\leq (1-\nu)A(f^2)B(g^2)+\nu A(g^2)B(f^2)-A(f^{2(1-\nu)}g^{2\nu})B(f^{2\nu}g^{2(1-\nu)}) \\ &\leq \frac{1}{2}\nu(1-\nu)\frac{M^2}{m^2}\left(A\left(\frac{g^4}{f^2}\right)B\left(\frac{f^4}{g^2}\right)-2A(g^2)B(f^2)+A(f^2)B(g^2)\right) \\ &\leq \frac{1}{2}\nu(1-\nu)\frac{M^2}{m^2}\left(\frac{M^2}{m^2}-1\right)^2A(f^2)B(g^2). \end{aligned} \quad (3.7)$$

Proof. For any $x, y \in E$ we have

$$m^2 \leq \frac{f^2(x)}{g^2(x)}, \frac{f^2(y)}{g^2(y)} \leq M^2.$$

Consider

$$a = \frac{f^2(x)}{g^2(x)}, \quad b = \frac{f^2(y)}{g^2(y)},$$

then $\frac{b}{a} \in \left[\frac{m^2}{M^2}, \frac{M^2}{m^2}\right]$ and by (3.6) we get

$$\begin{aligned} 0 &\leq \frac{1}{2}\nu(1-\nu)\frac{\left(\frac{f^2(y)}{g^2(y)}-\frac{f^2(x)}{g^2(x)}\right)^2}{\frac{M^2}{m^2}\frac{f^2(x)}{g^2(x)}} \\ &\leq (1-\nu)\frac{f^2(x)}{g^2(x)}+\nu\frac{f^2(y)}{g^2(y)}-\left(\frac{f^2(y)}{g^2(y)}\right)^\nu\left(\frac{f^2(x)}{g^2(x)}\right)^{1-\nu} \\ &\leq \frac{1}{2}\nu(1-\nu)\frac{\left(\frac{f^2(y)}{g^2(y)}-\frac{f^2(x)}{g^2(x)}\right)^2}{\frac{m^2}{M^2}\frac{f^2(x)}{g^2(x)}} \\ &\leq \frac{1}{2}\nu(1-\nu)\frac{M^2}{m^2}\max\left\{\left(\frac{m^2}{M^2}-1\right)^2,\left(\frac{M^2}{m^2}-1\right)^2\right\}\frac{f^2(x)}{g^2(x)} \\ &= \frac{1}{2}\nu(1-\nu)\frac{M^2}{m^2}\left(\frac{M^2}{m^2}-1\right)^2\frac{f^2(x)}{g^2(x)} \end{aligned}$$

for any $x, y \in E$ and $\nu \in [0, 1]$.

This inequality is equivalent to

$$\begin{aligned} 0 &\leq \frac{1}{2}\nu(1-\nu)\frac{m^2}{M^2}\frac{\left(\frac{f^2(y)}{g^2(y)}-\frac{f^2(x)}{g^2(x)}\right)^2}{f^2(x)}g^2(x) \\ &\leq (1-\nu)\frac{f^2(x)}{g^2(x)}+\nu\frac{f^2(y)}{g^2(y)}-\left(\frac{f^2(y)}{g^2(y)}\right)^\nu\left(\frac{f^2(x)}{g^2(x)}\right)^{1-\nu} \\ &\leq \frac{1}{2}\nu(1-\nu)\frac{M^2}{m^2}\frac{\left(\frac{f^2(y)}{g^2(y)}-\frac{f^2(x)}{g^2(x)}\right)^2}{f^2(x)}g^2(x) \\ &\leq \frac{1}{2}\nu(1-\nu)\frac{M^2}{m^2}\left(\frac{M^2}{m^2}-1\right)^2\frac{f^2(x)}{g^2(x)} \end{aligned} \quad (3.8)$$

for any $x, y \in E$ and $\nu \in [0, 1]$.

Now, if we multiply (3.8) by $g^2(x)g^2(y) > 0$ then we get

$$\begin{aligned}
 0 &\leq \frac{1}{2} \nu(1-\nu) \frac{m^2}{M^2} \frac{\left(\frac{f^2(y)}{g^2(y)} - \frac{f^2(x)}{g^2(x)}\right)^2 g^4(x) g^2(y)}{f^2(x)} \\
 &\leq (1-\nu) g^2(y) f^2(x) + \nu f^2(y) g^2(x) - f^{2\nu}(y) g^{2(1-\nu)}(y) f^{2(1-\nu)}(x) g^{2\nu}(x) \\
 &\leq \frac{1}{2} \nu(1-\nu) \frac{M^2}{m^2} \frac{\left(\frac{f^2(y)}{g^2(y)} - \frac{f^2(x)}{g^2(x)}\right)^2 g^4(x) g^2(y)}{f^2(x)} \\
 &\leq \frac{1}{2} \nu(1-\nu) \frac{M^2}{m^2} \left(\frac{M^2}{m^2} - 1\right)^2 f^2(x) g^2(y)
 \end{aligned} \tag{3.9}$$

for any $x, y \in E$ and $\nu \in [0, 1]$.

Observe that

$$\begin{aligned}
 \frac{\left(\frac{f^2(y)}{g^2(y)} - \frac{f^2(x)}{g^2(x)}\right)^2 g^4(x) g^2(y)}{f^2(x)} &= \frac{\left(\frac{f^4(y)}{g^4(y)} - 2\frac{f^2(y)}{g^2(y)}\frac{f^2(x)}{g^2(x)} + \frac{f^4(x)}{g^4(x)}\right) g^4(x) g^2(y)}{f^2(x)} \\
 &= \frac{\frac{f^4(y)g^4(x)}{g^2(y)} - 2f^2(y)f^2(x)g^2(x) + f^4(x)g^2(y)}{f^2(x)} \\
 &= \frac{f^4(y)g^4(x)}{g^2(y)f^2(x)} - 2f^2(y)g^2(x) + f^2(x)g^2(y)
 \end{aligned}$$

and by (3.9) we get

$$\begin{aligned}
 0 &\leq \frac{1}{2} \nu(1-\nu) \frac{m^2}{M^2} \left(\frac{f^4(y)g^4(x)}{g^2(y)f^2(x)} - 2f^2(y)g^2(x) + f^2(x)g^2(y)\right) \\
 &\leq (1-\nu) g^2(y) f^2(x) + \nu f^2(y) g^2(x) - f^{2\nu}(y) g^{2(1-\nu)}(y) f^{2(1-\nu)}(x) g^{2\nu}(x) \\
 &\leq \frac{1}{2} \nu(1-\nu) \frac{M^2}{m^2} \left(\frac{f^4(y)g^4(x)}{g^2(y)f^2(x)} - 2f^2(y)g^2(x) + f^2(x)g^2(y)\right) \\
 &\leq \frac{1}{2} \nu(1-\nu) \frac{M^2}{m^2} \left(\frac{M^2}{m^2} - 1\right)^2 f^2(x) g^2(y)
 \end{aligned}$$

for any $x, y \in E$ and $\nu \in [0, 1]$.

Now, if we use a similar argument to the one from the proof of Theorem 2.1 we deduce the desired result (3.7). $\square$

Corollary 3.2. Let $A : L \rightarrow \mathbb{R}$ be a normalized isotonic functional. If $f, g : E \rightarrow \mathbb{R}$ are such that, $f^2, g^2, \frac{g^4}{f^2}, \frac{f^4}{g^2}, f^{2(1-\nu)}g^{2\nu}, f^{2\nu}g^{2(1-\nu)} \in L$ for some $\nu \in [0, 1]$ and the condition (2.1) holds, then

$$\begin{aligned}
 0 &\leq \frac{1}{2} \nu(1-\nu) \frac{m^2}{M^2} \left(A\left(\frac{g^4}{f^2}\right)A\left(\frac{f^4}{g^2}\right) - A(g^2)A(f^2)\right) \\
 &\leq A(f^2)A(g^2) - A\left(f^{2(1-\nu)}g^{2\nu}\right)A\left(f^{2\nu}g^{2(1-\nu)}\right) \\
 &\leq \frac{1}{2} \nu(1-\nu) \frac{M^2}{m^2} \left(A\left(\frac{g^4}{f^2}\right)A\left(\frac{f^4}{g^2}\right) - A(g^2)A(f^2)\right) \\
 &\leq \frac{1}{2} \nu(1-\nu) \frac{M^2}{m^2} \left(\frac{M^2}{m^2} - 1\right)^2 A(f^2)A(g^2).
 \end{aligned} \tag{3.10}$$

In particular, if $f^2, g^2, \frac{g^4}{f^2}, \frac{f^4}{g^2}, fg \in L$, then we have

$$\begin{aligned}
 0 &\leq \frac{1}{8} \frac{m^2}{M^2} \left(A\left(\frac{g^4}{f^2}\right)A\left(\frac{f^4}{g^2}\right) - A(g^2)A(f^2)\right) \\
 &\leq A(f^2)A(g^2) - A^2(fg) \\
 &\leq \frac{1}{8} \frac{M^2}{m^2} \left(A\left(\frac{g^4}{f^2}\right)A\left(\frac{f^4}{g^2}\right) - A(g^2)A(f^2)\right) \\
 &\leq \frac{1}{8} \frac{M^2}{m^2} \left(\frac{M^2}{m^2} - 1\right)^2 A(f^2)A(g^2).
 \end{aligned} \tag{3.11}$$

We observe that the inequality (3.11) can be written as

$$\begin{aligned} 0 &\leq \frac{1}{8} \frac{m^2}{M^2} \left(\frac{A\left(\frac{g^4}{f^2}\right) A\left(\frac{f^4}{g^2}\right)}{A(g^2) A(f^2)} - 1 \right) \leq 1 - \frac{A^2(fg)}{A(f^2) A(g^2)} \\ &\leq \frac{1}{8} \frac{M^2}{m^2} \left(\frac{A\left(\frac{g^4}{f^2}\right) A\left(\frac{f^4}{g^2}\right)}{A(g^2) A(f^2)} - 1 \right) \leq \frac{1}{8} \frac{M^2}{m^2} \left(\frac{M^2}{m^2} - 1 \right)^2. \end{aligned} \quad (3.12)$$

4. Applications for Integrals

Let $(\Omega, \mathcal{A}, \mu)$ be a measurable space consisting of a set Ω , a σ -algebra $\mathcal{A}$ of subsets of Ω and a countably additive and positive measure μ on $\mathcal{A}$ with values in $\mathbb{R} \cup \{\infty\}$. For a μ -measurable function $w : \Omega \rightarrow \mathbb{R}$, with $w(x) \geq 0$ for μ -a.e. (almost every) $x \in \Omega$ and $p \geq 1$ consider the Lebesgue space

$$L_w^p(\Omega, \mu) := \{f : \Omega \rightarrow \mathbb{R}, f \text{ is } \mu\text{-measurable and } \int_{\Omega} |f(x)|^p w(x) d\mu(x) < \infty\}.$$

For simplicity of notation we write everywhere in the sequel $\int_{\Omega} w d\mu$ instead of $\int_{\Omega} w(x) d\mu(x)$. The same for other integrals involved below. We assume that $\int_{\Omega} w d\mu = 1$.

Let f, g be μ -measurable functions with the property that there exists the constants $M, m > 0$ such that

$$0 < m \leq \frac{f}{g} \leq M < \infty \text{ } \mu\text{-almost everywhere (a.e.) on } \Omega. \quad (4.1)$$

If $f^2, g^2, \frac{g^4}{f^2}, f^{2(1-\nu)} g^{2\nu}, f^{2\nu} g^{2(1-\nu)} \in L_w(\Omega, \mu)$ for some $\nu \in [0, 1]$ and the condition (4.1) holds, then by (2.7) we have

$$\begin{aligned} \nu(1-\nu)m^2 \left(\int_{\Omega} w \frac{g^4}{f^2} d\mu \int_{\Omega} w f^2 d\mu - 1 \right) &\leq \int_{\Omega} w f^2 d\mu \int_{\Omega} w g^2 d\mu - \int_{\Omega} w f^{2(1-\nu)} g^{2\nu} d\mu \int_{\Omega} w f^{2\nu} g^{2(1-\nu)} d\mu \\ &\leq \nu(1-\nu)M^2 \left(\int_{\Omega} w \frac{g^4}{f^2} d\mu \int_{\Omega} w f^2 d\mu - 1 \right). \end{aligned} \quad (4.2)$$

In particular, if $f^2, g^2, \frac{g^4}{f^2}, fg \in L_w(\Omega, \mu)$ and the condition (4.1) holds, then

$$\begin{aligned} \frac{1}{4}m^2 \left(\int_{\Omega} w \frac{g^4}{f^2} d\mu \int_{\Omega} w f^2 d\mu - 1 \right) &\leq \int_{\Omega} w f^2 d\mu \int_{\Omega} w g^2 d\mu - \left(\int_{\Omega} w f g d\mu \right)^2 \\ &\leq \frac{1}{4}M^2 \left(\int_{\Omega} w \frac{g^4}{f^2} d\mu \int_{\Omega} w f^2 d\mu - 1 \right). \end{aligned} \quad (4.3)$$

If $f^2, g^2, \frac{f^4}{g^2}, f^{2(1-\nu)} g^{2\nu}, f^{2\nu} g^{2(1-\nu)} \in L_w(\Omega, \mu)$ for some $\nu \in [0, 1]$ and the condition (4.1) holds, then by (2.15) we have

$$\begin{aligned} \frac{1}{M^2} \nu(1-\nu) \left(\int_{\Omega} w g^2 d\mu \int_{\Omega} w \frac{f^4}{g^2} d\mu - \left(\int_{\Omega} w f^2 d\mu \right)^2 \right) &\leq \int_{\Omega} w f^2 d\mu \int_{\Omega} w g^2 d\mu - \int_{\Omega} w f^{2(1-\nu)} g^{2\nu} d\mu \int_{\Omega} w f^{2\nu} g^{2(1-\nu)} d\mu \\ &\leq \frac{1}{m^2} \nu(1-\nu) \left(\int_{\Omega} w g^2 d\mu \int_{\Omega} w \frac{f^4}{g^2} d\mu - \left(\int_{\Omega} w f^2 d\mu \right)^2 \right). \end{aligned} \quad (4.4)$$

In particular, if $f^2, g^2, \frac{f^4}{g^2}, fg \in L_w(\Omega, \mu)$ and the condition (4.1) is valid, then we have

$$\begin{aligned} \frac{1}{4M^2} \left(\int_{\Omega} w g^2 d\mu \int_{\Omega} w \frac{f^4}{g^2} d\mu - \left(\int_{\Omega} w f^2 d\mu \right)^2 \right) &\leq \int_{\Omega} w f^2 d\mu \int_{\Omega} w g^2 d\mu - \left(\int_{\Omega} w f g d\mu \right)^2 \\ &\leq \frac{1}{4m^2} \left(\int_{\Omega} w g^2 d\mu \int_{\Omega} w \frac{f^4}{g^2} d\mu - \left(\int_{\Omega} w f^2 d\mu \right)^2 \right). \end{aligned} \quad (4.5)$$

If $f^2, g^2, \frac{g^4}{f^2}, \frac{f^4}{g^2}, f^{2(1-\nu)}g^{2\nu}, f^{2\nu}g^{2(1-\nu)} \in L_w(\Omega, \mu)$ for some $\nu \in [0, 1]$, and the condition (4.1) holds, then

$$\begin{aligned} 0 &\leq \frac{1}{2} \nu (1-\nu) \frac{m^2}{M^2} \left(\int_{\Omega} w \frac{g^4}{f^2} d\mu \int_{\Omega} w \frac{f^4}{g^2} d\mu - \int_{\Omega} w g^2 d\mu \int_{\Omega} w f^2 d\mu \right) \\ &\leq \int_{\Omega} w f^2 d\mu \int_{\Omega} w g^2 d\mu - \int_{\Omega} w f^{2(1-\nu)} g^{2\nu} d\mu \int_{\Omega} w f^{2\nu} g^{2(1-\nu)} d\mu \\ &\leq \frac{1}{2} \nu (1-\nu) \frac{M^2}{m^2} \left(\int_{\Omega} w \frac{g^4}{f^2} d\mu \int_{\Omega} w \frac{f^4}{g^2} d\mu - \int_{\Omega} w g^2 d\mu \int_{\Omega} w f^2 d\mu \right) \\ &\leq \frac{1}{2} \nu (1-\nu) \frac{M^2}{m^2} \left(\frac{M^2}{m^2} - 1 \right)^2 \int_{\Omega} w f^2 d\mu \int_{\Omega} w g^2 d\mu. \end{aligned} \quad (4.6)$$

In particular, if $f^2, g^2, \frac{g^4}{f^2}, \frac{f^4}{g^2}, fg \in L_w(\Omega, \mu)$, then we have

$$\begin{aligned} 0 &\leq \frac{1}{8} \frac{m^2}{M^2} \left(\int_{\Omega} w \frac{g^4}{f^2} d\mu \int_{\Omega} w \frac{f^4}{g^2} d\mu - \int_{\Omega} w g^2 d\mu \int_{\Omega} w f^2 d\mu \right) \\ &\leq \int_{\Omega} w f^2 d\mu \int_{\Omega} w g^2 d\mu - \left(\int_{\Omega} w f g d\mu \right)^2 \\ &\leq \frac{1}{8} \frac{M^2}{m^2} \left(\int_{\Omega} w \frac{g^4}{f^2} d\mu \int_{\Omega} w \frac{f^4}{g^2} d\mu - \int_{\Omega} w g^2 d\mu \int_{\Omega} w f^2 d\mu \right) \\ &\leq \frac{1}{8} \frac{M^2}{m^2} \left(\frac{M^2}{m^2} - 1 \right)^2 \int_{\Omega} w f^2 d\mu \int_{\Omega} w g^2 d\mu. \end{aligned} \quad (4.7)$$

5. Applications for Real Numbers

We consider the n -tuples of positive numbers $a = (a_1, \dots, a_n)$, $b = (b_1, \dots, b_n)$ and the probability distribution $p = (p_1, \dots, p_n)$, i.e. $p_i \geq 0$ for any $i \in \{1, \dots, n\}$ with $\sum_{i=1}^n p_i = 1$. If there exist the constants $m, M > 0$ such that

$$0 < m \leq \frac{a_i}{b_i} \leq M < \infty \text{ for any } i \in \{1, \dots, n\}, \quad (5.1)$$

then by (4.2) and (4.3) for the counting discrete measure, we have

$$\begin{aligned} \nu(1-\nu)m^2 \left(\sum_{i=1}^n p_i \frac{b_i^4}{a_i^2} \sum_{i=1}^n p_i a_i^2 - 1 \right) &\leq \sum_{i=1}^n p_i a_i^2 \sum_{i=1}^n p_i b_i^2 - \sum_{i=1}^n p_i a_i^{2(1-\nu)} b_i^{2\nu} \sum_{i=1}^n p_i a_i^{2\nu} b_i^{2(1-\nu)} \\ &\leq \nu(1-\nu)M^2 \left(\sum_{i=1}^n p_i \frac{b_i^4}{a_i^2} \sum_{i=1}^n p_i a_i^2 - 1 \right) \end{aligned} \quad (5.2)$$

for any $\nu \in [0, 1]$ and

$$\begin{aligned} \frac{1}{4} m^2 \left(\sum_{i=1}^n p_i \frac{b_i^4}{a_i^2} \sum_{i=1}^n p_i a_i^2 - 1 \right) &\leq \sum_{i=1}^n p_i a_i^2 \sum_{i=1}^n p_i b_i^2 - \left(\sum_{i=1}^n p_i a_i b_i \right)^2 \\ &\leq \frac{1}{4} M^2 \left(\sum_{i=1}^n p_i \frac{b_i^4}{a_i^2} \sum_{i=1}^n p_i a_i^2 - 1 \right). \end{aligned} \quad (5.3)$$

If $a = (a_1, \dots, a_n)$, $b = (b_1, \dots, b_n)$ satisfy (5.1), then by (4.4) and (4.5) for the counting discrete measure, we have

$$\begin{aligned} \frac{1}{M^2} \nu(1-\nu) \left(\sum_{i=1}^n p_i b_i^2 \sum_{i=1}^n p_i \frac{a_i^4}{b_i^2} - \left(\sum_{i=1}^n p_i a_i^2 \right)^2 \right) &\leq \sum_{i=1}^n p_i a_i^2 \sum_{i=1}^n p_i b_i^2 - \sum_{i=1}^n p_i a_i^{2(1-\nu)} b_i^{2\nu} \sum_{i=1}^n p_i a_i^{2\nu} b_i^{2(1-\nu)} \\ &\leq \frac{1}{m^2} \nu(1-\nu) \left(\sum_{i=1}^n p_i b_i^2 \sum_{i=1}^n p_i \frac{a_i^4}{b_i^2} - \left(\sum_{i=1}^n p_i a_i^2 \right)^2 \right) \end{aligned} \quad (5.4)$$

for any $\nu \in [0, 1]$ and

$$\begin{aligned} \frac{1}{4M^2} \left(\sum_{i=1}^n p_i b_i^2 \sum_{i=1}^n p_i \frac{a_i^4}{b_i^2} - \left(\sum_{i=1}^n p_i a_i^2 \right)^2 \right) &\leq \sum_{i=1}^n p_i a_i^2 \sum_{i=1}^n p_i b_i^2 - \left(\sum_{i=1}^n p_i a_i b_i \right)^2 \\ &\leq \frac{1}{4m^2} \left(\sum_{i=1}^n p_i b_i^2 \sum_{i=1}^n p_i \frac{a_i^4}{b_i^2} - \left(\sum_{i=1}^n p_i a_i^2 \right)^2 \right). \end{aligned} \quad (5.5)$$

If $a = (a_1, \dots, a_n)$, $b = (b_1, \dots, b_n)$ satisfy (5.1), then by (4.6) and (4.7) for the counting discrete measure, we have

$$\begin{aligned}
 0 &\leq \frac{1}{2} \nu(1-\nu) \frac{m^2}{M^2} \left(\sum_{i=1}^n p_i \frac{b_i^4}{a_i^2} \sum_{i=1}^n p_i \frac{a_i^4}{b_i^2} - \sum_{i=1}^n p_i b_i^2 \sum_{i=1}^n p_i a_i^2 \right) \\
 &\leq \sum_{i=1}^n p_i a_i^2 \sum_{i=1}^n p_i b_i^2 - \sum_{i=1}^n p_i a_i^{2(1-\nu)} b_i^{2\nu} \sum_{i=1}^n p_i a_i^{2\nu} b_i^{2(1-\nu)} \\
 &\leq \frac{1}{2} \nu(1-\nu) \frac{M^2}{m^2} \left(\sum_{i=1}^n p_i \frac{b_i^4}{a_i^2} \sum_{i=1}^n p_i \frac{a_i^4}{b_i^2} - \sum_{i=1}^n p_i b_i^2 \sum_{i=1}^n p_i a_i^2 \right) \\
 &\leq \frac{1}{2} \nu(1-\nu) \frac{M^2}{m^2} \left(\frac{M^2}{m^2} - 1 \right)^2 \sum_{i=1}^n p_i a_i^2 \sum_{i=1}^n p_i b_i^2
 \end{aligned} \tag{5.6}$$

for any $\nu \in [0, 1]$ and

$$\begin{aligned}
 0 &\leq \frac{1}{8} \frac{m^2}{M^2} \left(\sum_{i=1}^n p_i \frac{b_i^4}{a_i^2} \sum_{i=1}^n p_i \frac{a_i^4}{b_i^2} - \sum_{i=1}^n p_i b_i^2 \sum_{i=1}^n p_i a_i^2 \right) \\
 &\leq \sum_{i=1}^n p_i a_i^2 \sum_{i=1}^n p_i b_i^2 - \left(\sum_{i=1}^n p_i a_i b_i \right)^2 \\
 &\leq \frac{1}{8} \frac{M^2}{m^2} \left(\sum_{i=1}^n p_i \frac{b_i^4}{a_i^2} \sum_{i=1}^n p_i \frac{a_i^4}{b_i^2} - \sum_{i=1}^n p_i b_i^2 \sum_{i=1}^n p_i a_i^2 \right) \\
 &\leq \frac{1}{8} \frac{M^2}{m^2} \left(\frac{M^2}{m^2} - 1 \right)^2 \sum_{i=1}^n p_i a_i^2 \sum_{i=1}^n p_i b_i^2.
 \end{aligned} \tag{5.7}$$

6. Conclusion

In this paper, by making use of some reverses and refinements of Young's inequality (1.1), we obtained some inequalities for isotonic functionals that are related to the second part of Callebaut's inequality (1.2). Natural applications for integrals and n -tuples of real numbers were also provided.

Declarations

Acknowledgements: The author would like to express their sincere thanks to the editor and the anonymous reviewers for their helpful comments and suggestions.

Conflict of Interest Disclosure: The author declares no conflict of interest.

Copyright Statement: Author own the copyright of their work published in the journal and their work is published under the CC BY-NC 4.0 license.

Supporting/Supporting Organizations: This research received no external funding.

Ethical Approval and Participant Consent: This article does not contain any studies with human or animal subjects. It is declared that during the preparation process of this study, scientific and ethical principles were followed and all the studies benefited from are stated in the bibliography.

Plagiarism Statement: This article was scanned by the plagiarism program. No plagiarism detected.

Availability of Data and Materials: Data sharing not applicable.

ORCID

Sever Silvestru Dragomir  <https://orcid.org/0000-0003-2902-6805>

References

- [1] P.R. Beesack and J.E. Pečarić, *On Jessen's inequality for convex functions*, J. Math. Anal. Appl. **110** (1985), 536-552. [\[CrossRef\]](#) [\[Scopus\]](#)
- [2] J.E. Pečarić, *On Jessen's inequality for convex functions (III)*, J. Math. Anal. Appl., **156** (1991), 231-239. [\[CrossRef\]](#) [\[Scopus\]](#)
- [3] J.E. Pečarić and P.R. Beesack, *On Jessen's inequality for convex functions (II)*, J. Math. Anal. Appl., **156** (1991), 231-239. [\[CrossRef\]](#) [\[Scopus\]](#)
- [4] D. Andrica and C. Badea, *Grüss' inequality for positive linear functionals*, Periodica Math. Hung., **19** (1998), 155-167. [\[CrossRef\]](#) [\[Scopus\]](#)
- [5] S.S. Dragomir, *A refinement of Hadamard's inequality for isotonic linear functionals*, Tamkang J. Math., **24**(1) (1992), 101-106.
- [6] S.S. Dragomir, *On a reverse of Jessen's inequality for isotonic linear functionals*, J. Ineq. Pure & Appl. Math., **2**(3)(2001), Article 36.
- [7] S.S. Dragomir, *On the Jessen's inequality for isotonic linear functionals*, Nonlinear Anal. Forum, **7**(2)(2002), 139-151.

- [8] S.S. Dragomir, *On the Lupaş-Beesack-Pečarić inequality for isotonic linear functionals*, Nonlinear Funct. Anal. & Appl., **7**(2)(2002), 285-298.
- [9] S.S. Dragomir, *Bounds for the normalized Jensen functional*, Bull. Austral. Math. Soc. **74**(3)(2006), 417-478. [CrossRef] [Scopus]
- [10] S.S. Dragomir and N.M. Ionescu, *On some inequalities for convex-dominated functions*, L'Anal. Num. Théor. L'Approx., **19**(1) (1990), 21-27.
- [11] S.S. Dragomir and C.E.M. Pearce, *Selected Topics on Hermite-Hadamard Inequalities and Applications*, RGMIA Monographs, Victoria University, 2000. <http://rgmia.vu.edu.au/monographs.html>
- [12] S.S. Dragomir, C.E.M. Pearce and J.E. Pečarić, *On Jessen's and related inequalities for isotonic sublinear functionals*, Acta. Sci. Math., **61** (1995), 373-382.
- [13] F. Kittaneh and Y. Manasrah, *Improved Young and Heinz inequalities for matrix*, J. Math. Anal. Appl., **361**(1) (2010), 262-269 [CrossRef] [Scopus] [Web of Science]
- [14] F. Kittaneh and Y. Manasrah, *Reverse Young and Heinz inequalities for matrices*, Linear Multilinear Algebra., **59**(9) (2011), 1031-1037. [CrossRef] [Scopus] [Web of Science]
- [15] A. Lupaş, *A generalisation of Hadamard's inequalities for convex functions*, Univ. Beograd. Elek. Fak., 577-579 (1976), 115-121.
- [16] D.I. Cartwright and M.J. Field, *A refinement of the arithmetic mean-geometric mean inequality*, Proc. Amer. Math. Soc., **71** (1978), 36-38. [CrossRef]
- [17] D.K. Callebaut, *Generalization of Cauchy-Schwarz inequality*, J. Math. Anal. Appl. **12** (1965), 491-494. [CrossRef]

Fundamental Journal of Mathematics and Applications (FUJMA), (Fundam. J. Math. Appl.)

<https://dergipark.org.tr/en/pub/fujma>



All open access articles published are distributed under the terms of the CC BY-NC 4.0 license (Creative Commons Attribution-Non-Commercial 4.0 International Public License as currently displayed at <http://creativecommons.org/licenses/by-nc/4.0/legalcode>) which permits unrestricted use, distribution, and reproduction in any medium, for non-commercial purposes, provided the original work is properly cited.

How to cite this article: S.S. Dragomir, *Some refinements and reverses of Callebaut's inequality for isotonic functionals via a result due to Cartwright and Field*, Fundam. J. Math. Appl., **7**(1) (2024), 1-11. DOI 10.33401/fujma.1362681