

## PAPER DETAILS

TITLE: A General Fixed Point Theorem for Pairs of Mappings Satisfying a  $\phi$ -Implicit Relation in G - Metric Spaces

AUTHORS: Valeriu POPA, Alina Mihaela PATRICIU

PAGES: 1031-1038

ORIGINAL PDF URL: <https://dergipark.org.tr/tr/download/article-file/83654>



# A General Fixed Point Theorem for Pairs of Mappings Satisfying a $\phi$ - Implicit Relation in $G$ - Metric Spaces

Valeriu POPA<sup>1,\*</sup>, Alina-Mihaela PATRICIU<sup>1</sup>

<sup>1</sup>Vasile Alecsandri University of Bacău, Department of Mathematics, Informatics and Educational Sciences, Romania

Received: 03.07.2012 Revised: 27.03.2013 Accepted: 26.07.2013

## ABSTRACT

In this paper a general fixed point theorem for two pairs of weakly compatible mappings satisfying  $\phi$  - implicit relations in  $G$  - metric spaces, which generalize and improve Theorem 2.1 [9], is proved.

**Key Words:**  $G$  - metric spaces, common fixed point, weakly compatible mappings,  $\phi$  - implicit relation.

## 1. INTRODUCTION

Let  $(X, d)$  be a metric space and  $S, T : (X, d) \rightarrow (X, d)$  be two mappings. In 1994, Pant [17] introduced the notion of pointwise  $R$  - weakly commuting mappings. It is proved in [18] that pointwise  $R$  - weakly commutativity is equivalent to commutativity in coincidence points. Jungck [8] defined  $S$  and  $T$  to be weakly compatible if  $Sx = Tx$  implies  $STx = TStx$ . Thus,  $S$  and  $T$  are weakly compatible if and only if  $S$  and  $T$  are pointwise  $R$  - weakly commuting.

In [5], [6], Dhage introduced a new class of generalized metric space, named  $D$  - metric spaces. Mustafa and Sims [11], [12] proved that most of the claims concerning the fundamental topological structures on  $D$  - metric

spaces are incorrect and introduced an appropriate notion of generalized metric space, named  $G$  - metric space. In fact, Mustafa, Sims and other authors studied many fixed point results for self mappings in  $G$  - metric spaces under certain conditions [13], [14], [15], [16], [28] and other papers.

Several classical fixed point theorems and common fixed point theorems have been recently unified by considering a general condition by an implicit relation in [19], [20] and other papers. Actually, the method is used in the study of fixed points in metric spaces, symmetric spaces, quasi - metric spaces, ultra - metric spaces, convex metric spaces, reflexive spaces, compact metric spaces, paracompact metric spaces, in two or three metric spaces, for single valued mappings, hybrid pairs of mappings and set valued mappings. Quite recently, this method is used in the study of fixed points for mappings satisfying an contractive condition of integral type, in fuzzy metric spaces, probabilistic metric spaces and intuitionistic metric spaces.

\*Corresponding author, e-mail: vpopa@ub.ro

The notion of  $\phi$  - implicit relation is introduced in [4]. The study of fixed points for mappings satisfying implicit relations in  $G$  - metric spaces is initiated in [21], [22], [24], [26] and in other papers.

The study of fixed points for mappings satisfying a  $\phi$  - contractive implicit relation in  $G$  - metric spaces is initiated in [23], [25]. In [23], the authors proved a fixed point theorem for a mapping satisfying a  $\phi$  - implicit relation. Quite recently, in [27] the authors extended this results for a pair of mappings. In [9], a common fixed point theorem for two pairs of weakly compatible mappings in  $G$  - metric spaces is proved.

In this paper a general fixed point theorem for two pairs of weakly compatible mappings in  $G$  - metric spaces, satisfying  $\phi$  - implicit relations which generalizes and improves the results from Theorem 2.1 [9].

## 2. PRELIMINARIES

Definition 2.1 [12] Let  $X$  be a nonempty set and  $G : X^3 \rightarrow P_+$  be a function satisfying the following properties:

- $(G_1) : G(x, y, z) = 0$  if  $x = y = z$ ,
- $(G_2) : 0 < G(x, x, y)$  for all  $x, y \in X$  with  $x \neq y$ ,
- $(G_3) : G(x, x, y) \leq G(x, y, z)$  for all  $x, y, z \in X$  with  $z \neq y$ ,
- $(G_4) : G(x, y, z) = G(y, z, x) = \dots$  (symmetry in all three variables),
- $(G_5) : G(x, y, z) \leq G(x, a, a) + G(a, y, z)$  for all  $x, y, z, a \in X$  (rectangle inequality).

The function  $G$  is called a  $G$  - metric on  $X$  and the pair  $(X, G)$  is called a  $G$  - metric space.

Note that if  $G(x, y, z) = 0$  then  $x = y = z$ .

Definition 2.2 [12] Let  $(X, G)$  be a  $G$  - metric space.

A sequence  $(x_n)$  in  $(X, G)$  is said to be:

- a)  $G$  - convergent if for  $\varepsilon > 0$ , there is an  $x \in X$  and  $k \in \mathbb{N}$  such that for all  $n, m \in \mathbb{N}, n, m \geq k$ ,  $G(x, x_n, x_m) < \varepsilon$ .
- b)  $G$  - Cauchy if for  $\varepsilon > 0$ , there is  $k \in \mathbb{N}$  such that for all  $n, m, p \in \mathbb{N}$ , with  $n, m, p \geq k$ ,  $G(x_n, x_m, x_p) < \varepsilon$ , that is  $G(x_n, x_m, x_p) \rightarrow 0$  as  $n, m, p \rightarrow \infty$ .

A  $G$  - metric space  $(X, G)$  is said to be  $G$  - complete if every  $G$  - Cauchy sequence is  $G$  - convergent.

Lemma 2.1 [12] Let  $(X, G)$  be a  $G$  - metric space. Then, the following properties are equivalent:

- 1)  $(x_n)$  is  $G$  - convergent to  $x$ ;
- 2)  $G(x_n, x_n, x) \rightarrow 0$  as  $n \rightarrow \infty$ ;
- 3)  $G(x_n, x, x) \rightarrow 0$  as  $n \rightarrow \infty$ ;
- 4)  $G(x_n, x_m, x) \rightarrow 0$  as  $n, m \rightarrow \infty$ .

Lemma 2.2 [12] If  $(X, G)$  is a  $G$  - metric space, the following properties are equivalent:

- 1)  $(x_n)$  is  $G$  - Cauchy;
- 2) For  $\varepsilon > 0$ , there exists  $k \in \mathbb{N}$  such that  $G(x_n, x_m, x_m) < \varepsilon$  for all  $m, n \in \mathbb{N}, m, n \geq k$ .

Lemma 2.3 [12] Let  $(X, G)$  be a  $G$  - metric space. Then, the function  $G(x, y, z)$  is jointly continuous in all three of its variables.

Note that each  $G$  - metric generates a topology  $\tau_G$  on  $X$  [12], whose base is a family of open  $G$  - balls  $\{B_G(x, \varepsilon) : x \in X, \varepsilon > 0\}$ , where  $B_G(x, \varepsilon) = \{y \in X : G(x, y, y) < \varepsilon\}$  for all  $x, y \in X$  and  $\varepsilon > 0$ .

A nonempty set  $A \subset X$  is  $G$  - closed if  $A = \overline{A}$ .

Lemma 2.4 [10] Let  $(X, G)$  be a  $G$  - metric space and  $A$  a subset of  $X$ .  $A$  is  $G$  - closed if for any  $G$  - convergent sequence in  $A$  with  $\lim_{n \rightarrow \infty} x_n = x$ , then  $x \in A$ .

In [2], [22], [23] and other papers some fixed point theorems for weakly compatible mappings in  $G$  - metric spaces are proved. Quite recently, in [9] the following theorem is proved.

Theorem 2.1 (Theorem 2.1 [9]) Let  $(X, G)$  be a complete  $G$  - metric space. Suppose that  $(f, S)$  and  $(g, T)$  are two pairs weakly compatible self - mappings on  $X$  satisfying

$$G(fx, fx, gy) \leq h \max\{G(Sx, Sx, Ty), G(fx, fx, Sx), G(gy, gy, Ty), \frac{1}{2}[G(fx, fx, Ty) + G(gy, gy, Sx)]\} \quad (2.1)$$

and

$$G(fx, gy, gy) \leq h \max\{G(Sx, Ty, Ty), G(fx, Sx, Sx), G(gy, Ty, Ty), \frac{1}{2}[G(fx, Ty, Ty) + G(gy, Sx, Sx)]\} \quad (2.2)$$

for all  $x, y \in X$ , where  $h \in \left[0, \frac{1}{2}\right)$ . Suppose that  $f(X) \subset T(X)$  and  $g(X) \subset S(X)$ . If one of  $T(X)$  or  $S(X)$  is a  $G$ -closed subspace of  $X$ , then  $f, g, S$  and  $T$  have a unique common fixed point.

The purpose of this paper is to prove a general fixed point theorem for two pairs of weakly compatible self-mappings of  $(X, G)$  satisfying  $\phi$ -implicit relations in  $G$ -metric spaces which generalize and improve Theorem 2.1.

### 3. IMPLICIT RELATIONS

**Definition 3.1** [23] A function  $f: [0, \infty) \rightarrow [0, \infty)$  is a  $\phi$ -function,  $f \in \phi$ , if  $f$  is nondecreasing function such that  $\sum_{n=1}^{\infty} f^n(t) < +\infty$ , for all  $t > 0$  and  $f(0) = 0$ .

**Definition 3.2** [23] Let  $F_{\phi}$  be the set of all continuous functions  $F(t_1, \dots, t_6): P_+^6 \rightarrow P$  such that

$(F_1)$ :  $F$  is nonincreasing in variable  $t_5$ ,

$(F_2)$ : there exists a function  $\phi_1 \in \phi$  such that for all  $u, v \geq 0$  with  $F(u, v, v, u, u+v, 0) \leq 0$  implies  $u \leq \phi_1(v)$ ,

$(F_3)$ : there exists a function  $\phi_2 \in \phi$  such that for all  $t, t' > 0$ ,  $F(t, t, 0, 0, t, t') \leq 0$  implies  $t \leq \phi_2(t')$ .

The following examples are presented in [23].

**Example 3.1.**

$$F(t_1, \dots, t_6) = t_1 - k \max\left\{t_2, t_3, t_4, \frac{t_5 + t_6}{2}\right\},$$

where  $k \in (0, 1)$ .

**Example 3.2.**

$$F(t_1, \dots, t_6) = t_1 - at_2 - bt_3 - ct_4 - dt_5 - et_6,$$

where  $a > 0, b, c, d, e \geq 0$  and  $a + b + c + 2d + e < 1$ .

**Example 3.3.**

$$F(t_1, \dots, t_6) = t_1 - k \max\{t_2, t_3, t_4, t_5, t_6\},$$

where  $k \in \left[0, \frac{1}{2}\right)$ .

**Example 3.4.**

$$F(t_1, \dots, t_6) = t_1^2 - t_1(at_2 + bt_3 + ct_4) - dt_5t_6,$$

where  $a > 0, b, c, d \geq 0$  and  $a + b + c + d < 1$ .

**Example 3.5.**

$$F(t_1, \dots, t_6) = t_1 - k \max\left\{t_2, \frac{t_3 + t_4}{2}, \frac{t_5 + t_6}{2}\right\},$$

where  $k \in (0, 1)$ .

**Example 3.6.**

$$F(t_1, \dots, t_6) = t_1^3 - c \frac{t_3^2 t_4^2 + t_5^2 t_6^2}{1 + t_2 + t_3 + t_4},$$

where  $c \in [0, 1)$ .

**Example 3.7.**

$$F(t_1, \dots, t_6) = t_1^2 - at_2^2 - c \frac{t_5 t_6}{1 + t_3 + t_4},$$

where  $a \geq 0$  and  $a + c < 1$ .

**Example 3.8.**

$$F(t_1, \dots, t_6) = t_1 - at_2 - bt_3 - c \max\{2t_4, t_5 + t_6\},$$

where  $a > 0, b, c \geq 0$  and  $a + b + 2c < 1$ .

**Example 3.9.**

$$F(t_1, \dots, t_6) = t_1 - at_2 - bt_3 - c \max\{t_4 + t_5, 2t_6\},$$

where  $a > 0, b, c \geq 0$  and  $a + b + 3c < 1$ .

**Example 3.10.**

$F(t_1, \dots, t_6) = t_1 - c \max\{t_2, t_3, \sqrt{t_4 t_6}, \sqrt{t_5 t_6}\}$ ,  
where  $c \in (0, 1)$ .

#### 4. MAIN RESULTS

Theorem 4.1 [1] Let  $f, g$  be weakly compatible self mappings on a nonempty set  $X$ . If  $f$  and  $g$  have an unique point of coincidence  $w = fx = gx$ , then  $w$  is the unique common fixed point of  $f$  and  $g$ .

Theorem 4.2 Let  $f, g, S$  and  $T$  be self mappings of a  $G$ -metric space such that

$$H_1(G(Sx, Ty, Ty), G(fx, gy, gy), G(fx, Sx, Sx), G(gy, Ty, Ty), G(fx, Ty, Ty), G(gy, Sx, Sx)) \leq 0, \quad (4.1)$$

$$H_2(G(Tx, Sy, Sy), G(gx, fy, fy), G(gx, Tx, Tx), G(fy, Sy, Sy), G(gx, Sy, Sy), G(fy, Tx, Tx)) \leq 0, \quad (4.2)$$

for all  $x, y \in X$ , where  $H_1, H_2$  satisfy property  $(F_3)$ .

If there exist  $u, v \in X$  such that  $fu = Su$  and  $gv = Tv$ , then there exists  $t \in X$  such that  $t$  is the unique point of coincidence of  $f$  and  $S$ , as well  $t$  is the unique point of coincidence of  $g$  and  $T$ .

Proof. First we prove that  $fu = gv$ . Suppose that  $fu \neq gv$ . Then, by (4.1) we have successively

$$H_1(G(Su, Tv, Tv), G(fu, gv, gv), G(fu, Su, Su), G(gv, Tv, Tv), G(fu, Tv, Tv), G(gv, Su, Su)) \leq 0,$$

$$H_1(G(fu, gv, gv), G(fu, gv, gv), 0, 0, G(fu, gv, gv), G(gv, fu, fu)) \leq 0,$$

By  $(F_3)$  we obtain

$$G(fu, gv, gv) \leq \phi_2(G(gv, fu, fu)) < G(gv, fu, fu).$$

Similarly, by (4.2) we have successively

$$H_2(G(Tv, Su, Su), G(gv, fu, fu), G(gv, Tv, Tv), G(fv, Su, Su), G(gv, Su, Su), G(fu, Tv, Tv)) \leq 0,$$

$$H_2(G(gv, fu, fu), G(gv, fu, fu), 0, 0, G(gv, fu, fu), G(fu, gv, gv)) \leq 0,$$

By  $(F_3)$  we obtain

$$G(gv, fu, fu) \leq \phi_2(G(fu, gv, gv)) < G(fu, gv, gv).$$

Hence,

$$G(fu, gv, gv) \leq G(gv, fu, fu) < G(fu, gv, gv),$$

a contradiction. Hence,  $fu = gv$  which implies  $t = fu = gv = Su = Tv$  for some  $t \in X$ . Therefore  $t$  is a common point of coincidence of  $(f, S)$  and  $(g, T)$ . We prove that  $t$  is the unique point of coincidence of  $(f, S)$ . Suppose that there is other point of coincidence  $z = fw = Sw$ ,  $z \neq t$ .

By (4.1) we have successively

$$H_1(G(Sw, Tv, Tv), G(fw, gv, gv), G(fw, Sw, Sw), G(gv, Tv, Tv), G(fw, Tv, Tv), G(gv, Sw, Sw)) \leq 0,$$

$$H_1(G(fw, gv, gv), G(fw, gv, gv), 0, 0, G(fw, gv, gv), G(gv, fw, fw)) \leq 0.$$

By  $(F_3)$  we have

$$G(fv, gv, gv) \leq \phi_2(G(gv, fw, fw)) < G(gv, fw, fw).$$

Similarly, by (4.2) and  $(F_3)$  we obtain

$$G(gv, fw, fw) < G(fw, gv, gv).$$

Hence,

$$G(fw, gv, gv) < G(gv, fw, fw) < G(fw, gv, gv),$$

a contradiction. Hence,  $fw = gv = fu = t$ . Therefore,

$$fw = Sw = t = z.$$

Similarly, we prove that  $t$  is the unique point of coincidence of  $g$  and  $T$ .

Theorem 4.3. Let  $(X, G)$  be a  $G$ -metric space and  $\{f, S\}$  and  $\{g, T\}$  be two pairs of self mappings of  $X$  satisfying the inequalities (4.1), (4.2) for all  $x, y \in X$ ,  $H_1, H_2 \in \mathbf{F}_\phi$  and

$$S(X) \subset g(X) \text{ and } T(X) \subset f(X).$$

(4.3)

If one of  $f(X)$  or  $g(X)$  is a  $G$ -closed subset of  $X$ , then

a)  $(f, S)$  have a coincidence point,

b)  $(g, T)$  have a coincidence point.

Moreover, if  $(f, S)$  and  $(g, T)$  are weakly compatible, then  $f, g, S$  and  $T$  have an unique common fixed point.

Proof. Let  $x_0 \in X$  be an arbitrary point of  $X$ . Since  $S(X) \subset g(X)$  and  $T(X) \subset f(X)$ , there exist  $x_1, x_2 \in X$  such that  $Sx_0 = gx_1$  and  $Tx_1 = fx_2$ .

By continuing this process, for  $n = 0, 1, 2, \dots$  we choose  $x_n, y_n \in X$  such that

$$y_{2n} = Sx_{2n} = gx_{2n+1}, \quad y_{2n+1} = Tx_{2n+1} = fx_{2n+2}.$$

By (4.1) we have successively

$$H_1(G(Sx_{2n}, Tx_{2n+1}, Tx_{2n+1}), G(fx_{2n}, gx_{2n+1}, gx_{2n+1}), G(fx_{2n}, Sx_{2n}, Sx_{2n}), \\ G(gx_{2n+1}, Tx_{2n+1}, Tx_{2n+1}), G(fx_{2n}, Tx_{2n+1}, Tx_{2n+1}), G(gx_{2n+1}, Sx_{2n}, Sx_{2n})) \leq 0,$$

$$H_1(G(y_{2n}, y_{2n+1}, y_{2n+1}), G(yx_{2n-1}, y_{2n}, y_{2n}), G(y_{2n-1}, y_{2n}, y_{2n}), \\ G(y_{2n}, y_{2n+1}, y_{2n+1}), G(y_{2n-1}, y_{2n+1}, y_{2n+1}), 0) \leq 0.$$

By  $(F_1)$  and  $(G_5)$  we obtain

$$H_1(G(y_{2n}, y_{2n+1}, y_{2n+1}), G(y_{2n-1}, y_{2n}, y_{2n}), G(y_{2n-1}, y_{2n}, y_{2n}), \\ G(y_{2n}, y_{2n+1}, y_{2n+1}), G(y_{2n-1}, y_{2n}, y_{2n}) + G(y_{2n}, y_{2n+1}, y_{2n+1}), 0) \leq 0.$$

By  $(F_2)$  we obtain

$$G(y_{2n}, y_{2n+1}, y_{2n+1}) \leq \phi_1(G(y_{2n-1}, y_{2n}, y_{2n})).$$

Again, by (4.2) we have successively

$$H_2(G(Tx_{2n+1}, Sx_{2n+2}, Sx_{2n+2}), G(gx_{2n+1}, fx_{2n+2}, fx_{2n+2}), G(gx_{2n+1}, Tx_{2n+1}, Tx_{2n+1}), \\ G(fx_{2n+1}, Sx_{2n+2}, Sx_{2n+2}), G(gx_{2n+1}, Sx_{2n+2}, Sx_{2n+2}), G(fx_{2n+2}, Tx_{2n+1}, Tx_{2n+1})) \leq 0,$$

$$H_2(G(y_{2n+1}, y_{2n+2}, y_{2n+2}), G(yx_{2n}, y_{2n+1}, y_{2n+1}), G(y_{2n}, y_{2n+1}, y_{2n+1}), \\ G(y_{2n+1}, y_{2n+2}, y_{2n+2}), G(y_{2n}, y_{2n+2}, y_{2n+2}), 0) \leq 0.$$

By  $(F_1)$  and  $(G_5)$  we have that

$$H_2(G(y_{2n+1}, y_{2n+2}, y_{2n+2}), G(y_{2n}, y_{2n+1}, y_{2n+1}), G(y_{2n}, y_{2n+1}, y_{2n+1}), \\ G(y_{2n+1}, y_{2n+2}, y_{2n+2}), G(y_{2n}, y_{2n+1}, y_{2n+1}) + G(y_{2n}, y_{2n+2}, y_{2n+2}), 0) \leq 0.$$

By  $(F_2)$  we have

$$G(y_{2n+1}, y_{2n+2}, y_{2n+2}) \leq \phi_1(G(y_{2n}, y_{2n+1}, y_{2n+1})),$$

which implies

$$G(y_n, y_{n+1}, y_{n+1}) \leq \phi_1(G(y_{n-1}, y_n, y_n)), n = 1, 2, \dots$$

Then

$$G(y_n, y_{n+1}, y_{n+1}) \leq \phi_1^n(G(y_0, y_1, y_1)).$$

We prove that  $(y_n)$  is a  $G$  - Cauchy sequence in  $X$ .

For  $n, m \in \mathbb{N}$  with  $m > n$ , we have repeating  $(G_5)$  that

$$\begin{aligned} G(y_n, y_m, y_m) &\leq G(y_n, y_{n+1}, y_{n+1}) + G(y_{n+1}, y_{n+2}, y_{n+2}) + \dots + G(y_{m-1}, y_m, y_m) \\ &\leq \sum_{k=n}^{m-1} \phi_1^k(G(y_0, y_1, y_1)). \end{aligned}$$

Since  $\sum_{k=1}^{\infty} \phi_1^k(G(y_0, y_1, y_1)) < +\infty$ , then for any  $\varepsilon > 0$ , there exists  $k \in \mathbb{N}$  such that for  $m, n \geq k$ ,

$\sum_{k=n}^{m-1} \phi_1^k(G(y_0, y_1, y_1)) < \varepsilon$ . Hence, by Lemma 2.2,  $y_n$  is a  $G$  - Cauchy sequence. Since  $(X, G)$  is  $G$  - complete,

there exists  $z \in X$  such that  $y_n \rightarrow z$  as  $n \rightarrow \infty$ . This implies that  $\lim_{n \rightarrow \infty} y_{2n} = \lim_{n \rightarrow \infty} y_{2n+1} = z$ .

Suppose that  $g(X)$  is closed. It follows that  $z = gu$ , for some  $u \in X$ . Using (4.1) we have successively

$$\begin{aligned} &H_1(G(Sx_{2n}, Tu, Tu), G(fx_{2n}, gu, gu), G(fx_{2n}, Sx_{2n}, Sx_{2n}), \\ &G(gu, Tu, Tu), G(fx_{2n}, Tu, Tu), G(gu, Sx_{2n}, Sx_{2n})) \leq 0, \\ &H_1(G(y_{2n}, Tu, Tu), G(y_{2n-1}, gu, gu), G(y_{2n-1}, y_{2n}, y_{2n}), \\ &G(gu, Tu, Tu), G(y_{2n-1}, Tu, Tu), G(gu, y_{2n}, y_{2n})) \leq 0. \end{aligned}$$

Letting  $n$  tend to infinity we obtain

$$H_1(G(z, Tu, Tu), 0, 0, G(z, Tu, Tu), G(z, Tu, Tu), 0) \leq 0,$$

which implies by  $(F_2)$  that  $G(z, Tu, Tu) \leq \phi(0) = 0$ . Hence,  $z = Tu = gu$  and  $u$  is a coincidence point of  $g$  and  $T$ .

Since  $T(X) \subset f(X)$ , there exists  $v \in X$  such that  $z = Tu = fv = gu$ . Then by (4.2) we have successively

$$\begin{aligned} &H_2(G(Tu, Sv, Sv), G(gu, fv, fv), G(gu, Tu, Tu), \\ &G(fv, Sv, Sv), G(gu, Sv, Sv), G(fv, Tu, Tu)) \leq 0, \\ &H_2(G(gu, Sv, Sv), 0, 0, G(gu, Sv, Sv), G(gu, Sv, Sv), 0) \leq 0 \end{aligned}$$

which implies by  $(F_2)$  that

$$G(gu, Sv, Sv) \leq \phi(0) = 0,$$

i.e.  $z = gu = Sv = fv$ . Hence,  $v$  is a coincidence point of  $S$  and  $f$ .

Therefore,  $z$  is the common point of coincidence of  $(f, S)$  and  $(g, T)$ . By Theorem 4.1,  $z$  is the unique point of coincidence of  $(f, S)$  and  $(g, T)$ . By Theorem 4.1,  $z$  is the unique common fixed point of  $f, g, S$  and  $T$ .

Corollary 4.1 Let  $(X, G)$  be a complete  $G$  - metric space. Suppose that  $(f, S)$  and  $(g, T)$  are weakly compatible pairs of self mappings of  $X$  satisfying

$$\begin{aligned} G(Sx, Ty, Ty) &\leq h \max\{G(fx, gy, gy), G(fx, Sx, Sx), \\ G(gy, Ty, Ty), &\frac{1}{2}[G(fx, Ty, Ty) + G(gy, Sx, Sx)]\}, \end{aligned} \quad (4.4)$$

$$\begin{aligned} G(Tx, Sy, Sy) &\leq h \max\{G(gx, fy, fy), G(gx, Tx, Tx), \\ G(fy, Sy, Sy), &\frac{1}{2}[G(gx, Sy, Sy) + G(fy, Tx, Tx)]\} \leq 0, \end{aligned} \quad (4.5)$$

for all  $x, y \in X$  and  $h \in (0, 1)$ . Suppose that  $S(X) \subset g(X)$  and  $T(X) \subset f(X)$ . If one of  $g(X)$  or  $f(X)$  is a  $G$  - closed subspace of  $X$ , then  $f, g, S$  and  $T$  have a unique common fixed point.

Proof. The proof it follows from Theorem 4.3 and Example 3.1 with  $\phi_1(t) = \phi_2(t) = kt$ .

Remark 4.1

1. In the proof of Theorem 2.1 [3], page 4, lines 10 and 11 from the bottom, there exists some written mistakes and hence the proof of the fact that the sequence  $(y_n)$  is a  $G$  - Cauchy sequence is not correct. Similarly, in the proof of Theorem 2.1 [9]. For a correct form of Theorem 2.1, we suggest the inequality

$$\begin{aligned} G(fx, gy, gy) &\leq h \max\{G(Sx, Ty, Ty), G(Sx, fx, fx), G(Ty, gy, gy), \\ &\frac{1}{2}[G(Sx, gy, gy) + G(Ty, fx, fx)]\} \end{aligned}$$

instead inequality (2.2).

2. Corollary 4.1 is a generalization and the correct form of Theorem 2.1 because  $h \in (0, 1)$  instead  $h \in \left[0, \frac{1}{2}\right)$  and the

fact that  $(y_n)$  is a  $G$  - Cauchy sequence is correct.

3. By Examples 3.2 – 3.10 we obtain new particular results.

## REFERENCES

- [1] Abbas, M. and Rhoades, B. E., "Common fixed point theorems under strict contractive condition", Math. Anal. Appl., 270: 181-188 (2002).
- [2] Abbas, M. and Rhoades, B. E., "Common fixed point results for noncommuting mappings without continuity in generalized metric spaces", Appl. Math. Comput., 215: 262-269 (2009).
- [3] Abbas, M., Khan, S. H. and Nazir, T., "Common fixed points for  $R$  - weakly commuting maps in generalized metric spaces", Fixed Point Theory Appl., 2011, 2011:41.
- [4] Altun, I. and Turkoglu, D., "Some fixed point theorems for weakly compatible mappings satisfying an implicit relation", Taiwanese J. Math., 13, 4: 1291-1304 (2009).
- [5] Dhage, B. C., "Generalized metric spaces and mappings with fixed point", Bull. Calcutta Math. Soc., 84: 329-336 (1992).
- [6] Dhage, B. C., "Generalized metric spaces and topological structures I", An. Ştiinţ. Univ. Al. I. Cuza Iaşi, Mat., 46 (1): 3-24 (2000).
- [7] Imdad, M., Kumar, S. and Khan, M. S., "Remarks on some fixed point theorems satisfying implicit relations", Rad. Mat., 11: 135-143 (2002).
- [8] Jungck, G., "Common fixed points for noncontinuous nonself maps on nonnumeric spaces", Far East J. Math. Sci., 4, 2: 195-215 (1996).
- [9] Kaewcharoen, A., "Common fixed points for four mappings in  $G$  - metric spaces", Int. J. Math. Anal., 6 (47): 2345-2356 (2012).
- [10] Karpinar, E., Yildiz – Ulus, A. and Erhan, I. M., "Cyclic contractions on  $G$  - metric spaces", Abstr. Appl. Anal., Volume 2013, Article ID 182747.
- [11] Mustafa, Z. and Sims, B., "Some remarks concerning  $D$  - metric spaces", Proc. Conf. Fixed Point Theory and Applications, Valencia (Spain), 189-198 (2003).
- [12] Mustafa, Z. and Sims, B., "A new approach to generalized metric spaces", J. Nonlinear Convex Anal., 7: 287-297 (2006).

- [13] Mustafa, Z., Obiedat, H. and Awawdeh, F., "Some fixed point theorems for mappings on  $G$  - complete metric spaces", Fixed Point Theory Appl., Volume 2008, Art. ID 189879, 12 pages.
- [14] Mustafa, Z. and Sims, B., "Fixed point theorems for contractive mappings in complete  $G$  - metric spaces", Fixed Point Theory Appl., Volume 2009, Art. ID 917175, 10 pages.
- [15] Mustafa, Z. and Obiedat, H., "A fixed point theorem of Reich in  $G$  - metric spaces", Cubo, 12: 73-93 (2010).
- [16] Mustafa, Z., Kandagji, M. and Shatanawi, W., "Fixed point results on complete  $G$  - metric spaces", Stud. Sci. Math. Hung., 48 (3): 304-315 (2011).
- [17] Pant, R. P., "Common fixed points for noncommuting mappings", J. Math. Anal. Appl., 188: 436-440 (1994).
- [18] Pant, R. P., "Common fixed points for four mappings", Bull. Calcutta Math. Soc., 9: 281-286 (1998).
- [19] Popa, V., "Some fixed point theorems for implicit contractive mappings", Stud. Cercet. Ştiinţ., Ser. Mat., Univ. Bacău, 7: 129-133 (1997).
- [20] Popa, V., "Some fixed point theorems for compatible mappings satisfying an implicit relation", Demonstr. Math., 32: 157-163 (1999).
- [21] Popa, V., "A general fixed point theorem for several mappings in  $G$  - metric spaces", Sci. Stud. Res., Ser. Math. Inform., 21 (1): 205-214 (2011).
- [22] Popa, V. and Patriciu, A.-M., "A general fixed point theorem for pairs of weakly compatible mappings in  $G$  - metric spaces", J. Nonlinear Sci. Appl., 5 (2): 151-160 (2012).
- [23] Popa, V. and Patriciu, A.-M., "A general fixed point theorem for mappings satisfying an  $\phi$  - implicit relation in complete  $G$  - metric spaces", GU J. Sci., 25 (2): 403-408 (2012).
- [24] Popa, V. and Patriciu, A.-M., "Two general fixed point theorems of pairs of weakly compatible mappings in  $G$  - metric spaces", Novi Sad J. Math., 42 (2): 49-60 (2012).
- [25] Popa, V. and Patriciu, A.-M., "Two common fixed point theorems for three mappings satisfying a  $\phi$  - implicit relation in complete  $G$  - metric spaces", Vietnam J. Math., 41 (2): 233-249 (2013).
- [26] Popa, V. and Patriciu, A.-M., "Fixed point results on complete  $G$  - metric spaces satisfying an implicit relation of new type", Ukr. Math. J., 65 (6): 816-821 (2013).
- [27] Raswan, R. A. and Saleh, S. M., "Common fixed point theorems for mappings satisfying  $\phi$  - implicit relation in complete  $G$  - metric spaces", Proc. Pakistan Acad. Sci., 5 (3): 227-234 (2012).
- [28] Shatanawi, W., "Fixed point theory for contractive mappings satisfying  $\phi$  - maps in  $G$  - metric spaces", Fixed Point Theory Appl., 2010, Article ID 189650, 9 pages.