

PAPER DETAILS

TITLE: q -Harmonic mappings for which analytic part is q -convex functions of complex order

AUTHORS: Asena ÇETINKAYA, Yasar POLATO GLU

PAGES: 813-820

ORIGINAL PDF URL: <https://dergipark.org.tr/tr/download/article-file/522556>

q – Harmonic mappings for which analytic part is q – convex functions of complex order

Asena Çetinkaya^{*†} and Yaşar Polatoğlu[‡]

Abstract

We introduce a new class of harmonic function f , that is subclass of planar harmonic mapping associated with q – difference operator. Let h and g are analytic functions in the open unit disc $\mathbb{D} = \{z : |z| < 1\}$. If $f = h + \bar{g}$ is the solution of the non-linear partial differential equation $w_q(z) = \frac{D_q g(z)}{D_q h(z)} = \frac{\bar{f}_z}{f_z}$ with $|w_q(z)| < 1$, $w_q(z) \prec b_1 \frac{1+z}{1-qz}$ and h is q – convex function of complex order, then the class of such functions are called q – harmonic functions for which analytic part is q – convex functions of complex order denoted by $\mathcal{S}_{\mathcal{H}\mathcal{C}_q(b)}$. Obviously that the class $\mathcal{S}_{\mathcal{H}\mathcal{C}_q(b)}$ is the subclass of $\mathcal{S}_{\mathcal{H}}$. In this paper, we investigate properties of the class $\mathcal{S}_{\mathcal{H}\mathcal{C}_q(b)}$ by using subordination techniques.

Keywords: q – difference operator, q – harmonic mapping, q – convex function of complex order.

Mathematics Subject Classification (2010): 30C45

Received : 21.03.2017 *Accepted :* 31.05.2017 *Doi :* 10.15672/HJMS.2017.480

^{*}Department of Mathematics and Computer Sciences, İstanbul Kültür University, İstanbul, TURKEY ,

Email : asnfigen@hotmail.com

[†]Corresponding Author.

[‡]Department of Mathematics and Computer Sciences, İstanbul Kültür University, İstanbul, TURKEY,

Email : y.polatoglu@iku.edu.tr

1. Introduction

A planar harmonic mapping in the open unit disc \mathbb{D} is a complex valued harmonic function f , which maps \mathbb{D} onto the some planar domain $f(\mathbb{D})$. Since \mathbb{D} is a simply connected domain, the mapping f has a canonical decomposition $f = h + \bar{g}$, where h and g are analytic in \mathbb{D} and have the following power series expansions

$$h(z) = z + \sum_{n=2}^{\infty} a_n z^n \quad \text{and} \quad g(z) = \sum_{n=1}^{\infty} b_n z^n$$

where $a_n, b_n \in \mathbb{C}$, $n = 0, 1, 2, 3, \dots$. As usual, we call h the analytic part of f and g the co-analytic part of f , respectively. An elegant and complete treatment theory of the harmonic mapping is given in Duren's monograph [3]. Lewy [11] proved in 1936 that the harmonic mapping f is locally univalent in \mathbb{D} if and only if its Jacobian $J_f = |h'(z)|^2 - |g'(z)|^2$ is different from zero in \mathbb{D} . In view of this result, locally univalent harmonic mappings in the open unit disc are either sense-preserving if $|g'(z)| < |h'(z)|$ or sense-reversing if $|g'(z)| > |h'(z)|$ in \mathbb{D} . Throughout this paper, we will restrict ourselves to the study of sense-preserving harmonic mappings. We also note that $f = h + \bar{g}$ is sense-preserving in \mathbb{D} if and only if h' does not vanish in \mathbb{D} and the second dilatation $w(z) = \frac{g'(z)}{h'(z)}$ has the property $|w(z)| < 1$ for all $z \in \mathbb{D}$. Therefore the class of all sense-preserving harmonic mappings in \mathbb{D} with $a_0 = b_0 = 0$ and $a_1 = 1$ will be denoted by $\mathcal{S}_{\mathcal{H}}$. Thus $\mathcal{S}_{\mathcal{H}}$ contains standard class \mathcal{S} of analytic univalent functions. The family of all mappings $f \in \mathcal{S}_{\mathcal{H}}$ with the additional property that $g'(0) = 0$, i.e., $b_1 = 0$ are denoted by $\mathcal{S}_{\mathcal{H}}^0$. Hence it is clear that $\mathcal{S} \subset \mathcal{S}_{\mathcal{H}}^0 \subset \mathcal{S}_{\mathcal{H}}$.

In 1908 and 1910 Jackson [8, 9] initiated a study of q - difference operator by

$$(1.1) \quad D_q f(z) = \frac{f(z) - f(qz)}{(1-q)z} \quad \text{for } z \in B \setminus \{0\}$$

where B is a subset of complex plane \mathbb{C} , called q - geometric set if $qz \in B$, whenever $z \in B$. Note that if a subset B of \mathbb{C} is q - geometric, then it contains all geometric sequences $\{zq^n\}_0^\infty$, $zq \in B$. Obviously, $D_q f(z) \rightarrow f'(z)$ as $q \rightarrow 1^-$. The q - difference operator (1.1) is sometimes called Jackson q - difference operator. Note that such an operator plays an important role in the theory of hypergeometric series and quantum physics (see for instance [1, 4, 5, 10]).

Also, note that $D_q f(0) \rightarrow f'(0)$ as $q \rightarrow 1^-$ and $D_q^2 f(z) = D_q(D_q f(z))$. In fact, q - calculus is ordinary classical calculus without the notion of limits. Recent interest in q - calculus is because of its applications in various branches of mathematics and physics. For definition and properties of q - difference operator and q - calculus, one may refer to [1, 4, 5, 10].

Under the hypothesis of the definition of q - difference operator, then we have the following rules:

- (1) For a function $f(z) = z^n$, we observe that

$$D_q z^n = \frac{1 - q^n}{1 - q} z^{n-1}.$$

Therefore we have

$$D_q f(z) = 1 + \sum_{n=2}^{\infty} a_n \frac{1 - q^n}{1 - q} z^{n-1}.$$

- (2) If functions f and g are defined on a q - geometric set $B \subset \mathbb{C}$ such that q - derivatives of f and g exist for all $z \in B$, then

- (i) $D_q(af(z) \pm bg(z)) = aD_q f(z) \pm bD_q g(z)$ where a and b are real or complex constants.

- (ii) $D_q(f(z).g(z)) = g(z)D_qf(z) + f(qz)D_qg(z).$
- (iii) $D_q\left(\frac{f(z)}{g(z)}\right) = \frac{g(qz)D_qf(z) - f(qz)D_qg(z)}{g(z)g(qz)}, \quad g(z)g(qz) \neq 0.$
- (iv) As a right inverse, Jackson introduced q - integral

$$\int_0^z f(t)d_qt = z(1-q) \sum_{n=0}^{\infty} q^n f(zq^n)$$

provided that the series converges.

The following theorem is an analogue of the fundamental theorem of calculus.

A. Theorem. ([10]) Let f be a q - regular at zero, defined on q - geometric set B containing zero. Define

$$F(z) = \int_c^z f(\zeta)d_q\zeta, \quad (\zeta \in B)$$

where c is a fixed point in B , then F is q - regular at zero. Furthermore $D_qF(z)$ exists for every $z \in B$ and

$$D_qF(z) = f(z)$$

for every $z \in B$.

Conversely, if a and b are two points in B , then

$$\int_a^b D_qf(\zeta)d_q\zeta = f(b) - f(a).$$

- (3) The q - differential is defined as

$$d_qf(z) = f(z) - f(qz).$$

Therefore we can write

$$d_qf(z) = \frac{f(z) - f(qz)}{(1-q)z} d_qz.$$

- (4) The partial q - derivative of a multivariable real continuous functions $f(x_1, x_2, \dots, x_{i-1}, x_i, x_{i+1}, \dots, x_n)$ to a variable x_i is defined by

$$D_{q,x_i}f(\vec{x}) = \frac{f(\vec{x}) - \varepsilon_{q,x_i}f(\vec{x})}{(1-q)x_i}, \quad x_i \neq 0, q \in (0,1)$$

$$\left[D_{q,x_i}f(\vec{x}) \right]_{x_i=0} = \lim_{x_i \rightarrow 0} D_{q,x_i}f(\vec{x})$$

where $\varepsilon_{q,x_i}f(\vec{x}) = f(x_1, x_2, \dots, x_{i-1}, qx_i, x_{i+1}, \dots, x_n)$ and we use $D_{k,x}^k$ instead of operator $\frac{\partial^k}{\partial_q x^k}$ for some simplification.

Finally, let Ω be the family of functions ϕ analytic in \mathbb{D} , and satisfy the conditions $\phi(0) = 0$, $|\phi(z)| < 1$ for all $z \in \mathbb{D}$. Denote by \mathcal{P}_q the family of functions p of the form $p(z) = 1 + p_1z + p_2z^2 + \dots$, analytic in \mathbb{D} and satisfy the condition

$$(1.2) \quad \left| p(z) - \frac{1}{1-q} \right| \leq \frac{1}{1-q}, \quad z \in \mathbb{D}$$

where $q \in (0,1)$ is a fixed real number. Let \mathcal{A} be the family of functions f , defined by $f(z) = z + a_2z^2 + a_3z^3 + \dots$, that are analytic in \mathbb{D} and satisfy the conditions $f(0) = 0$, $f'(0) = 1$. If f satisfies the condition

$$1 + \frac{1}{b} \left(qz \frac{D_q(D_qf(z))}{D_qf(z)} \right) \prec \frac{1+z}{1-qz},$$

where $b \in \mathbb{C}$, $b \neq 0$, then f is called q - convex function of complex order, and the class of such functions are denoted by $\mathcal{C}_q(b)$. If f_1 and f_2 are analytic functions in \mathbb{D} , then we say

that f_1 is subordinate to f_2 , written as $f_1 \prec f_2$ if there exists a Schwarz function $\phi \in \Omega$ such that $f_1(z) = f_2(\phi(z))$, $z \in \mathbb{D}$. We also note that if f_2 univalent in \mathbb{D} , then $f_1 \prec f_2$ if and only if $f_1(0) = f_2(0)$ and $f_1(\mathbb{D}) \subset f_2(\mathbb{D})$. This implies that $f_1(\mathbb{D}_r) \subset f_2(\mathbb{D}_r)$, where $\mathbb{D}_r = \{z : |z| < r, 0 < r < 1\}$ (Subordination principle [6]).

We also need the following lemmas:

1.1. Lemma. Let ϕ be analytic in \mathbb{D} with $\phi(0) = 0$ and $|\phi(z)| < 1$, $z \in \mathbb{D}$. If $|\phi(z)|$ attains its maximum value on the circle $|z| = r$ at a point z_0 , then we have

$$z_0 \phi'(z_0) = m \phi(z_0), \quad m \geq 1.$$

For more details of Jack's lemma, one may refer to [7].

1.2. Lemma. ([12]) If h is an element of $\mathcal{C}_q(b)$, then

$$F_2(|b|, Reb, q, r) \leq |D_q h(z)| \leq F_1(|b|, Reb, q, r)$$

where

$$F_1(|b|, Reb, q, r) = \left[(1 - qr)^{Reb+|b|} \cdot (1 + qr)^{Reb-|b|} \right]^{-\frac{1-q^2}{2q^2 \log q^{-1}}},$$

$$F_2(|b|, Reb, q, r) = \left[(1 - qr)^{Reb-|b|} \cdot (1 + qr)^{Reb+|b|} \right]^{-\frac{1-q^2}{2q^2 \log q^{-1}}}.$$

The aim of this paper is to investigate properties of the class of q -harmonic functions for which analytic part is q -convex functions of complex order defined by

$$\mathcal{S}_{\mathcal{H}\mathcal{C}_q(b)} = \left\{ f = h + \bar{g} : w_q(z) = \frac{D_q g(z)}{D_q h(z)} = \frac{\bar{f}_{\bar{z}}}{f_z}, w_q(z) \prec b_1 \frac{1+z}{1-qz}, |w_q(z)| < 1, h \in \mathcal{C}_q(b) \right\},$$

where

$$D_q h(z) = \frac{h(z) - h(qz)}{(1-q)z} = f_z \quad \text{and} \quad D_q g(z) = \frac{g(z) - g(qz)}{(1-q)z} = \bar{f}_{\bar{z}}.$$

2. Main Results

In this section, we first assume that the function f is sense-preserving q -harmonic function if and only if $w_q(z) = \frac{\bar{f}_{\bar{z}}}{f_z}$ is analytic. To show that

(\Rightarrow) Let $f = h + \bar{g}$ be sense-preserving q -harmonic function, then we will show that w_q is analytic. Since $h(z) = z + \sum_{n=2}^{\infty} a_n z^n$ and $g(z) = \sum_{n=1}^{\infty} b_n z^n$ are analytic functions, then we can write q -derivatives of these functions as

$$D_q h(z) = 1 + \sum_{n=2}^{\infty} \frac{1-q^n}{1-q} a_n z^{n-1} \quad \text{and} \quad D_q g(z) = b_1 + \sum_{n=2}^{\infty} \frac{1-q^n}{1-q} b_n z^{n-1}.$$

We must note that when $q \rightarrow 1^-$, $D_q h(z)$ reduces to $h'(z)$ and $D_q g(z)$ reduces to $g'(z)$. The second q -dilatation and q -Jacobian are defined by

$$w_q(z) = \frac{D_q g(z)}{D_q h(z)} = \frac{\bar{f}_{\bar{z}}}{f_z},$$

$$J_{fq}(z) = |D_q h(z)|^2 - |D_q g(z)|^2.$$

Also, the total q -differential of $f(\vec{x})$ can be written in the following manner,

$$d_q f(\vec{x}) = D_{q,x_1} d_q x_1 + D_{q,x_2} d_q x_2 + D_{q,x_3} d_q x_3 + \cdots + D_{q,x_n} d_q x_n.$$

Therefore the q -differential can be written as

$$d_q f = D_{q,z} d_q z + D_{q,\bar{z}} d_q \bar{z}.$$

Consequently, f is locally univalent and sense-preserving if $|D_q h(z)| > |D_q g(z)|$ and sense-reversing if $|D_q g(z)| > |D_q h(z)|$. Note that $f_z \neq 0$ whenever $J_{fq}(z) > 0$. For sense-preserving f , one sees that

$$(|D_q h(z)| - |D_q g(z)|)|d_q z| \leq |d_q f| \leq (|D_q h(z)| + |D_q g(z)|)|d_q z|.$$

With aid of these definitions, let $f = h + \bar{g}$ be the solution of the non-linear elliptic partial differential equation

$$w_q(z)f_z = \bar{f}_{\bar{z}}$$

under the condition $|w_q(z)| < 1$ for all $z \in \mathbb{D}$. A non-constant complex -valued function f is q -harmonic and orientation sense-preserving mapping on \mathbb{D} if and only if f is the solution of the non-linear elliptic partial differential equation

$$(2.1) \quad w_q(z)f_z = \bar{f}_{\bar{z}}$$

where

$$f_z = D_q h(z) = \frac{h(z) - h(qz)}{(1-q)z} \quad \text{and} \quad \bar{f}_{\bar{z}} = D_q g(z) = \frac{g(z) - g(qz)}{(1-q)z}.$$

If we take the q -derivative of equation (2.1) with respect to \bar{z} , we get

$$(2.2) \quad \bar{f}_{\bar{z}z} = f_{z\bar{z}}w_q(z) + f_z \frac{\partial w_q}{\partial \bar{z}}.$$

On the other hand, since f is q -harmonic, then we have $\triangle f = 4 \frac{\partial^2 f}{\partial z \partial \bar{z}} = 4f_{z\bar{z}} = 0$ and $\bar{f}_{\bar{z}z} = 0$. Therefore the equality (2.2) reduces to

$$(2.3) \quad f_z \frac{\partial w_q}{\partial \bar{z}} = 0$$

and this shows that $\frac{\partial w_q}{\partial \bar{z}} = 0$, that is, w_q is analytic.

(\Leftarrow) Conversely, if w_q is analytic in \mathbb{D} , then $\frac{\partial w_q}{\partial \bar{z}} = 0$. Therefore equality (2.2) reduces to

$$(2.4) \quad \bar{f}_{\bar{z}z} = f_{z\bar{z}}w_q(z).$$

On the other hand, using the definition of w_q , we have $|w_q(z)| < 1$. Thus, we get

$$(2.5) \quad 1 - |w_q(z)| \neq 0.$$

Considering (2.4) and (2.5), we obtain

$$(2.6) \quad \bar{f}_{\bar{z}z} = f_{z\bar{z}}w_q(z) \Rightarrow f_{z\bar{z}} = 0$$

and the equality (2.6) shows that f is q -harmonic. This proves our assumption.

We now investigate properties of the class $\mathcal{S}_{\mathcal{H}\mathcal{C}_q(b)}$. For Theorem 2.4, we need the following results. The first theorem is very important in order to obtain subordination of the analytic functions involving q -difference operator.

2.1. Theorem. ([2]) p is an element of \mathcal{P}_q if and only if $p(z) \prec \frac{1+z}{1-qz}$. This result is

sharp for the functions $p(z) = \frac{1+\phi(z)}{1-q\phi(z)}$, where ϕ is a Schwarz function.

Proof. If p is an element of \mathcal{P}_q , then we have

$$\left| p(z) - \frac{1}{1-q} \right| \leq \frac{1}{1-q} \Leftrightarrow |p(z) - m| \leq m,$$

where $m = \frac{1}{1-q} > 1$. Therefore we can write

$$\left| \frac{1}{m}p(z) - 1 \right| \leq 1.$$

Thus the function $\psi(z) = \frac{1}{m}p(z) - 1$ is analytic and has modulus at most 1 in \mathbb{D} , and so

$$\phi(z) = \frac{\psi(z) - \psi(0)}{1 - \overline{\psi(0)}\psi(z)} = \frac{(\frac{1}{m}p(z) - 1) - (\frac{1}{m} - 1)}{1 - (\frac{1}{m} - 1)(\frac{1}{m}p(z) - 1)}$$

satisfies the conditions of Schwarz lemma. This shows that we can write

$$p(z) = \frac{1 + \phi(z)}{1 - (1 - \frac{1}{m})\phi(z)} \Rightarrow p(z) \prec \frac{1 + z}{1 - qz}.$$

Conversely, suppose that the function p is analytic in \mathbb{D} and satisfies the condition $p(0) = 1$ and

$$\begin{aligned} p(z) \prec \frac{1 + z}{1 - qz} &\Rightarrow p(z) = \frac{1 + \phi(z)}{1 - (1 - \frac{1}{m})\phi(z)} \\ p(z) - m &= m \frac{\frac{1-m}{m} + \phi(z)}{1 + \frac{1-m}{m}\phi(z)}. \end{aligned}$$

On the other hand the function $\frac{\frac{1-m}{m} + \phi(z)}{1 + \frac{1-m}{m}\phi(z)}$ maps the unit circle onto itself, then we have

$$|p(z) - m| \leq m \Leftrightarrow \left| p(z) - \frac{1}{1 - q} \right| \leq \frac{1}{1 - q}.$$

This shows that $p \in \mathcal{P}_q$. □

We must note that the linear transformation $\frac{1+z}{1-qz}$ maps $|z| = r$ onto the disc with centre $C(r) = \frac{1+qr^2}{1-q^2r^2}$ and radius $\rho(r) = \frac{(1+q)r}{1-q^2r^2}$.

2.2. Lemma. If f is a function (real or complex valued) defined on q -geometric set \mathbb{B} with $|q| \neq 1$, then

$$D_q(\log f(z)) = \frac{D_q f(z)}{f(z)}.$$

Proof. Using the definition of q -difference operator, then we have

$$D_q(\log f(z)) = \frac{\log f(qz) - \log f(z)}{qz - z} = \log \left(1 + h \frac{D_q f(z)}{f(z)} \right)^{\frac{1}{h}}.$$

If we take limit for $h \rightarrow 0$, we obtain the desired result. □

2.3. Lemma. (q -Jack's Lemma) Let ϕ be analytic in \mathbb{D} with $\phi(0) = 0$. If $|\phi(z)|$ attains its maximum value on the circle $|z| = r$ at a point $z_0 \in \mathbb{D}$, then we have

$$z_0 D_q \phi(z_0) = m \phi(z_0)$$

where $m \geq 1$ is a real number.

Proof. Using the definition of q -difference operator and Jack's lemma, then we can write

$$D_q \phi(z) = \frac{\phi(z) - \phi(qz)}{z - qz} = \frac{\phi(z) - \phi(z_0)}{z - z_0}, \quad qz = z_0.$$

If we take limit for $z \rightarrow z_0$, we get

$$\lim_{z \rightarrow z_0} D_q \phi(z) = D_q \phi(z_0) = \lim_{z \rightarrow z_0} \frac{\phi(z) - \phi(z_0)}{z - z_0} = \phi'(z_0).$$

Therefore we have

$$z_0 D_q \phi(z_0) = m \phi(z_0).$$

□

2.4. Theorem. If $f = h + \bar{g}$ is an element of $\mathcal{S}_{\mathcal{H}\mathcal{C}_q(b)}$, then

$$(2.7) \quad \frac{g(z)}{h(z)} \prec b_1 \frac{1+z}{1-qz}.$$

Proof. Since $f = h + \bar{g} \in \mathcal{S}_{\mathcal{H}\mathcal{C}_q(b)}$, then we have

$$\frac{D_q g(z)}{D_q h(z)} \prec b_1 \frac{1+z}{1-qz}.$$

The linear transformation $w = b_1 \frac{1+z}{1-qz}$ maps $|z| = r$ onto the disc with centre $C(r) = (\frac{\alpha_1(1+qr^2)}{1-q^2r^2}, \frac{\alpha_2(1+qr^2)}{1-q^2r^2})$ and radius $\rho(r) = \frac{|b_1|(1+q)r}{1-q^2r^2}$, where $\alpha_1 = Reb_1$ and $\alpha_2 = Reb_2$. Thus using the subordination principle and the definition of the class $\mathcal{S}_{\mathcal{H}\mathcal{C}_q(b)}$, we can write

$$(2.8) \quad w_q(\mathbb{D}_r) = \left\{ \frac{D_q g(z)}{D_q h(z)} : \left| \frac{D_q g(z)}{D_q h(z)} - \frac{b_1(1+qr^2)}{1-q^2r^2} \right| \leq \frac{|b_1|(1+q)r}{1-q^2r^2}, q \in (0, 1) \right\}.$$

In order to verify Schwarz function conditions, we define the function ϕ by

$$(2.9) \quad \frac{g(z)}{h(z)} = b_1 \frac{1+\phi(z)}{1-q\phi(z)}.$$

Note that ϕ is a well defined analytic function and

$$\left. \frac{g(z)}{h(z)} \right|_{z=0} = b_1 = b_1 \frac{1+\phi(0)}{1-q\phi(0)}.$$

This proves that $\phi(0) = 0$. We now need to show that $|\phi(z)| < 1$ for all $z \in \mathbb{D}$. If we take q - derivative of both sides of (2.9) and simplify, we get

$$\frac{D_q g(z)}{h(z)} - \frac{g(qz)D_q h(z)}{h(z)h(qz)} = b_1 \frac{D_q \phi(z) - q\phi(qz)D_q \phi(z) + qD_q \phi(z) + q\phi(qz)D_q \phi(z)}{(1-\phi(z))(1-\phi(qz))}.$$

Multiplying both sides of this equation by $h(z)/D_q h(z)$ and simplifying, we obtain

$$(2.10) \quad \frac{D_q g(z)}{D_q h(z)} = b_1 \left(\frac{1+\phi(qz)}{1-q\phi(qz)} + \frac{(1+q)zD_q \phi(z)}{(1-q\phi(z))(1-q\phi(qz))} \cdot \frac{h(z)}{zD_q h(z)} \right).$$

Applying Lemma 2.2 in the equation (2.10), we can write the following form

$$(2.11) \quad \frac{D_q g(z)}{D_q h(z)} = b_1 \left(\frac{1+\phi(qz)}{1-q\phi(qz)} + \frac{(1+q)zD_q \phi(z)}{(1-q\phi(z))(1-q\phi(qz))} (1-q\phi(z))^{\frac{1-q^2}{q^2 \log q - 1}} \right).$$

Assume to the contrary that there exists a point $z_0 \in \mathbb{D}_r$ such that $|\phi(z_0)| = 1$. In view of Lemma 2.3, equation (2.11) gives

$$\frac{D_q g(z_0)}{D_q h(z_0)} = b_1 \left(\frac{1+\phi(qz_0)}{1-q\phi(qz_0)} + \frac{(1+q)m\phi(z_0)}{(1-q\phi(z_0))(1-q\phi(qz_0))} (1-q\phi(z_0))^{\frac{1-q^2}{q^2 \log q - 1}} \right) \notin w_q(\mathbb{D}_r).$$

This contradicts our assumption (2.8) and therefore $|\phi(z)| < 1$ for all $z \in \mathbb{D}$. This completes the proof of our theorem. \square

2.5. Corollary. If $f = h + \bar{g} \in \mathcal{S}_{\mathcal{H}\mathcal{C}_q(b)}$, then we have

$$(2.12) \quad F_2(|b|, Reb, |b_1|, q, r) \leq |D_q g(z)| \leq F_1(|b|, Reb, |b_1|, q, r),$$

where

$$F_1(|b|, Reb, |b_1|, q, r) = \left[(1-qr)^{Reb+|b|} (1+qr)^{Reb-|b|} \right]^{-\frac{1-q^2}{2q^2 \log q - 1}} \frac{|b_1|(1+r)}{1-qr},$$

$$F_2(|b|, Reb, |b_1|, q, r) = \left[(1-qr)^{Reb-|b|} (1+qr)^{Reb+|b|} \right]^{-\frac{1-q^2}{2q^2 \log q - 1}} \frac{|b_1|(1-r)}{1+qr}.$$

Proof. Since $f = h + \bar{g}$ is an element of $\mathcal{S}_{\mathcal{HC}_q(b)}$, from Theorem 2.4 we write $\frac{D_q g(z)}{D_q h(z)} \prec b_1 \frac{1+z}{1-qz}$, where $h \in \mathcal{C}_q(b)$. Therefore we have

$$\left| \frac{D_q g(z)}{D_q h(z)} - \frac{b_1(1+qr^2)}{1-q^2r^2} \right| \leq \frac{|b_1|(1+q)r}{1-q^2r^2}.$$

This inequality yields

$$|D_q g(z)| \leq |D_q h(z)| \frac{|b_1|(1+r)}{1-qr}.$$

If we use Lemma 1.2, we get the right side of (2.12). Similarly, we can prove the other side of the inequality (2.12). \square

2.6. Corollary. If $f = h + \bar{g} \in \mathcal{S}_{\mathcal{HC}_q(b)}$, then we have

$$(2.13) \quad f = h(z) + \overline{h(z)b_1 \frac{1+\phi(z)}{1-q\phi(z)}},$$

where ϕ is a Schwarz function.

Proof. Using Theorem 2.4, then we can write

$$\frac{g(z)}{h(z)} \prec b_1 \frac{1+z}{1-qz} \Rightarrow \frac{g(z)}{h(z)} = b_1 \frac{1+\phi(z)}{1-q\phi(z)}.$$

Therefore we obtain

$$g(z) = h(z)b_1 \frac{1+\phi(z)}{1-q\phi(z)},$$

which gives (2.13). \square

References

- [1] Andrews, G.E. *Applications of basic hypergeometric functions*, SIAM Rev. 16 (1974), 441-484.
- [2] Çetinkaya, A. and Mert, O. *A certain class of harmonic mappings related to functions of bounded boundary rotation*, Proc. of 12th Symposium on Geometric Function Theory and Applications (2016), 67-76.
- [3] Duren, P. *Harmonic mappings in the plane*, Cambridge Tracts in Math. 2004.
- [4] Fine, N.J. *Basic hypergeometric series and applications*, Math. Surveys Monogr. 1988.
- [5] Gasper, G. and Rahman, M. *Basic hypergeometric series*, Cambridge University Press, 2004.
- [6] Goodman, A.W. *Univalent functions Volume I and II*, Polygonal Pub. House, 1983.
- [7] Jack, I.S. *Functions starlike and convex of order α* , J. Lond. Math. Soc. (2), 3 (1971), 469-474.
- [8] Jackson, F.H. *On q -functions and a certain difference operator*, Trans. Roy. Soc. Edin. 46 (1908), 253-281.
- [9] Jackson, F.H. *On q -difference integrals*, Quart. J. Pure Appl. Math. 41 (1910), 193-203.
- [10] Kac, V. and Cheung, P. *Quantum calculus*, Springer, 2001.
- [11] Lewy, H. *On the non-vanishing of the Jacobian in certain one-to-one mappings*, Bull. Amer. Math. Soc. 42 (1936), 689-692.
- [12] Polatoğlu, Y., Aydoğan, M. and Mert, O. *Some properties of q -convex functions of complex order*, Sarajevo J. Math. Submitted, 2016.