

PAPER DETAILS

TITLE: The fatty acid composition of *Nigella sativa* from Turkey

AUTHORS: Mozhgan ZARIFIKHOSROSHAHI, Münüre TANUR ERKOYUNCU, Zeynep ERGUN

PAGES: 39-43

ORIGINAL PDF URL: <https://dergipark.org.tr/tr/download/article-file/2676826>

The Fatty Acid Composition of *Nigella sativa* from Turkey

Mozhagn ZARIFKHOSROSHAHI^{1*}

Münüre TANUR ERKOYUNCU²

Zeynep ERGUN³

¹Department of Horticulture, Faculty of Agriculture, Yozgat Bozok University, Yozgat, Turkey.

²Department of Field crops, Selcuk University, Konya, Turkey.

³Department of Bioengineering, Faculty of Engineering, Adana Alparslan Türkeş Science and Technology University, Adana, Turkey.

*Correspondence: mn_zarifi@yahoo.com

Citation

Zarifkhosroshahi, M., Tanur Erkoyncu, M., Ergun, Z. (2022). The Fatty Acid Composition of *Nigella sativa* from Turkey. Int J Agric For Life Sci (2022) 6(2): 39-43

Received: 28 September 2022

Accepted: 15 December 2022

Published Online: 19 December 2022

Year: 2022

Volume: 6

Issue: 2 (December)

Pages: 39-43



This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY-NC) license

<https://creativecommons.org/licenses/by-nc/4.0/>

Copyright © 2022

International Journal of Agriculture, Forestry and Life Sciences; Edit Publishing, Eskişehir, Türkiye.

Available online

<https://dergipark.org.tr/ijafsl>

Abstract

Nigella sativa, “black seed” or “black cumin” belongs to the Ranunculaceae family. It has been applied in folk medicine since ancient times. The phytochemical contents of black cumin are of interest to researchers in agriculture, pharmacology, and energy. Nowadays, it is so popular to be considered losing weight diets. In this study, the oil content and fatty acid composition of a commercial genotype of *Nigella sativa* L. from Turkey besides the fixed oil antioxidant activity were evaluated. Omega-6 fatty acid was the major fatty acid in the seed oil of studied black cumin, followed by Omega-9, free acid. However, the major composition of the oil was unsaturated fatty acids and could be considered a healthy fat for daily consumption. Moreover, the seed oil showed relatively high antioxidant activity by DPPH scavenging capacity.

Key words

Black cumin, Fatty acids, Antioxidants, Fat.

Introduction

The presence of medicinal plants in nature is one of the great blessings of God. Since man is a part of nature, nature has certainly provided medicinal plants for every disease. For this reason, the more human beings turn to nature and benefit from its blessings, the faster, better and more assuredly they will be treated for their illness. One of the magic medicinal plants which have been used throughout history is black cumin. Black cumin (*Nigella sativa* L.) is an important perennial medicinal and aromatic plant from the Ranunculaceae family and is one of the three commonly known species of the *Nigella* genus.

The plant, also known as “karamuk” and “siyah susam” in Turkish, is an annual and herbaceous plant that grows to about 20-30 cm and has black seeds. Among 12 different species of black cumin grown in the rich flora of Turkey, only *Nigella sativa* L. is cultivated (Baytop, 1984; Aslan, 2019).

The seeds of black caraway are widely used as spices besides their application in the food, medicine, and cosmetics industries (Baytöre, 2011).

Black cumin is a miraculous medicinal plant with pharmacological, biological, and therapeutic properties due to containing compounds such as fixed oils, essential oils, alkaloids, saponins, and proteins (Moghimi et al., 2018; Mukhtar et al., 2019). Based on previous studies, the seeds of black cumin have been accepted as an important medicine in Asian and Middle Eastern countries for centuries in traditional medicine and are useful for many diseases such as chronic headache, backache, hypertension, diabetes, infection, paralysis, and inflammation (Randhawa and Alghamdi, 2011; Yimer ve ark., 2019). Besides its seed oil which contains high levels of oleic, linoleic, and linolenic fatty acids has antioxidant, anti-inflammatory, anticancer, antidiabetic, antitumoral, antimicrobial, and immune-stimulating activities and gastroprotective properties. (Khan et al., 2011; Ahmad et al., 2013; Adam et al., 2016; Mazaheri et al., 2019). In Turkey, the plant is grown extensively in Mersin, Burdur, Istanbul, Gaziantep, Amasya, and Kahramanmaraş provinces in our country (Baytöre, 2011; Kılıç and Arabacı, 2016).

In a study by Piras et al. (2013), linoleic acid was higher than other fatty acids in three Turkish and one Egyptian local breeds of *N. Sativa*. Based on previous studies, the essential oil composition and fatty acid profiles differ by region (Hosseini et al., 2019). Moreover, studies reported that *N. sativa* L. seed oil can be a good source of biodiesel replacing fossil fuel which is a big challenge for fighting global warming besides meeting increasing population needs. *N. sativa* L. seed oil had a

As black caraway is a precious plant with pharmaceutical properties and is consumed in daily health nutrition in Turkey, this study aimed to investigate the fatty acid composition of commercial genotypes of black caraway from Turkey.

The physicochemical and nutritional characteristics of each oilseed are influenced by its fatty acid composition. The total fat, fatty acid composition, and antioxidant activity of the studied black caraway are shown in Table 1. The total fat obtained from the seeds of black caraway was 39.5 g/100g. Hosseini et al. (2019) determined the total fat of 16 black caraway from

different countries between 27.57-33.04 g/100g. Gharby et al., (2015) obtained the oil yield of black cumin 37% and 27% from solvent or cold press, respectively. Similarly, Atta (2003) compared “solvent” or “cold press” extraction methods and yielded 24.76% and 34.78% of oil content in black cumin, respectively. However, many studies on the total fat content of black cumin from different regions generally showed that the fat content differs in the range of %20-%30 (Burits and Bucar, 2000; Hamrouni-Sellami et al., 2008; Cheikh-Rouhou et al., 2007; Sultan et al., 2009; Ali et al., 2012; Hosseini et al., 2019; Kiani et al., 2020). The results obtained in this study are the highest among results from previous studies. As reported solvent extraction method was more successful in the determination of total fat in black cumin as it was the preferred method in this study. The high amount of obtained total fat may be due to the extraction method with equipped soxhlet besides the genotype and location of the material.

Totally 10 saturated and five unsaturated fatty acids were detected in the studied genotype of *Nigella sativa* L. The major fatty acids in *Nigella sativa* were found as linoleic, oleic, and palmitoleic acids (52.89%, 25.12%, and 13.62%), respectively. Approximately half percent of seed oils were polyunsaturated fatty acids mainly linoleic acid. The results are in agreement with previous studies (Burits and Bucar, 2000; Hamrouni-Sellami et al., 2008; Cheikh-Rouhou et al., 2007; Sultan et al 2009; Ali et al., 2012; Hosseini et al. 2019; Kiani et al., 2020). A high concentration of linoleic acid is a determining indicator of the nutritional value of plant oils and has healthy effects by reducing serum cholesterol leading to a remedy for cardiovascular disorders (Al-Jassir, 1992; Keles, 2020; Guney, 2020; Ergun 2021). Most percentages of seed oil (80.86%) in studied *Nigella sativa* L. were comprised of unsaturated fatty acids. Palmitoleic acid was found as the main SFA in the seed oil of black cumin. Likewise, measurable amounts of caproic acid (C6:0) an SFA, eicosenoic acid (C20:1n9c), cis-11,14-eicosadienoic acids (C21:2) MUFAs were detected. Pereira et al., (2014) reported

that unsaturated fatty acids eicosenoic and cis-11,14-eicosadienoic acids had inflammatory effects. However, genetic factors (cultivar, variety grown), environmental conditions, seed quality (maturity, harvesting, or storage conditions), oil extraction methods, and esterification methods during the determination of fatty acids may be the cause of possible differentiation among results from different studies on oil content and fatty acid composition of *Nigella sativa* L. (Ramadan and Mörsel 2002; Kiani et al., (2020).

Free radicals generated through routine metabolism pathways may be the basis of many human diseases (Gundesli 2020; Okatan et al., 2021). Antioxidants have the ability to scavenge free radicals by exchanging electrons (Gundesli et al., 2020; 2021). 4-terpineol, t-anethole, and thymoquinone carvacrol isolated from the essential oil of *N. sativa* have considerable free radical scavenging properties (Ali and Blunden, 2003). There are several studies on the antioxidant capacity of black cumin oil. Among various antioxidant capacity measuring assays, DPPH has been widely used in the scavenging assessment of free radicals due to being an easy method (Erkan et al., 2008; Scherer and Godoy, 2009). Erkan et al., (2008) reported that black cumin seeds inhibited DPPH radical formation as 515 of IC50 (μM) mean. Sultan et al (2009) also studied the antioxidant capacity of fixed and essential oils of black cumin via DPPH assay and reported their inhabitation strength of DPPH as 32.32% and 80.25%, respectively. In, this study, the DPPH antioxidant activity of black cumin was determined as 284.70 mg TE/100g. Al Turkmani et al (2015) reported that scavenging DPPH radicals of *N. sativa* essential oil showed a significant difference in samples from two different regions in Syria as 0.607 mg⁻¹ml and 0.240 mg⁻¹ml. Although there are differences in the DPPH antioxidant activity among previous studies, the results are in agreement with each other based on affecting factors such as region, genotype, and extraction solvents.

Table 1. Fatty acid composition, total fat, and antioxidant activity of *Nigella sativa* L.

Fatty Acids	%
Caproic acid (C6:0)	0.10±0.01
Caprylic acid (C8:0)	0.04±0.01
Myristic Acid (C14:0)	0.36±0.01
Pentadecanoic acid (C15:0)	0.03±0.00
Palmitic acid (C16:0)	13.62±0.36
Palmitoleic acid (C16:1)ω-7	0.17±0.00
Margaric Acid (C17:0)	0.07±0.00
Stearic acid (C18:0)	3.55±0.01
Arachidic acid (C20:0)	0.21±0.00
Behenic acid (C22:0)	0.05±0.01
ΣSFA	18.2
Oleic acid (C18:1n9c)ω-9	25.12±0.06
Eicosenoic acid (C20:1n9c)ω-9	0.32±0.01
cis-11,14-Eicosadienoic acid (C21:2)	2.33±0.04
ΣMUFA	27.77
Linoleic acid (C18:2n6c) ω-6	52.89±0.01
α-Linolenic acid (C18:3n3) ω-3	0.20±0.01
ΣPUFA	53.09
Total Fat	39.50 g/100g
DPPH Antioxidant Activity	284.70±8.90 mg TE/100g

Conclusion

The present study revealed that the commercial genotype from Turkey has a high content of fat besides a good source of linoleic–oleic type fatty acids which are so important for a healthy diet in the cosmetic industry. The study supported the findings of previous studies on the rich phytochemical content of *N.sativa* showing that commercial genotypes also can be used in breeding programs. However, studies on trueness to the name of these genotypes besides the breeding of genotypes with higher content of bioactive compounds should be carried out

Statement of Conflict of Interest

The author(s) declare no conflict of interest for this study.

Author's Contributions

Z.E. and M.Z. contributed equally in fatty acid analyses and the writing of the manuscript. M.T.E helped in writing the manuscript. The authors contributed to the discussion of the results and all of them read and approved the final manuscript.

References

- Adam, G. O., Rahman, M. M., Lee, S. J., Kim, G. B., Kang, H. S., Kim, J. S., Kim, S. J. (2016). Hepatoprotective effects of *Nigella sativa* seed extract against acetaminophen-induced oxidative stress. *Asian Pacific Journal of Tropical Medicine*, 9(3):221-227.
- Aghabarari, B., Dorostkar, N., Martinez-Huerta, M. V. (2014). Synthesis of biodiesel from *Nigella sativa* seed oil using surfactant-Brønsted acidic-combined ionic liquid as catalyst. *Fuel processing technology*, 118:296-301.
- Ahmad, A., Husain, A., Mujeeb, M., Khan, S. A., Najmi, A. K., Siddique, N. A., ... Anwar, F. (2013). A review on therapeutic potential of *Nigella sativa*: A miracle herb. *Asian Pacific journal of tropical biomedicine*, 3(5):337-352.
- Al Turkmani, M. O., Karabet, F., Mokrani, L., Soukkarieh, C. (2015). Chemical composition and *in vitro* antioxidant activities of essential oil from *Nigella sativa* L. seeds cultivated in Syria. *International Journal of ChemTech Research*, 8(10):76-82.
- Ali, B. H., Blunden, G. (2003). Pharmacological and toxicological properties of *Nigella sativa*. *Phytotherapy Research: An international journal devoted to pharmacological and toxicological evaluation of natural product derivatives*, 17(4):299-305.
- Ali, M. A., Sayeed, M. A., Alam, M. S., Yeasmin, M. S., Khan, A. M., Muhamad, I. I. (2012). Characteristics of oils and nutrient contents of *Nigella sativa* Linn. and *Trigonella foenum-graecum* seeds. *Bulletin of the Chemical Society of Ethiopia*, 26(1).
- Al-Jassir, M. S. (1992). Chemical composition and microflora of black cumin (*Nigella sativa* L.) seeds growing in Saudi Arabia. *Food Chemistry*, 45(4):239-242.
- Aslan, R. (2019). Siyah Reçete: Çörek Otu (*Nigella Sativa*). *Ayrıntı Dergisi*, 7(74).
- Atta, M. B. (2003). Some characteristics of nigella (*Nigella sativa* L.) seed cultivated in Egypt and its lipid profile. *Food chemistry*, 83(1):63-68.
- Aziz, S. A., Kurniawati, A., Faridah, D. N. (2017). Changes of thymoquinone, thymol, and malondialdehyde content of black cumin (*Nigella sativa* L.) in response to Indonesia tropical altitude variation. *Hayati Journal of Biosciences*, 24(3):156-161.
- Baytop T, 1984. Türkiye’de Bitkiler İle Tedavi (Geçmişte ve Bugün). İstanbul Üniversitesi Yayınları No: 3255, Sanal Matbaacılık, İstanbul, 520 s.
- Baytöre F, 2011. Bazı Çörekotu (*Nigella sativa* L.) Populasyonlarının Verim ve Verim Kriterlerinin Belirlenmesi. Namık Kemal Üniversitesi Fen Bilimleri Enstitüsü, Yüksek Lisans Tezi, Tarla Bitkileri Anabilim Dalı, Tekirdağ.
- Bozkurt, T., Ergun, Z. (2021). Fatty acid composition and antioxidant capacity of pomegranate seed oil. *GSC Biological and Pharmaceutical Sciences*, 15(2):103-110.
- Burits, M., Bucar, F. (2000). Antioxidant activity of *Nigella sativa* essential oil. *Phytotherapy research*, 14(5):323-328.
- Cheikh-Rouhou, S., Besbes, S., Hentati, B., Blecker, C., Deroanne, C., Attia, H. (2007). *Nigella sativa* L.: Chemical composition and physicochemical characteristics of lipid fraction. *Food chemistry*, 101(2):673-681.
- Darakhshan, S., Pour, A. B., Colagar, A. H., Sisakhtnezhad, S. (2015). Thymoquinone and its therapeutic potentials. *Pharmacological research*, 95:138-158.
- David, F., Sandra, P., Wylie, P. (2003). Improving the Analysis of Fatty Acid Methyl Esters Using Retention Time Locked Method and Retention Time Databases,” Application Note 5990-4822EN, Agilent Technologies publication 5988-5871EN.
- De Souza, C. O., Vannice, G. K., Rosa Neto, J. C., Calder, P. C. (2018). Is palmitoleic acid a plausible nonpharmacological strategy to prevent or control chronic metabolic and inflammatory disorders?. *Molecular Nutrition Food Research*, 62(1):1700504.
- Ergun, Z. (2021). Seed oil content and fatty acid profiles of endemic Phoenix theophrasti Greuter, Phoenixroebelenii O'Brien, Phoenix canariensis Hort. Ex Chabaud, and Phoenix dactylifera L. grown in the same locality in Turkey. *Turkish Journal of Agriculture and Forestry*, 45(5):557-564.
- Ergun, Z., Bozkurt, T. (2020). Determination of fatty acid composition and antioxidant activity of fig seed oil. *International Journal of Agricultural and Natural Sciences*, 13(2):101-107.
- Erkan, N., Ayranci, G. Ayranci, E.(2008). Antioxidant activities of rosemary (*Rosmarinus officinalis* L.) extract, blackseed (*Nigella sativa* L.) essential oil, carnosic acid, rosmarinic acid and sesamol, *Food Chem.*, 110: 76-82.
- Gharby, S., Harhar, H., Guillaume, D., Roudani, A., Boulbaroud, S., Ibrahimi, M., ... Charrouf, Z. (2015). Chemical investigation of *Nigella sativa* L. seed oil produced in

- Morocco. Journal of the Saudi Society of Agricultural Sciences, 14(2):172-177.
- Gholamnezhad, Z., Keyhanmanesh, R., Boskabady, M. H. (2015). Anti-inflammatory, antioxidant, and immunomodulatory aspects of *Nigella sativa* for its preventive and bronchodilatory effects on obstructive respiratory diseases: A review of basic and clinical evidence. Journal of Functional Foods, 17:910-927.
- Guney, M. (2020). Determination of fatty acid profile and antioxidant activity of Rosehip seeds from Turkey. International Journal of Agriculture Environment and Food Sciences, 4(1):114-118.
- Gundesli, M.A., (2020). Determination of Sugar contents, Total Phenol and Antioxidant Activity of various parts 'Uzun' pistachio cultivar (*Pistacia vera* L.). International Journal of Agriculture Environment and Food Sciences, 4(1):52-58.
- Gundesli, M.A., Kafkas, N.E., Güney, M., Ercisli, S. (2021). Determination of phytochemicals from fresh fruits of fig (*Ficus carica* L.) at different maturity stages. Acta Sci. Pol. Hortorum Cultus, 20(2):73–81.
- Gundesli, M.A., Kafkas, N.E., Kafkas, S., Guney, M., Okatan, V., Usanmaz, S., (2020). Identification and Characterisation of Volatile Compounds Determined by HS/GC-MS Technique in Pulp of 'Abbas' Fig (*Ficus carica* L.) Variety. Pakistan Journal of Agricultural Sciences. 57(3):623-629.
- Hamrouni-Sellami, I. B. T. I. S. S. E. M., Elyes Khouk, M. O. H. A. M. E. D., Marzouk, B. (2008). Lipid and aroma composition of black cumin (*Nigella sativa* L.) seeds from Tunisia. Journal of Food Biochemistry, 32(3):335-352.
- Hosseini, S. S., Rezadoost, H., Nadjafi, F., Asareh, M. H. (2019). Comparative essential oil composition and fatty acid profiling of some Iranian black cumin landraces. Industrial Crops and Products, 140:111628.
- Keles, H. (2020). Changes of some horticultural characteristics in jujube (*Ziziphus jujube* Mill.) fruit at different ripening stages. Turkish Journal of Agriculture and Forestry, 44(4):391-398.
- Khan, A., Chen, H. C., Tania, M., Zhang, D. Z. (2011). Anticancer activities of *Nigella sativa* (black cumin). African Journal of Traditional, Complementary and Alternative Medicines, 8(5S).
- Khan, T. Y., Atabani, A. E., Badruddin, I. A., Ankalgi, R. F., Khan, T. M., Badarudin, A. (2015). Ceiba pentandra, *Nigella sativa* and their blend as prospective feedstocks for biodiesel. Industrial Crops and Products, 65:367-373.
- Kılıc, C., Arabaci, O. (2016). Çörek Otu (*Nigella Sativa* L.)'Nda Farklı Ekim Zamanı Ve Tohumluk Miktarının Verim Ve Kaliteye Etkisi. Adnan Menderes Üniversitesi Ziraat Fakültesi Dergisi, 13(2):49-56.
- Kiani, M., Alahdadi, I., Soltani, E., Boelt, B., Benakashani, F. (2020). Variation of seed oil content, oil yield, and fatty acids profile in Iranian *Nigella sativa* L. landraces. Industrial crops and products, 149:112367.
- Kostiæ, D. A., Dimitrijeviæ, D. S., Mitiaæ, S. S., Mitiaæ, M. N., Stojanoviæ, G. S., Živanoviæ, A. V. (2013). Phenolic content and antioxidant activities of fruit extracts of *Morus nigra* L (Moraceae) from Southeast Serbia. Tropical Journal of Pharmaceutical Research, 12(1):105-110.
- Maideen, N. M. P. (2020). Prophetic medicine-*Nigella Sativa* (Black cumin seeds)–potential herb for COVID-19?. Journal of pharmacopuncture, 23(2), 62.
- Mazaheri, Y., Torbati, M., Azadmard-Damirchi, S., Savage, G. P. (2019). Effect of roasting and microwave pre-treatments of *Nigella sativa* L. seeds on lipase activity and the quality of the oil. Food Chemistry, 274:480-486.
- Moghimi, M., Farzaneh, V., Bakhshabadi, H. (2018). The effect of ultrasound pretreatment on some selected physicochemical properties of black cumin (*Nigella Sativa*). Nutrire, 43(1):1-8.
- Mukhtar, H., Qureshi, A. S., Anwar, F., Mumtaz, M. W., Marcu, M. (2019). *Nigella sativa* L. seed and seed oil: Potential sources of high-value components for development of functional foods and nutraceuticals/pharmaceuticals. Journal of Essential Oil Research, 31(3):171-183.
- Okatan, V., Bulduk, I., Kaki, B., Gundesli M.A. Usanmaz, S., Alas, T., Helvacı, M., Kahramanoğlu, I., Hajizadeh, H.S., (2021). Identification and Quantification of Biochemical Composition and Antioxidant Activity of Walnut Pollens. Pak. J. Bot, 53(6): 2241-2250.
- Pereira, D. M., Correia-da-Silva, G., Valentão, P., Teixeira, N., Andrade, P. B. (2014). Anti-inflammatory effect of unsaturated fatty acids and ergosta-7, 22-dien-3-ol from *Marthasterias glacialis*: Prevention of CHOP-mediated ER-stress and NF-κB activation. PLoS One, 9(2):e88341.
- Piras, A., A. Rosa, Marongiu, B., S. Porcedda, Falconieri, D., M.A. Dessi, Ozcelik, B., U. Koca. (2013). Chemical Composition and In Vitro Bioactivity of the Volatile and Fixed Oils of *Nigella sativa* L. Extracted by Supercritical Carbon Dioxide. Industrial Crops and Products. 46: 317-323.
- Ramadan, M.F., Mörsel, J.T. (2002). Neutral lipid classes of black cumin (*Nigella sativa* L.) seed oils. Eur. Food Res. Technol. 214:202– 206.
- Randhawa MA, Alghamdi MS. (2011). Anticancer activity of *Nigella sativa* (black seed) —A review. Am J Chin Med. 39(6): 1075-1091.
- Salem, M. L. (2005). Immunomodulatory and therapeutic properties of the *Nigella sativa* L. seed. International immunopharmacology, 5(13-14):1749-1770.
- Scherer, R. and H.T. Godoy. 2009. Antioxidant activity index (AAI) by the 2,2-diphenyl-1- picrylhydrazyl method. Food Chem., 112: 654-658
- Sultan, M. T., Butt, M. S., Anjum, F. M., Jamil, A., Akhtar, S., Nasir, M. (2009). Nutritional profile of indigenous cultivar of black cumin seeds and antioxidant potential of its fixed and essential oil. Pak. J. Bot, 41(3):1321-1330.
- Yimer, E. M., Tuem, K. B., Karim, A., Ur-Rehman, N., Anwar, F. (2019). *Nigella sativa* L.(black cumin): a promising natural remedy for wide range of illnesses. Evidence-Based Complementary and Alternative Medicine, 2019.