# PAPER DETAILS

TITLE: A Boundary Value Problem with Retarded Argument Containing an Eigenparameter in the

Transmission Condition

AUTHORS: Özge Isik Gülmezler,Özgür Mizrak

PAGES: 282-288

ORIGINAL PDF URL: https://dergipark.org.tr/tr/download/article-file/3440906



## A Boundary Value Problem with Retarded Argument Containing an Eigenparameter in the Transmission Condition

Özge Işık Gülmezler, Özgür Mızrak\*

Mersin University, Faculty of Science, Departmant of Mathematics, Mersin, Türkiye ozgeisik58@gmail.com<sup>®</sup>, \*mizrak@mersin.edu.tr <sup>®</sup> Received date:28.09.2023, Accepted date: 17.10.2023

#### Abstract

In this work, a discontinuous boundary value problem with retarded argument is studied. At the discontinuity point there is a transmission condition that contains a parameter. Asymptotic properties of eigenvalues and corresponding eigenfunctions of the boundary value problem are studied.

Keywords: Retarded argument, transmission condition, eigenvalue, eigenfunction, boundary value problem

## Geçiş koşulunda Özdeğer Parametresi İçeren Geç Kalan Argümanlı Bir Sınır Değer Problemi

Öz

Bu çalışmada, süreksizlik içeren geç kalan argümanlı bir sınır değer problemi ele alınmıştır. Süreksizlik noktasında parametre içeren bir geçiş koşulu vardır. Ele alınan sınır değer probleminin özdeğer ve özfonksiyonlarının asimptotik özellikleri incelenecektir.

Anahtar Kelimeler: Geç kalan argüman, geçiş koşulu, özdeğer, özfonksiyon, sınır değer problemi

#### **INTRODUCTION**

Differential equations with retarded argument becomes the popular branches of functional differential equations. After the development of control systems in engineering retarded equations become important. Before that scientists were aware of this type of delays in the control systems but there was not enough theory about this subject. Because of that this type of affects were ignored in the models. Delays have an important role to explain complex models mathematically and it also has important affects.

Standart problem in this area was given in 1956 by Norkin (Norkin,1956). And in 1958 Norkin studied the same equation with more general boundary conditions(Norkin, 1958). Most of the work done in this area is seperated into two parts as continuous and discontinuous problems. Continuous problems have either standart boundary conditions or eigenparameters at the boundary conditions (Norkin, 1972; Bayramoğlu, Köklü, Baykal 2002; Koparan 2019) Discontinuous problems varries as having eigenparameter at the boundary condition and having transmission conditions at the discontinuity points (Şen, Bayramov 2011; Şen, Bayramov 2011b; Yang 2012; Aydin Akgun, Bayramov, Bayramoğlu 2013; Şen, Seo, Araci 2013, Hira 20017). There is only one work done about the problem that has eigenparameter at the transmission condition (Şen, Bayramov 2013). Applications of differential equations with retarded argument is given in the book by Kolmanovskii, and Myshkis (Kolmanovskii and Myshkis, 1999).

In this work, below differential equation with retarded argument will be studied. Here the problem has discontinuity and at the discontinuity point transmission conditions have eigenparameter.

$$y''(x) + \lambda^2 y(x) + q(x)y(x - \Delta(x)) = 0, \qquad x \in \begin{bmatrix} 0 & \pi \end{bmatrix}$$

$$\left[0,\frac{-}{2}\right) \cup \left(\frac{-}{2},\pi\right] \tag{1}$$

$$y(0) + \alpha y'(0) = 0,$$
 (2)

$$y(\pi) = 0, \tag{3}$$

and the transmission conditions

$$y\left(\frac{\pi}{2}+0\right) = \frac{\delta}{\lambda}y\left(\frac{\pi}{2}-0\right),\tag{4}$$

$$y'\left(\frac{\pi}{2}+0\right) = \frac{\delta}{\lambda}y'\left(\frac{\pi}{2}-0\right),\tag{5}$$

where q(x) and  $\Delta(x) \ge 0$  are real valued functions continuous in  $\left[0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right]$  and have finite left and right limits at each side of  $\frac{\pi}{2}$ , if  $x \in \left[0, \frac{\pi}{2}\right)$  then  $x - \Delta x \ge 0$ , if  $x \in \left(\frac{\pi}{2}, \pi\right]$  then  $x - \Delta x \ge \frac{\pi}{2}$ ,  $\lambda$  is an eigenparameter and  $\alpha, \delta \ne 0$  are arbitrary real numbers.

Consider the solution of equation (1) on  $\left[0, \frac{\pi}{2}\right]$  such that it satisfies the initial conditions

$$\omega_1(0,\lambda) = \alpha, \qquad \omega'_1(0,\lambda) = -1. \tag{6}$$

These initial conditions (6) define a unique solution of equation (1) on  $\left[0,\frac{\pi}{2}\right)$  (Norkin,1972). By transmission conditions, solution of equation (1) on  $\left(\frac{\pi}{2},\pi\right]$  can be defined in terms of  $\omega_1(x,\lambda)$ .  $\omega_2(x,\lambda)$ can be written as

$$\omega_2\left(\frac{\pi}{2},\lambda\right) = \frac{\delta}{\lambda}\omega_1\left(\frac{\pi}{2},\lambda\right),\tag{7}$$

$$\omega'_{2}\left(\frac{\pi}{2},\lambda\right) = \frac{\sigma}{\lambda}\omega'_{1}\left(\frac{\pi}{2},\lambda\right). \tag{8}$$

These initial contions (7) define a unique solution of equation (1) on  $\left(\frac{\pi}{2}, \pi\right]$ .

Now, we can define the function  $\omega(x, \lambda)$  on  $\left[0, \frac{\pi}{2}\right] \cup \left(\frac{\pi}{2}, \pi\right]$  such that

$$\omega(x,\lambda) = \begin{cases} \omega_1(x,\lambda), & x \in [0\frac{\pi}{2}] \\ \omega_2(x,\lambda), & x \in (\frac{\pi}{2}\pi] \end{cases}.$$
(9)

This function is a solution of equation (1) on  $\left[0,\frac{\pi}{2}\right] \cup \left(\frac{\pi}{2},\pi\right]$ , that satisfies left boundary condition (2) and both transmission conditions (4) and (5).

**Lemma 1:** Let  $\omega(x, \lambda)$  be a solution of equation (1) and  $\lambda > 0$ . Then  $\omega_1(x, \lambda)$  and  $\omega_2(x, \lambda)$  have the following forms:

$$\omega_{1}(x,\lambda) = \alpha cos\lambda x - \frac{sin\lambda x}{\lambda} + \frac{1}{\lambda} \int_{0}^{x} q(\tau) sin \lambda (x-\tau) \omega_{1}(\tau - \Delta(\tau), \lambda) d\tau,$$
(10)

$$\omega_{2}(x,\lambda) = \frac{\delta}{\lambda} \omega_{1}\left(\frac{\pi}{2},\lambda\right) cos\lambda\left(x-\frac{\pi}{2}\right) + \\ + \frac{\delta}{\lambda^{2}} \omega_{1}'\left(\frac{\pi}{2},\lambda\right) sin\lambda(x-\frac{\pi}{2}) + \\ + \frac{1}{\lambda} \int_{\frac{\pi}{2}}^{x} q(\tau) sin\lambda(x-\tau) \omega_{2}(\tau-\Delta(\tau),\lambda) d\tau.$$
(11)

**Proof:** By differentiating  $\omega_1(x,\lambda)$  and  $\omega_2(x,\lambda)$  twice and applying integration by parts we see that (10) and (11) satisfy equation (1).

**Theorem 1:** The eigenvalue problem (1)-(5) have only simple eigenvalues.

**Proof:** Let  $\tilde{\lambda}$  be an eigenvalue of the problem (1)-(5) and

$$\tilde{u}(x,\tilde{\lambda}) = \begin{cases} \tilde{u}_1(x,\tilde{\lambda}), & x \in [0\frac{\pi}{2}] \\ \tilde{u}_2(x,\tilde{\lambda}), & x \in (\frac{\pi}{2}\pi] \end{cases}$$

be a corresponding eigenfunction. Then from (2) and (6) it follows that the Wronskian

$$W[\widetilde{u}_1(0,\widetilde{\lambda}),\widetilde{\omega}_1(0,\widetilde{\lambda})] = \begin{vmatrix} \widetilde{u}_1(0,\widetilde{\lambda}) & \alpha \\ \widetilde{u}'_1(0,\widetilde{\lambda}) & -1 \end{vmatrix} = 0.$$

Then it means that  $\tilde{u}_1(x, \tilde{\lambda})$  and  $\tilde{\omega}_1(0, \tilde{\lambda})$  are linearly dependent on  $[0\frac{\pi}{2}]$ . Similarly it can be proved that  $\tilde{u}_2(x, \tilde{\lambda})$  and  $\tilde{\omega}_2(0, \tilde{\lambda})$  are linearly dependent. Consequently, it follows that  $\omega(x, \tilde{\lambda})$  is an eigenfunction for the boundary value problem (1)-(5) and all the eigenfunctions to this problem corresponding to  $\tilde{\lambda}$  are linearly dependent. Hence eigenvalues of the problem are simple.





### MATERIAL AND METHODS

The function  $\omega(x,\lambda)$  defined by (10) is a solution of equation (1) satisfying left boundary condition (2) and transmission conditions (4) and (5). Writing  $\omega(x,\lambda)$  into (3), characteristic equation is obtained.

$$F(\lambda) = \omega(\pi, \lambda) = \omega_2(\pi, \lambda) = 0.$$
(12)

Theorem 1 guarantees that the set of eigenvalues of the boundary value problem (1)-(5) coincides with the set of real roots of equation (12).

#### Lemma 2:

1. Let  $\lambda \ge 2q_1$ . Then  $\omega_1(x, \lambda)$  given by equation (10) satisfy the following inequality:

$$\begin{split} |\omega_1(x,\lambda)| &\leq \frac{1}{q_1} \sqrt{4q_1^2 \alpha^2 + 1}. \end{split} \tag{13}$$
  
2. Let  $\lambda \geq max\{2q_1, 2q_2\}$ . Then  $\omega_2(x,\lambda)$  given by equation (10) satisfies the following inequality:

$$|\omega_2(x,\lambda)| \le \frac{4|\delta|}{\lambda q_1} \sqrt{4q_1^2 \alpha^2 + 1}.$$
(14)

**Proof:** Let  $B_{1\lambda} = max_{[0,\frac{\pi}{2}]} |\omega_1(x,\lambda)|$ . Then from (10), it follows that for every  $\lambda > 0$ , the following inequality holds:

$$B_{1\lambda} \le \sqrt{\alpha^2 + \frac{1}{\lambda^2}} + \frac{1}{\lambda} q_1 B_{1\lambda}.$$
(15)

For  $\lambda \ge 2q_1$ , (13) is obtained. Differentiating (10) with respect to x,

$$\omega_1'(x,\lambda) = -\alpha\lambda \sin\lambda x - \cos\lambda x - -\int_0^x q(\tau)\cos\lambda(x-\tau)\omega_1(\tau-\Delta(\tau),\lambda)d\tau \quad (16)$$

is obtained. From (13) and (16), for  $\lambda \ge 2q_1$ , we have  $|\omega'_1(x,\lambda)| \le \sqrt{\alpha^2 \lambda^2 + 1} + \sqrt{4q_1^2 \alpha^2 + 1}$ .

Hence for  $\lambda \ge 2q_1$ , and  $x \in [0\frac{\pi}{2}]$ ,

$$\frac{|\omega_1'(x,\lambda)|}{\lambda} \le \frac{1}{q_1}\sqrt{4q_1^2\alpha^2 + 1}.$$
(17)

Let  $B_{2\lambda} = max_{\left[\frac{\pi}{2},\pi\right]} |\omega_2(x,\lambda)|$ , then from (10), (13) and (17), for  $\lambda \ge 2q_1$ , the following inequality holds:

$$B_{2\lambda} \leq \frac{2|\delta|}{\lambda q_1} \sqrt{4q_1^2 \alpha^2 + 1} + \frac{1}{\lambda} q_2 B_{2\lambda}.$$
  
Hence for  $\lambda \geq max\{2q_1, 2q_2\}$ , (14) is obtained.

**Theorem 2:** Eigenvalues of the problem (1)-(5) form an infinite set of positive real numbers.

**Proof:** Differentiating (10) with respect to *x*,

$$\omega_{2}'(x,\lambda) = -\delta\omega_{1}\left(\frac{\pi}{2},\lambda\right) \sin\lambda\left(x-\frac{\pi}{2}\right) + \frac{\delta}{\lambda}\omega_{1}'\left(\frac{\pi}{2},\lambda\right) \cos\lambda\left(x-\frac{\pi}{2}\right) + \frac{1}{\lambda}\int_{\frac{\pi}{2}}^{x}q(\tau)\cos\lambda(x-\tau)\omega_{2}(\tau-\Delta(\tau),\lambda)d\tau.$$
(18)

Writing (10), (11), (16) and (18) into the characteristic equation (12), we obtain,

$$\begin{aligned} \frac{\alpha\delta}{\lambda}\cos\lambda\pi &-\frac{\delta}{\lambda^2}\sin\lambda\pi + \\ &+\frac{\delta}{\lambda^2}\int_0^{\frac{\pi}{2}}q(\tau)\sin\lambda(\pi-\tau)\omega_1(\tau-\Delta(\tau),\lambda)d\tau \\ &+\frac{1}{\lambda}\int_{\frac{\pi}{2}}^{\pi}q(\tau)\sin\lambda(\pi-\tau)\omega_2(\tau-\Delta(\tau),\lambda)d\tau = 0. \end{aligned}$$
(19)

For  $\alpha \neq 0$  and sufficiently large  $\lambda$ , by (13) and (14) equation (19) may be written as

$$\lambda \cos \lambda \pi + O(1) = 0. \tag{20}$$

Clearly there are infinitely many real numbers  $\lambda$  that satisfy (20).

# Asymptotic Formulas for Eigenvalues and Eigenfunctions

Now we will study the asymptotic behavior of eigenvalues and eigenfunctions. From now on we will assume  $\lambda$  is sufficiently large. On  $\left[0\frac{\pi}{2}\right)$ , from (9) and (13)

$$ω_1(x, \lambda) = O(1).$$
 (21)  
On  $(\frac{\pi}{2}\pi]$ , from (10) and (14)



$$\omega_2(x,\lambda) = O\left(\frac{1}{\lambda}\right). \tag{22}$$

Derivatives of of  $\omega_1(x, \lambda)$  and  $\omega_2(x, \lambda)$  with respect to  $\lambda$  on  $x \in [0\frac{\pi}{2})$ , and  $x \in (\frac{\pi}{2}\pi]$  respectively, exist and are continuous for  $|\lambda| < \infty$  by Norkin (Norkin, 1972).

### Lemma 3:

$$\omega'_{1\lambda}(x,\lambda) = O(1), for \ x \in \left[0, \frac{\pi}{2}\right], \tag{23}$$

$$\omega'_{2\lambda}(x,\lambda) = O\left(\frac{1}{\lambda}\right), \text{ for } x \in \left[\pi, \frac{\pi}{2}\right],$$
 (24)

**Proof:** Differentiating (9) with respect to  $\lambda$  and by (21),

$$\omega'_{1\lambda}(x,\lambda) = \frac{1}{\lambda} \int_0^x q(\tau) \sin \lambda(x-\tau) \omega'_{1\lambda}(\tau-\lambda) d\tau + K(x,\lambda), \quad (|K(x,\lambda)| \le K_0).$$
(25)

Let  $C_{1\lambda} = max_{[0,\frac{\pi}{2}]} |\omega'_{1\lambda}(x,\lambda)|$ . The existence of  $C_{1\lambda}$  follows from the continuity of derivative for  $x \in \left[0, \frac{\pi}{2}\right]$ . From (25)

$$C_{1\lambda} = \frac{1}{\lambda} q_1 C_{1\lambda} + K_0$$

is obtained. So for  $\lambda \ge 2q_1$ ,  $C_{1\lambda} \le 2K_0$ . Therefore (23) is proved. Similarly (24) may be proved.

**Theorem 3:** Let *n* be a sufficiently large natural number. Then there is only one eigevalue of the problem (1)-(5) around  $n + \frac{1}{2}$ .

**Proof:** First consider the O(1) term in equation (20)

$$-\frac{\lambda}{\alpha}\sin\lambda\pi + \frac{1}{\alpha}\int_{0}^{\frac{\pi}{2}}q(\tau)\sin\lambda(\pi-\tau)\omega_{1}(\tau-\Delta(\tau),\lambda)d\tau + \frac{\lambda}{\alpha}\int_{\frac{\pi}{2}}^{\frac{\pi}{2}}q(\tau)\sin\lambda(\pi-\tau)\omega_{2}(\tau-\Delta(\tau),\lambda)d\tau$$

From (21)-(24), this expression has bounded derivative with respect to  $\lambda$ . It is clear that for  $\lambda$  big enough roots of equation (20) are located close to  $n + \lambda$ 

 $\frac{1}{2}$ . We need to show that there is only one solution of (20) around each  $n + \frac{1}{2}$ .

Consider the function  $F(\lambda) = \lambda cos\lambda \pi + O(1)$ . Its derivative  $F'(\lambda) = cos\lambda \pi - \lambda \pi sin\lambda \pi + O(1)$  is equal to zero for  $\lambda$  close to  $n + \frac{1}{2}$  for sufficiently large *n*. Consequently by Rolle's Theorem proof is completed.

From equation (20) eigenvalues of the boundary value problem (1)-(5) are obtained as:

$$\lambda_n = n + \frac{1}{2} + O\left(\frac{1}{n}\right),\tag{26}$$

From equation (10), (16) and (21)

$$\omega_1(x,\lambda) = \alpha \cos \lambda x + O\left(\frac{1}{\lambda}\right),$$
 (27)

$$\omega'_1(x,\lambda) = -\alpha\lambda\sin\lambda x + O(1), \qquad (28)$$

From equation (11), (22), (27) and (28),

$$\omega_2(x,\lambda) = \frac{\alpha\delta}{\lambda}\cos\lambda x + O\left(\frac{1}{\lambda^2}\right).$$
(29)

Hence eigenfunctions  $u_n(x)$  have the following asymptotic representation.

$$u_n(x) = \begin{cases} \alpha \cos\left(n + \frac{1}{2}\right)x + O\left(\frac{1}{n}\right), \ x \in [0, \frac{\pi}{2}] \\ \frac{\delta}{n} \alpha \cos\left(n + \frac{1}{2}\right)x + O\left(\frac{1}{n^2}\right), x \in \left(\frac{\pi}{2}, \pi\right] \end{cases}$$

### **RESULTS AND DISCUSSION (Main title)**

Under additional assumptions we can improve these formulas for the eigenvalues and the eigenfunctions.

# Improved Asymptotic Representations for the Eigenvalues and Eigenfunctions

**Theorem 4:** Assume that q'(x) and  $\Delta''(x)$  exist, are bounded on  $[0, \frac{\pi}{2}) \cup (\frac{\pi}{2}, \pi]$  and have left and right limits at  $\frac{\pi}{2}$ , on  $[0, \frac{\pi}{2}) \cup (\frac{\pi}{2}, \pi]$ ,  $\Delta'(x) \le 1$ ,  $\Delta(0) = 0$ and  $\lim_{x \to \frac{\pi}{2} \neq 0} \Delta(x) = 0$ . Then for  $n \to \infty$ , eigenvalues of the problem (1)-(5) are as follows:

$$\lambda_n = n + \frac{1}{2} + \frac{\alpha B(\pi, n, \Delta(\tau)) - 1}{\alpha n \pi} + O\left(\frac{1}{n^2}\right)$$
(30)

**Proof:** It is clear that from (27) and (29)

$$\omega_1(\tau - \Delta(\tau), \lambda) = \alpha \cos \lambda(\tau - \Delta(\tau)) + O\left(\frac{1}{\lambda}\right) (31)$$
  
$$\omega_2(\tau - \Delta(\tau), \lambda) = \frac{\alpha \delta}{\lambda} \cos \lambda(\tau - \Delta(\tau)) + O\left(\frac{1}{\lambda^2}\right) (32)$$

Writing these expressions into the equation (19) we obtain

 $\lambda\alpha\delta\cos\lambda\pi - \delta\sin\lambda\pi +$ 

$$+\alpha\delta \int_{0}^{\pi} q(\tau)\sin\lambda\pi \frac{1}{2} [\cos\lambda\Delta(\tau) + \cos\lambda(2\tau - \Delta(\tau))]d\tau$$
$$-\delta \int_{0}^{\pi} q(\tau)\sin\lambda\tau \frac{1}{2} [\sin\lambda\Delta(\tau) + \sin\lambda(2\tau - \Delta(\tau))]d\tau + O\left(\frac{1}{\lambda}\right) = 0.$$
(33)

Defining

$$A(x,\lambda,\Delta(\tau)) = \frac{1}{2} \int_0^x q(\tau) \sin\lambda \,\Delta(\tau) d\tau,$$
  

$$B(x,\lambda,\Delta(\tau)) = \frac{1}{2} \int_0^x q(\tau) \cos\lambda \,\Delta(\tau) d\tau,$$
(34)

and the Lemma III.3.3 in (Norkin, 1972),

$$\int_{0}^{x} q(\tau) \cos \lambda \left( 2\tau - \Delta(\tau) \right) d\tau = O\left(\frac{1}{\lambda}\right),$$
  
$$\int_{0}^{x} q(\tau) \sin \lambda \left( 2\tau - \Delta(\tau) \right) d\tau = O\left(\frac{1}{\lambda}\right).$$
 (35)

Using (34) and (35), equation (33) can be written as

 $\lambda\alpha\delta\cos\lambda\pi - \delta\sin\lambda\pi +$ 

$$\alpha\delta\sin\lambda\pi B(\pi,\lambda,\Delta(\tau)) - \alpha\delta\cos\lambda\pi A(\pi,\lambda,\Delta(\tau)) + O\left(\frac{1}{\lambda}\right) = 0.$$

Hence

$$\cot \lambda \pi = \frac{1 - \alpha B(\pi, \lambda, \Delta(\tau))}{\alpha \lambda} + O\left(\frac{1}{\lambda^2}\right).$$

Writing  $\lambda = \lambda_n = n + \frac{1}{2} + \delta_n$  into above equation

$$\delta_n = \frac{\alpha B(\pi, n, \Delta(\tau)) - 1}{\alpha n \pi} + O\left(\frac{1}{n^2}\right)$$

and

$$\lambda_n = n + \frac{1}{2} + \frac{\alpha B(\pi, n, \Delta(\tau)) - 1}{\alpha n \pi} + O\left(\frac{1}{n^2}\right)$$
are obtained.

**Theorem 5:** Under the same hypothesis of Theorem 4 eigenfunctions of the boundary value problem (1)-(5) are as follows:

$$u_{1n}(x) = \alpha \left( 1 - \frac{A(x, n, \Delta(\tau))}{n} \right) \cos\left(n + \frac{1}{2}\right) x$$
$$- \frac{(\alpha B(\pi, n, \Delta(\tau)) - 1)x}{n\pi} \sin\left(n + \frac{1}{2}\right) x$$
$$+ \left(\frac{\alpha B(x, n, \Delta(\tau) - 1)}{n}\right) \sin\left(n + \frac{1}{2}\right) x + O\left(\frac{1}{n^2}\right), \tag{36}$$

$$u_{2n}(x) = \frac{\alpha\delta}{n} \left( 1 - \frac{A(x, n, \Delta(\tau))}{n} \right) \cos\left(n + \frac{1}{2}\right) x$$
$$- \frac{\delta(\alpha B(\pi, n, \Delta(\tau)) - 1)x}{n^2\pi} \sin\left(n + \frac{1}{2}\right) x$$
$$+ \frac{\delta(\alpha B(x, n, \Delta(\tau)) - 1)}{n^2} \sin\left(n + \frac{1}{2}\right) x + O\left(\frac{1}{n^3}\right). \tag{37}$$

Proof: From (10) and (31) we obtain

$$\begin{split} \omega_1(x,\lambda) &= \alpha cos\lambda x - \frac{sin\lambda x}{\lambda} + \\ &+ \frac{\alpha}{\lambda} \int_0^x q(\tau) sin\,\lambda(x-\tau)\cos\lambda(\tau-\Delta(\tau))\,d\tau + \\ &+ O\left(\frac{1}{\lambda^2}\right). \end{split}$$

Using (34) and (35),

$$\omega_{1}(x,\lambda) = \alpha \left(1 - \frac{A(x,\lambda,\Delta(\tau))}{\lambda}\right) \cos\lambda x + \left(\frac{\alpha B(x,\lambda,\Delta(\tau)) - 1}{\lambda}\right) \sin\lambda x + O\left(\frac{1}{\lambda^{2}}\right).$$
(38)

Replacing  $\lambda$  with  $\lambda_n$  and using (30),

$$u_{1n}(x) = \alpha \left( 1 - \frac{A(x, n, \Delta(\tau))}{n} \right) \cos\left(n + \frac{1}{2}\right) x$$
$$- \frac{(\alpha B(\pi, n, \Delta(\tau)) - 1)x}{n\pi} \sin\left(n + \frac{1}{2}\right) x$$
$$+ \left(\frac{\alpha B(x, n, \Delta(\tau) - 1)}{n} \right) \sin\left(n + \frac{1}{2}\right) x + O\left(\frac{1}{n^2}\right).$$





Writing (10), (16), (31) and (32) into the equation (11), we have

$$\omega_{2}(x,\lambda) = \frac{\alpha\delta}{\lambda} \cos\lambda x - \frac{\sin\lambda x}{\lambda^{2}} + \frac{\alpha\delta}{\lambda^{2}} \int_{0}^{x} q(\tau) \sin\lambda(x-\tau) \cos\lambda(\tau) - \Delta(\tau) d\tau + + O\left(\frac{1}{\lambda^{3}}\right).$$

Using (34) and (35),

$$\omega_{2}(x,\lambda) = \frac{\alpha\delta}{\lambda} \left( 1 - \frac{A(x,\lambda,\Delta(\tau))}{\lambda} \right) \cos\lambda x + \delta\left(\frac{\alpha B(x,\lambda,\Delta(\tau)) - 1}{\lambda^{2}}\right) \sin\lambda x + O\left(\frac{1}{\lambda^{3}}\right).$$
(39)

Replacing  $\lambda$  with  $\lambda_n$  and using (30),

$$u_{2n}(x) = \frac{\alpha\delta}{n} \left( 1 - \frac{A(x, n, \Delta(\tau))}{n} \right) \cos\left(n + \frac{1}{2}\right) x$$
$$- \frac{\delta(\alpha B(\pi, n, \Delta(\tau)) - 1)x}{n^2 \pi} \sin\left(n + \frac{1}{2}\right) x$$
$$+ \frac{\delta(\alpha B(x, n, \Delta(\tau)) - 1)}{n^2} \sin\left(n + \frac{1}{2}\right) x + O\left(\frac{1}{n^3}\right)$$

is obtained.

### CONCLUSION

Eigenvalues and eigenfunctions of the (1)-(5) boundary value problem have the following asymptotic representation:

$$\lambda_n = n + \frac{1}{2} + \frac{\alpha B(\pi, n, \Delta(\tau)) - 1}{\alpha n \pi} + O\left(\frac{1}{n^2}\right)$$
$$u_n(x) = \begin{cases} u_{1n}(x), & x \in [0, \frac{\pi}{2}] \\ u_{2n}(x), & x \in \left(\frac{\pi}{2}, \pi\right] \end{cases}$$

here  $u_{1n}(x)$  and  $u_{2n}(x)$  are given by the equation (36) and (37) respectively.

## **CONFLICT OF INTEREST**

The authors have no conflicts of interest to declare.

# **RESEARCH AND PUBLICATION ETHICS STATEMENT**

The authors declare that this study complies with research and publication ethics.

### REFERENCES

- Aydin Akgun, F., Bayramov, A. and Bayramoğlu, M. (2013). Discontinuous boundary value problems with retarded argument and eigenparameter-dependent boundary conditions. Mediterranean journal of mathematics, 10(1), 277-288.
- Bayramoğlu, M., Köklü, K. Ö. and Baykal, O. (2002). On the spectral properties of the regular Sturm-Liouville Problem with the lag argument for which its boundary conditions depends on the spectral parameter. Turkish Journal of Mathematics, 26(4), 421-432.
- Cetinkaya, F. A. and Mamedov, K. R. (2017). A boundary value problem with retarded argument and discontinuous coefficient in the differential equation. Azerbaijan Journal of Mathematics, 7(1), 135-145.
- Hira, F. (2017). A trace formula for the Sturm-Liouville type equation with retarded argument. Commun. Fac. Sci. Univ. Ank. Sér. A1 Math. Stat, 66(1), 124-132.
- Kolmanovskii, V. and Myshkis, A. (1999). Introduction to the Theory and Applications of Functional Differential Equations. Dordrecht, The Nederlands: Kluwer Academic Publishers.
- Koparan, K. (2019). Sınırda parametre içeren geç kalan argümanlı Sturm Liouville probleminin özdeğerlerinin ve özfonksiyonlarının asimtotik özellikleri, Yüksek Lisans Tezi.
- Norkin, S. B. (1956). Boundary problem for a secondorder differential equation with a retarded argument. Uchenye Zapiski Moskovskogo Gosudarstvennogo Universiteta, 181, 59-72.
- Norkin, S. B. (1958). On periodic solutions of a linear homogeneous differential equation of second order with retarded argument. Matematicheskii Sbornik, 87(1), 71-104.
- Norkin, S. B. (1972). Differential Equations of the Second Order with Retarded Argument, Providence, Rhode Island: American Mathematical Society.
- Şen, E. and Bayramov, A. (2011). On calculation of eigenvalues and eigenfunctions of a Sturm-Liouville type problem with retarded argument which contains a spectral parameter in the boundary condition. Journal of Inequalities and Applications, 2011(1), 1-9.
- Şen, E. and Bayramov, A. (2011). Calculation of eigenvalues and eigenfunctions of a discontinuous boundary value problem with retarded argument which contains a spectral parameter in the boundary



condition. Mathematical and Computer Modelling, 54(11-12), 3090-3097.

- Şen, E. and Bayramov, A. (2013). Asymptotic formulations of the eigenvalues and eigenfunctions for a boundary value problem. Mathematical Methods in the Applied Sciences, 36(12), 1512-1519.
- Şen, E., Seo, J. J. and Araci, S. (2013). Asymptotic behaviour of eigenvalues and eigenfunctions of a Sturm-Liouville problem with retarded argument. Journal of Applied Mathematics, 2013.
- Yang, C. F. (2012). Trace and inverse problem of a discontinuous Sturm–Liouville operator with retarded argument. Journal of Mathematical Analysis and Applications, 395(1), 30-41.