PAPER DETAILS

TITLE: Chemical Composition and Insecticidal Activity of Origanum syriacum L. Essential Oil

Against Sitophilus oryzae and Rhyzopertha dominica

AUTHORS: Tunay Karan, Seyda SIMSEK, Ilyas Yildiz, Ramazan Erenler

PAGES: 87-93

ORIGINAL PDF URL: https://dergipark.org.tr/tr/download/article-file/438645



DOI: 10.21448/ijsm.404114

Published at http://www.ijate.net

http://dergipark.gov.tr/ijsm

Research Article

Chemical Composition and Insecticidal Activity of Origanum syriacum L. Essential Oil Against Sitophilus oryzae and Rhyzopertha dominica

Tunay Karan¹⁰, Seyda Simsek¹⁰, Ilyas Yildiz¹⁰, Ramazan Erenler^{10*1}

¹ Plant Research Laboratory, Department of Chemistry, Faculty of Arts and Sciences, Gaziosmanpasa University, Tokat, Turkey

² Department of Plant Protection, Faculty of Agriculture, Bozok University, Yozgat, Turkey

Abstract: *Origanum* genus belonging to the Lamiaceae family is aromatic and medicinal plant. It has been used in many countries for medicinal and pharmaceutical purpose. Aerial part of the *Origanum syriacum* L. was dried at shade. The essential oil was generated by steam distillation and compounds were identified by GC-MS analysis. γ -terpinene (26.7%), thymol (26.6%) and carvacrol (22.9%) was detected as the mail constituents. The essential oil was tested for insecticidal activity against adult of *Sitophilus oryzae* (L.) (Coleoptera: Curculionidae) and *Rhyzopertha dominica* (F.) (Coleoptera: Bostrychidae) using fumigation method. Essential oil revealed the excellent fumigant effect on *R. dominica* adults with a median lethal concentration (LC₅₀) value of 0.124 µL/L and 0.107 µL/L for 48 h and 72 h respectively. LC₅₀ values of *S. oryzae* were found as 0.173 µL/L and 0.135 µL/L for 48 h and 72 h respectively. As a result, essential oil of *O. syriacum* has a potency to be a natural insecticide.

ARTICLE HISTORY

Received: 12 January 2018 Revised: 04 March 2018 Accepted: 11 March 2018

KEYWORDS

Origanum syriacum, Insecticidal activity, Fumigant toxicity, Essential oil

1. Introduction

Plants include numerous fascinating compounds revealing plenty of biological activities [1-4]. Therefore, they play an important role in drug discovery and development process. *Origanum* genus belonging to Lamiaceae family, mainly distributed through the Mediterranean region and the Balkans has been represented almost 50 species through the world, fourteen of which are endemic for Turkey [5]. *Origanum* plant consumed as herbal tea and has been used for traditional medicine since ancient times as stimulating, antirheumatic, antispasmodic, and has antibacterial effects to treat various illness such as revulsion, dyspepsia, muscle contraction, diarrhea and infection diseases [6]. The secondary metabolites isolated from *O. rotundifolium* and *O. majorana* displayed the significant antiproliferative and antioxidant activities [7, 8]. *Origanum* species are well known for their essential oils which employ in flavoring agent mainly for meat, fish, soup and bottled food [9]. Carvacrol, caryophyllene, terpinene and thymol are the dominant essential oil compounds of *Origanum* revealing the significant biological activities [10]. *Origanum* demand has been increasing steadily in the world market, on accounts of the significance in food, pharmaceutical and cosmetic industries [11].

CONTACT: Ramazan Erenler rerenler@gmail.com Gaziosmanpasa University, Department of Chemistry, Faculty of Arts and Sciences, Plant Research Laboratory, Tokat, Turkey

Insect pests have threatened the crop production all over the world. Synthetic insecticides such as methyl bromide and phosphine fumigants have been used extensively to destroy stored pets and to protect stored food, feedstuffs, and agricultural materials [12]. However, widespread usage of fumigants has brought about the development of resistance in stored-product insects [13]. Moreover, owing to the depleting of ozone layer, residue forming, and carcinogenicity, some synthetic insecticides have been prohibited [14, 15]. As a result, it is essential to find novel fumigants which should be harmless to the environment, ecosystem, and non-target organisms to combat stored product insects. Natural products, especially essential oils have gained the great interest to replace synthetic products in fighting pets recently [16]. The rice weevil, *Sitophilus oryzae* and bean weevil, *Rhyzopertha dominica* are the most widespread and destructive insect pests of stored grains as well as stored products [17, 18].

Essential oils (EOs) exhibiting a broad spectrum of biological activities are generated from various aromatic plants. EOs have been extensively used in food and pharmaceutical industry [19]. Moreover, due to the their insecticidal effect, they have been employed in agricultural purpose [20].

Herein, the essential oil of *O. syriacum* was generated by steam distillation and identified by GC-MS. In addition, insecticidal activity test of essential oil was carried out against *S. oryzae* and *R. dominica*. The results revealed that the essential oil exhibited excellent insecticidal activity against these pets.

2. Material and Methods

2.1. Plant Material

The aerial parts of cultivated *Origanum syriacum* were harvested from the Medicinal and Aromatic Plant Field of Gaziosmanpasa University, Tokat, Turkey and identified by Prof. Dr. Ozgur Eminagaoglu. A voucher specimen was deposited in the Herbarium of the Faculty of Forest Engineering (ARTH: 5256) Artvin Coruh University.

2.2. Isolation of the Essential Oils

Aerial parts of plant material (50 g) were diluted with distillated water (250 ml) then subjected to hydrodistillation for 4 h, using a Clevenger apparatus. The oil was dried over sodium sulphate and stored in the freeze (4 °C) until to be analyzed.

2.3. GC and GC–MS Analysis

GC analyses were performed on a Perkin–Elmer Clarus 500 Series, in divided mode, 50:1, equipped with a flame ionization detector (FID) (Perkin-Elmer Clarus 500) and a mass spectrometer-equipped (Perkin-Elmer Clarus 500) BPX-5 capillary column (30 m × 0.25 mm, 0.25 um i.d., SGE Analytical science- Trajan scientific and medicinal). The temperature of injection was steady and FID was performed at 250 °C. Helium was the carrier gas at a rate of 1.0 ml/min. The oven temperature was 50 °C at the beginning then was raised to 220 °C with a rate of 8 °C/min. In the mass spectrometer, ionization energy was 70 eV and the temperature of transfer line was at 250 °C. The standard components were used for the majority of the essential oil constituents and Kovats retention indices (RIs) were determined for all the sample components using the Van den Dool and Kratz equation according to homolog n-alkane series retention times.

2.4. Insect

The cultures of *S. oryzae* and *R. dominica* were supplied from Department of Plant Protection, Bozok University. One third of 5 L glasswares were filled with clean wheat, and then adult males and females were added to lay eggs to get a single aged population. After the adults were removed, cultures were incubated at 27 ± 2 °C in a dark climate chamber with $60 \pm$

5% r.h. (relative humidity) for 48 h. The new generation of adults came into view by 45 day and 3-4 week-old adults were used in tests [21].

2.5. Dose-Response Bio-Assay

Insecticidal activity of essential oil was determined by fumigation method. Conducted dose effect on *S. oryzae* and *R. dominica* of *O. syriacum* was carried out. In the experiments, 5, 10, 15 and 20% (v/v) (essential oil/acetone) concentrations of the essential oil were used. Compressed rubber-capped glass tubes (20 ml) were used. Small paper discs were cut and then they are fixed to the cover and 10 adult specimens were situated in the each glass. 10 μ l volume of the oil mixture was saturated to the filter papers by pipettes. After this process, it was waited for 5 minutes to volatilize the acetone. Then the caps were covered on the glasses and they were waited in the incubator at 25°C in dark conditions for 24- 48 and 72 hours. The dead specimens were counted and recorded for each 24 hours interval. The randomized block design was used for the study. All experiments were repeated 3 times and each repetition consists of 3 replications [22].

2.6. Statistical Analysis

Dose-mortality response data from the bioassays were tested by POLO-PC probit analysis to determine the LC_{50} , LC_{90} , confidence level. The differences of LC_{50} , LC_{90} were executed by comparison of confidence levels [23].

3. Results and Discussion

Origanum species are well known with their rich essential oils contents which are clear red color, a pleasant and slightly spicy odor. In this study, EO analysis was executed by GC-MS and 14 components were identified with a rate of higher than 0.1%, representing 99.8% of the oil (Table 1). γ -Terpinene, thymol were the main compounds, at 26.7% and 26.6% respectively; carvacrol, p-cymene, monoterpenes which are the second most abundant compounds, were present at 22.9% and 13.3% respectively.

No	Compounds	RI	Percent (%)	
1	α -Thujene	935	0.71	
2	α -Pinene	941	1.07	
3	Camphene	955	0.13	
4	Myrcene	991	2.69	
5	α -Terpinen	1021	4.27	
6	<i>p</i> -Cymene	1027	13.32	
7	y-Terpinene	1067	26.66	
8	α -Terpinolen	1092	0.15	
9	Borneol	1169	0.12	
10	Terpinen-4-ol	1181	0.44	
11	α -Terpineol	1192	0.12	
12	Thymol	1291	26.62	
13	Carvacrol	1305	22.91	
14	β -Caryophyllene	1418	0.58	
	Total		99.79	

Table 1. Chemical composition of O. syriacum essential oil analyzed by GC-MS

RI: Retention indices calculated against n-alkanes, % calculated from FID data

In this work, fumigation assay was executed to investigate the toxicity of *O. syriacum* essential oil against adults of *S. oryzae* and *R. dominica* at four different concentrations. EO revealed the most fumigant effect on *R. dominica* adults with a median lethal concentration

(LC₅₀) value of 0.124 μ L/L at the confidence level of 0.100-0.140 for 48 h. Fumigation effect of EO increased with the time-dependence. LC₅₀ value of *R. dominica* was 0.107 μ L/L which was the best activity among the trials for 72 h. However, LC₅₀ values of *S. oryzae* adults were calculated as 0.173 μ L/L and 0.135 μ L/L for 48 h and 72 h respectively. While comparing LC₉₀ for *R. dominica* adults and *S. oryzae* adults, significant differences were detected. LC₉₀ values of *R. dominica* adults and *S. oryzae* adults were calculated as 0.198 μ L/L and 0.272 μ L/L for 48 h respectively. Moreover, these values (LC₉₀) were found as 0.192 μ L/L and 0.222 μ L/L for *R. dominica and S. oryzae* for 72 h respectively (Table 2).

Insects	hours	Slope±SD	$LC_{50} (\mu L/L)$	LC ₉₀ (µL/L)
			(Confidence level)	(Confidence level)
R. dominica	48	17.166±2.790	0.124	0.198
			(0.100 - 0.140)	(0.181-0.226)
	72	15.114±2.233	0.107	0.192
			(0.082-0.125)	(0.174-0.218)
S. oryzae	48	13.014 ± 1.520	0.173	0.272
			(0.153-0.206)	(0.231-0.360)
	72	14.713 ± 1.508	0.135	0.222
			(0.123-0.147)	(0.202-0.251)

Table 2. Fumigant effect of O. syriacum essential oil on R. dominica and S. oryzae

These results revealed a few differences regarding to the compounds and quantities with the literatures. For instance, *O. syriacum* collected from August at altitudes of 250-1900 m consisted of carvacrol and *p*- cymene at 60.5% and 7.4% respectively. In the same work, it was reported that *O. onites* essential oil constituted carvacrol (69.0%) as a chief product. Furthermore, carvacrol (%39.1), thymol (22.2%), and *p*-cymene (9.3%) were found in EO of *O. vulgare* [24]. These results emphasized the important variability of chemical composition of *Origanum* oils through their components. These differences could be on account of various criterions such as part of the plant, harvest period, location, altitude, climate, environmental conditions. Indeed, the synthesis and secretion of oils are influenced by these factors [25].

Insecticidal activity of *Origanum* species EOs was executed against various insects that EOs displayed the considerable activity. For instance, *O. majorana* EO displayed the good fumigant activity against *Spodoptera littoralis* which was the serious pest of some crops such as cotton, chili, and tobacco [26]. The major compounds of *O. majorana* EO were detected as thymol (38.8%), carvacrol (32.9%), *p*-cymene (7.9%), and *γ*-terpinene (5.1%). The chemical compounds of *O. majorana* have been similar with *O. syriacum*. Therefore, the insecticidal activity of *O. syriacum* may be due to the major constituents of essential oil components. Another work carried out in Algeria revealed that *O.glandulosum* has highly toxic potential power against *Rhyzopertha dominica* by both contact and fumigant assays [27].

Insecticidal components of most essential oils are usually monoterpene [28]. Because of the high volatility, essential oils have fumigant and gaseous action that may be significance for stored product insecticides. The carvracrol, a monoterpene has extensive insecticidal and acaricidal activity against agricultural stored product, and medicinal pests and act as a fumigant [29].

A good natural insecticide has been accepted that it influences a limited range of pest insects, has no harmful effects on non-target organisms and the environment, act in many ways on various types of pest complex [30]. Most plant extract and essential oils have been known to possess repellent and insecticidal activity against various stored product insects [31]. Some essential oils such as mustard oil, cinnamon oil and horseradish oil were reported to have strong

effect against *S. oryzae* insect. In addition, essential oils generated from *Pinus longifolia*, *Eucalyptus obliqua* and *Coriandrum sativum* were tested for contact and fumigant activities against rice weevil, *Sitophilus oryzae*, *Callosobruchus chinensis* and *Corcyra cephalonica* and the results varied with the test material, insect species and exposure time. In fumigation assay, *Corcyra cephalonica* and *Eucalyptus obliqua* oils at 130 µg/cm², led to 100% toxicity to all the species within the 24 h of the treatment [32].

4. Conclusion

O. syriacum essential oil has a potential for applications for stored-grain pests due to its high volatility and fumigant activity. In addition, EO of *O. syriacum* might be applicable for managing *S. oryzae* and *R. dominica* insects in enclosed spaces such as storage bins, glasshouses, or buildings due to their fumigant action. These results suggested that the insecticidal action of the oils might be due to the fumigant action. In other word, the oils could be toxic by penetrating the insect body via the respiratory system. The significant insecticidal activity of *O. syriacum* could be caused from γ -terpinene, thymol and carvacrol which were the major compounds of EO content. The major three compounds have methyl and isopropyl moiety. Thymol and carvacrol are constitutional isomers. In other words, both compounds have same molecular formula, C₁₀H₁₄O but different structural formulas. Methyl, isopropyl and hydroxyl connected to the phenyl group in thymol and carvacrol. Hence, methyl, isopropyl and hydroxyl bonded to the phenyl may be effective for the activity.

Conflict of interest

The authors declare that they have no conflict of interest.

Orcid

Tunay Karan ^(b) https://orcid.org/0000-0002-9114-8400 Seyda Simsek ^(b) https://orcid.org/0000-0002-0096-8949 Ilyas Yıldız ^(b) https://orcid.org/0000-0003-1254-1069 Ramazan Erenler ^(b) https://orcid.org/0000-0002-0505-3190

5. References

- [1] Elmastas, M., Erenler, R., Isnac, B., Aksit, H., Sen, O., Genc, N., & Demirtas, I. (2016). Isolation and identification of a new neo-clerodane diterpenoid from *Teucrium chamaedrys* L. *Natural Product Research*, 30 (3): 299-304.
- [2] Erenler, R., Yilmaz, S., Aksit, H., Sen, O., Genc, N., Elmastas, M., & Demirtas, I. (2014). Antioxidant Activities of Chemical Constituents Isolated from Echinops orientalis Trauv. *Records of Natural Products*, 8 (1): 32-36.
- [3] Erenler, R., Sen, O., Yildiz, I., & Aydin, A. (2016). Antiproliferative activities of chemical constituents isolated from *Thymus praecox* subsp *grossheimii* (Ronniger) Jalas. *Records of Natural Products*, 10 (6), 766-770.
- [4] Demirtas, I., Erenler, R., Elmastas, M., & Goktasoglu, A. (2013). Studies on the antioxidant potential of flavones of *Allium vineale* isolated from its water-soluble fraction. *Food Chemistry*, 136 (1), 34-40.
- [5] Davis, P.H. (1998). Flora of Turkey and the East Aegean Islands. Edinburg, United Kingdom: University Press.
- [6] Jun, W.J., Han, B.K., Yu, K.W., Kim, M.S., Chang, I.S., Kim, H.Y., & Cho, H.Y. (2001). Antioxidant effects of *Origanum majorana* L. on superoxide anion radicals. *Food Chemistry*, 75 (4): 439-444.

- [7] Erenler, R., Sen, O., Aksit, H., Demirtas, I., Yaglioglu, A.S., Elmastas, M., & Telci, I. (2016). Isolation and identification of chemical constituents from *Origanum majorana* and investigation of antiproliferative and antioxidant activities. *Journal of the Science of Food and Agriculture*, 96 (3): 822-836.
- [8] Erenler, R., Meral, B., Sen, O., Elmastas, M., Aydin, A., Eminagaoglu, O., & Topcu, G. (2017). Bioassay-guided isolation, identification of compounds from *Origanum rotundifolium* and investigation of their antiproliferative and antioxidant activities. *Pharmaceutical Biology*, 55 (1): 1646-1653.
- [9] Busatta, C., Vidal, R.S., Popiolski, A.S., Mossi, A.J., Dariva, C., Rodrigues, M.R.A., Corazza, F.C., Corazza, M.L., Oliveira, J.V., & Cansian, R.L. (2008). Application of *Origanum majorana* L. essential oil as an antimicrobial agent in sausage. *Food Microbiology*, 25 (1): 207-211.
- [10] Economou, G., Panagopoulos, G., Tarantilis, P., Kalivas, D., Kotoulas, V., Travlos, I.S., Polysiou, M., & Karamanos, A. (2011). Variability in essential oil content and composition of *Origanum hirtum* L., *Origanum onites* L., *Coridothymus capitatus* (L.) and *Satureja thymbra* L. populations from the Greek island Ikaria. *Industrial Crops and Products*, 33 (1): 236-241.
- [11] Milos, M., Mastelic, J., & Jerkovic, I. (2000). Chemical composition and antioxidant effect of glycosidically bound volatile compounds from oregano (*Origanum vulgare* L. ssp *hirtum*). *Food Chemistry*, 71 (1), 79-83.
- [12] Athanassiou, C.G., Hasan, M.M., Phillips, T.W., Aikins, M.J., & Throne, J.E. (2015). Efficacy of methyl bromide for control of different life stages of stored-product psocids. *Journal of economic entomology*, 108 (3), 1422-1428.
- [13] Abouseadaa, H.H., Osman, G.H., Ramadan, A.M., Hassanein, S.E., Abdelsattar, M.T., Morsy, Y.B., Alameldin, H.F., El-Ghareeb, D.K., Nour-Eldin, H.A., & Salem, R. (2015). Development of transgenic wheat (*Triticum aestivum* L.) expressing avidin gene conferring resistance to stored product insects. *BMC plant biology*, 15 (1): 183.
- [14] Rajashekar, Y., Ravindra, K., & Bakthavatsalam, N. (2014). Leaves of *Lantana camara* Linn. (Verbenaceae) as a potential insecticide for the management of three species of stored grain insect pests. *Journal of food science and technology*, 51 (11), 3494-3499.
- [15] Jagadeesan, R., Nayak, M.K., Pavic, H., Chandra, K., & Collins, P.J. (2015). Susceptibility to sulfuryl fluoride and lack of cross-resistance to phosphine in developmental stages of the red flour beetle, *Tribolium castaneum* (Coleoptera: Tenebrionidae). *Pest management science*, 71 (10), 1379-1386.
- [16] Isman, M.B. (2006). Botanical insecticides, deterrents, and repellents in modern agriculture and an increasingly regulated world. *Annu Rev Entomol*, 51, 45-66.
- [17] Aref, S.P., Valizadegan, O., & Farashiani, M.E. (2015). *Eucalyptus dundasii* Maiden essential oil, chemical composition and insecticidal values against *Rhyzopertha dominica* (F.) and *Oryzaephilus surinamensis* (L.). *Journal of Plant Protection Research*, 55 (1), 35-41.
- [18] Mishra, B.B., Tripathi, S., & Tripathi, C. (2013). Bioactivity of two plant derived essential oils against the rice weevils *Sitophilus oryzae* (L.) (Coleoptera: Curculionidae). *Proceedings of the National Academy of Sciences, India Section B: Biological Sciences*, 83 (2), 171-175.
- [19] Shaaban, H.A.E., El-Ghorab, A.H., & Shibamoto, T. (2012). Bioactivity of essential oils and their volatile aroma components: Review. *Journal of Essential Oil Research*, 24 (2), 203-212.

- [20] Huang, Y., Lam, S., & Ho, S. (2000). Bioactivities of essential oil from *Elletaria cardamomum* (L.) Maton. to *Sitophilus zeamais* Motschulsky and *Tribolium castaneum* (Herbst). *Journal of Stored Products Research*, 36 (2), 107-117.
- [21] Abay, G., Karakoc, O.C., Tufekci, A.R., Koldas, S., & Demirtas, I. (2012). Insecticidal activity of *Hypnum cupressiforme* (Bryophyta) against *Sitophilus granarius* (Coleoptera: Curculionidae). *Journal of Stored Products Research*, 51, 6-10.
- [22] Polatoğlu, K., Karakoç, Ö.C., Yücel, Y.Y., Gücel, S., Demirci, B., Demirci, F., & Başer, K.H.C. (2017). Insecticidal activity of *Salvia veneris* Hedge. Essential oil against coleopteran stored product insects and *Spodoptera exigua* (Lepidoptera). *Industrial Crops and Products*, 97: 93-100.
- [23] LeOra Software (1994). Polo-PC a user's guide to probit or logit analysis. 1119 Shattuck Avenue, Berkeley, CA, 94707: USA.
- [24] Schulz, H., Özkan, G., Baranska, M., Krüger, H., & Özcan, M. (2005). Characterisation of essential oil plants from Turkey by IR and Raman spectroscopy. *Vibrational Spectroscopy*, 39 (2), 249-256.
- [25] Khalfi, O., Sahraoui, N., Bentahar, F., & Boutekedjiret, C. (2008). Chemical composition and insecticidal properties of *Origanum glandulosum* (Desf.) essential oil from Algeria. *Journal of the Science of Food and Agriculture*, 88 (9), 1562-1566.
- [26] Souguir, S., Chaieb, I., Cheikh, Z.B., & Laarif, A. (2013). Insecticidal activities of essential oils from some cultivated aromatic plants against *Spodoptera littoralis* (Boisd). *Journal* of Plant Protection Research, 53 (4), 388-391.
- [27] Sahraoui, N., Hazzit, M., & Boutekedjiret, C. (2017). Effects of microwave heating on the antioxidant and insecticidal activities of essential oil of *Origanum glandulosum* Desf. obtained by microwave steam distillation. *Journal of Essential Oil Research*: 1-10.
- [28] Konstantopoulou, I., Vassilopoulou, L., Mavragani-Tsipidou, P., & Scouras, Z. (1992). Insecticidal effects of essential oils. A study of the effects of essential oils extracted from eleven Greek aromatic plants on *Drosophila auraria*. *Experientia*, 48 (6), 616-619.
- [29] Ahn, Y.-J., Lee, S.-B., Lee, H.-S., & Kim, G.-H. (1998). Insecticidal and acaricidal activity of carvacrol and β-thujaplicine derived from *Thujopsis dolabrata* var. *hondai* Sawdust. *Journal of Chemical Ecology*, 24 (1), 81-90.
- [30] Regnault-Roger, C. (1997). The potential of botanical essential oils for insect pest control. Integrated Pest Management Reviews, 2 (1), 25-34.
- [31] El-Nahal, A., Schmidt, G., & Risha, E. (1989). Vapours of Acorus calamus oil—a space treatment for stored-product insects. *Journal of Stored Products Research*, 25 (4), 211-216.
- [32] Rani, P.U. (2012). Fumigant and contact toxic potential of essential oils from plant extracts against stored product pests. *Journal of Biopesticides*, 5 (2), 120.