PAPER DETAILS

TITLE: Microwave-assisted synthesis of N-benzylidene-4-fluoroaniline and

N-benzylidene-4-Nitroaniline and their inhibitory activities on hCA isoenzymes

AUTHORS: Hülya ÇELIK, Aynur BABAGIL

PAGES: 38-48

ORIGINAL PDF URL: https://dergipark.org.tr/tr/download/article-file/633078



2019, Vol. 6, No. 1, 38-48

https://dx.doi.org/10.21448/ijsm.479108

Published at http://www.ijate.net

http://dergipark.gov.tr/ijsm

Research Article

Microwave-assisted synthesis of N-benzylidene-4-fluoroaniline and N-benzylidene-4-Nitroaniline and their inhibitory activities on hCA isoenzymes

Hülya Celik ^{(1)*,1}, Aynur Babagil ²

¹Agri Ibrahim Cecen University, Faculty of Pharmacy, 04100, Agri, Turkey

² Ataturk University, Faculty of Engineering, Department of Nano-Science and Nano-Engineering, 25070, Erzurum, Turkey

Abstract: In this study, N-benzylidene-4-fluoroaniline (3a), N-benzylidene-4nitroaniline (3b) as a result of condensation of benzaldehyde (1) and 4fluoroaniline (2a), 4-nitroaniline (2b) using a microwave method was synthesized. The structures of the synthesized compounds were characterized by Fouirer Transform Infrared Spektrofotometre (FTIR spectroscopy), Nuclear Magnetic Resonance (NMR spectroscopy) and elemental analysis methods. The prepared compounds were tested for their inhibitory effects on carbonic anhydrase isoenzymes (hCA-I and hCA-II). The results showed that the synthesized compounds 3a and 3b had a strong inhibitory effect on hCA-I and hCA-II enzymes activity in vitro. By testing these compounds in other CA isoenzymes, more effective CA inhibitors can be formed. Thus, new therapeutic applications can be made for enzyme activators in the near future.

ARTICLE HISTORY

Received: November 05, 2018 Revised: January 14, 2019 Accepted: January 20, 2019

KEYWORDS

Schiff, Microwave, Fluoroaniline, Nitroaniline, Inhibitor

1. INTRODUCTION

Schiff bases derived from the reaction of primary amines with carbonyl compounds are used in a wide range of different fields. [1]. Schiff bases are indicated by the general formula RCH = NR. R and R 'are alkyl or aryl substituents [2]. Schiff bases are widely used in various fields such as pharmaceutical preparation, agricultural and plastic industry, polymer production, electronics industry, cosmetics, dyestuff production due to their biological and structural properties [3]. Schiff bases have antifungal properties as fungicides, insecticides, chemotherap eutic, anticancer agents in antioxidants, antimutagenic and antiviral drugs [4-8]. Nitro and halo derivatives of Schiff bases are known to have antimicrobial and anti-tumoral activity.

For the first time in 1986 Gedye used microwave in organic synthesis. The synthesis of microwave methods in scientific studies has started to attract great interest in recent years [9]. The microwave is a low frequency electromagnetic energy state at the end of the electromagnetic spectrum between the radio and infrared waves in the frequency range 300-300.000 MHz [10]. Microwave chemistry continues its development with scientific studies [11-

ISSN-e: 2148-6905 /© IJSM 2019

^{*}CONTACT: Hülya Çelik 🖾 hycelik@agri.edu.tr 🖃 Agri Ibrahim Cecen University, Faculty of Pharmacy, 04100, Agri, Turkey

Celik & Babagil

15]. Because of the direct heating of the substances in the microwave reactions, the products are obtained with pure and high yields in less time than the classical heating methods [16, 17].

Carbonic anhydrase enzymes (CA, EC 4.2.1.1) are metalloproteins containing zinc (Zn^{2+}) which catalyze the conversion of CO₂ to bicarbonate (HCO₃⁻) and proton (H⁺) release. In addition, the carbonic anhydrase enzyme plays a role in many important reactions such as transport of CO₂ in tissues, pH and CO₂ homeostasis, transport of ions, biosynthetic reactions, bone formation. CA enzymes have been found in many tissues such as salivary glands, muscles, brain, nerve myelin sheath, pancreas, prostate and endometrial tissues by histochemical methods [18].

$$CO_2 + H_2O \longrightarrow H_2CO_3 \implies H^+ + HCO_3^-$$

The carbonic anhydrase isoenzymes are encoded by five different family of genes that are evolutionally independent, including α -, β -, CA-, δ - and ϵ - CA. Six teen different CA isoenzymes and CA-linked proteins (CARPs) were identified in the mammals. hCA-I and hCA-II are two main cytosolic CA isoforms found in mammalian red blood cells [19, 20].

For this reason, in this study it is aimed to synthesize Schiff base derivative N-benzylidene-4-fluoroaniline (3a), N-benzylidene-4-nitroaniline (3b) by micro wave method and investigate the inhibition potentials on CA-I and hCA-II isoenzymes.

2. MATERIALS and METHODS

2.1. Chemicals

Benzaldehyde, 4-fluoroaniline are commercially available (Merck, Sigma-Aldrich) and were used without further purification. Sepharose-4B, sulphanylamide, L-tyrosine, Tris, Na₂SO₄, protein assay reagents, and chemicals for electrophoresis were purchased from Sigma-Aldrich Co. All other chemicals were of analytical grade and obtained from Merck. Reactions were monitored via thin-layer chromatography (TLC). ¹H NMR and ¹³C NMR spectra were recorded on a 400 (100) MHz Varian spectrometer using CDCl₃. Column chromatography was performed on silica gel 60 (70–230 mesh ASTM), and TLC was carried out on silica gel (254–366 mesh ASTM). Melting points were determined on a capillary melting apparatus (Buchi 530) and are uncorrected. Infrared (IR) spectra were obtained from solutions in 0.1-mm cells with a Perkin-Elmer spectrophotometer (Waltham, MA). Elemental analyzes were performed on a Leco CHNS-932 apparatus.

2.2. General Method for Synthesis of 3a, 3b

4-floroanilin (2a) (1mmol), 4-nitroanilin (2b) (1mmol) was added into the benzaldehyde (1) (1 mmol) mixture, and then the reaction mixture was exposed to microwave radiation at 2450 MHz. The progress of the reaction was monitored by thin layer chromatography (TLC) (Runner phase, *n*-hexane-ethylacetate (4: 1) was used). It was determined that the reactions were completed in 8 minutes for all aniline derivatives. The crude products 3a ve 3b were obtained in pure form (Scheme 1).



Scheme 1. The synthesis of the compounds 3a, 3b

2.3. Purification of Carbonic Anhydrase Isozymes (hCA-I and hCA-II) from Human Erythrocytes by Affinity Chromatography

Fresh human blood was obtained from Atatürk University, Blood Center, kept at 4°C and used within 2-3 days at most. Blood samples were centrifuged for 15 minutes at 2500 rpm to separate the erythrocytes, and the remaining plasma and leucocytes layers were carefully removed and discarded. Subsequently, the remaining erythrocytes were washed twice with 0.9% NaCl solution and the upper portions were discarded. Then, the erythrocytes were hemolyzed with pure water at 0°C and mixed for half an hour, and the cell membranes were separated by centrifuging the hemolyzate at 20,000 rpm at 4°C for half an hour. The pH was adjusted to 8.7 with solid Tris. Thus, the hemolyzate came to be applied to the colonic [21].

The affinity was prepared on gel Sepharose-4B matrix. After activating Sepharose-4B with CNBr, L-tyrosine was covalently attached as the extension arm. Tyrosine was then coupled by diazotization of the sulfanilamide as the ligand. The hemolysate was applied to the prepared Sepharose-4B-L-tyrosine-sulfanylamide affinity column equilibrated with 25 mM Tris–HCl/0.1 M Na₂SO₄ (pH 8.7). The affinity gel was washed with 25 mM Tris–HCl/22 mM Na₂SO₄ (pH 8.7). The affinity gel was washed with 25 mM Tris–HCl/22 mM Na₂SO₄ (pH 8.7). The human carbonic anhydrase (hCA I and hCA II) isozymes were eluted with 1 M NaCl/25 mM Na₂HPO₄ (pH 6.3) and 0.1 M CH₃COONa/0.5 M NaClO₄ (pH 5.6), respectively. All procedures were stored at 4 $^{\circ}$ C [22].

2.4. Hydratase Activity Assay

Carbonic anhydrase activity was determined using the Wilbur-Anderson Method modified by Rickli et al. [23]. This method is based on the determination of the pH change from the H⁺ ion resulting from the hydration of CO₂ to the bromine thymol blue indicator and measuring the permeation rate. The time (tc) interval was determined between addition of CO₂ solution and occurrence of a yellow-green color. The same interval was recorded without enzyme solution (to). The activity was calculated from the formula. One Wilbur – Anderson Unit = (to- tc) / tc

2.5. Inhibition Assays

In order to calculate the IC_{50} values of hCA I and hCA-II enzymes, inhibitor effects of Schiff bases derivatives were run at different concentrations, keeping the substrate concentration constant on hydratase activity. The activities of the enzymes in the uninhibited medium were used as 100% activity. Hydratase activities were measured in the presence of different concentrations of enzymes, and % activity values were calculated. IC_{50} values were calculated for each inhibitor using the scheme of % Activity- [inhibitor] [24].

2.6. Protein Determination

The protein assay for all purification steps was performed according to the Bradford method [25]. This method is based on the principle that coomassie brilliant blue G-250 binds

to proteins in a phosphoric acid medium. The resulting complex shows maximum absorbance at 595 nm. The sensitivity of this method is between 1 - 100 micrograms.

3. RESULTS

N-benzylidene-4-fluoroaniline (3a), N-benzylidene-4-nitroroaniline (3b), were prepared from reaction of benzaldehyde (1), 4-floroaniline (2a), and 4-nitroaniline (2b) by microwave method (Scheme 2). It was determined that the reactions were completed in 8 minutes for two Schiff bases derivatives. Its physical properties, analytical and spectral data of the imine's compounds are summarized below.

Scheme 2. The synthesis route of the compound (3a,3b)

3.1. N-Benzylidene-4-fluoroaniline (3a)

Yield: 95 %, M.p. 104 °C, yellow solid.¹H NMR (400 MHz, CDCl₃) δ 8.45 H (s, 1H), 7.90 (dd, *J* = 7.0, 2.6 Hz, 2H), 7.48 (dd, *J* = 5.2, 2.0 Hz, 3H), 7.21 (dd, *J* = 8.6, 5.0 Hz, 2H), 7.08 (t, *J* = 8.5 Hz, 2H) (Figure 1).¹³C NMR (400 MHz, CDCl₃) δ 160.4, 131.7, 129.0, 129.0, 122.6, 122.5, 116.2, 115.9, 29.9 (Figure 2). FTIR (CDCl₃, cm⁻¹): 3408 (aromatic CH), 3058 (aromatic CH), 2923, 2852 (CH=N), 1627 (aromattic C=C), 1505, 1261, 1222, 1094 (C-F), 801, 700 cm⁻¹ (Figure 3). C₁₃H₁₀FN Anal. calc. for: C,78.37; H, 5.06; N, 7.03 % Found: C, 78.71; H, 4.870; N, 7.093 %

Figure 1. ¹H-NMR spectrum (400 MHz) of Compound **3a** (CDCl₃, δ ppm)

Figure 2. ¹³C-NMR Spectrum (400 MHz) of Compound 3a (CDCl₃, δ ppm)

Figure 3. FTIR spectrum of Compound 3a (CDCl₃, cm⁻¹)

3.2. N-Benzylidene-4-nitroaniline (3b)

Yield: 93 %, M.p. 115°C, yellow solid. ¹H NMR (400 MHz, CDCl₃) δ 8.43 (s, 1H), 8.27 (d, J = 8.8 Hz, 2H), 7.92 (d, J = 7.1 Hz, 2H), 7.61 – 7.45 (m, 3H), 7.25 (d, J = 8.9 Hz, 2H). (Figure 4).¹³C NMR (400 MHz, CDCl₃) δ 162.95, 135.37, 132.43, 129.30, 128.99, 126.35, 125.05, 121.27, 113.37, 77.35, 77.03, 76.72. (Figure 5). FTIR (CDCl₃, cm⁻¹): 3482 (aromatic CH), 3369 (CH=N), 1629 (aromatic C=C), 1586, 1470, 1304, 1114, 838, 752, 697, 488 cm⁻¹ (Figure 6). C₁₃H₁₀N₂O₂ Anal. calc. for: C,69.02; H, 4.46; N, 12.38; O, 14.14 % Found: C, 69.50; H, 4.370; N, 12.02; O, 15.10 %

Figure 4. ¹H-NMR spectrum (400 MHz) of Compound **3b** (CDCl₃, δ ppm)

Figure 5. ¹³C-NMR Spectrum (400 MHz) of Compound 3b (CDCl₃, δ ppm)

Figure 6. FTIR spectrum of Compound 3b (CDCl₃, cm⁻¹)

3.3. Purification Results of Carbonic Anhydrase Isoenzymes (hCA-I and hCA-II)

hCA-I and hCA-II isoenzymes were purified in one step using human blood Sepharose 4B-L tyrosine-sulphanylamide affinity chromatography. hCA-I was purified with specific activity of 2953.1 fold and 60.2 %, hCA-II purified with 797.33 specific activity, 7497.2 fold and 52.3% yield (Table 1).

Purification steps	Volume	Activity	Total a	ctivity	Protein	Specific activity	Purification layer number
	(mL)	(EU/mL)	(EU)	(%)	(mg/mL)	(EU/mg)	_ ,
Hemolysate	100	44.67	4467	100	1.64×10^{3}	0.27	-
hCA-I	45	59.8	2691	60.2	0.075	797.33	2953.1
hCA-II	35	66.8	2338	52.3	0.033	2024.24	7497.2

Table 1. Purification of hCA I and hCA II from human erythrocytes

The carbonic anhydrase enzyme has been purified many times from numerous different organisms and the effects of various pesticides, chemicals and drugs on enzyme activities have been investigated. In this study, hCA-I and hCA-II isoenzymes were purified by Sepharose 4B-L-tyrosine-sulphanylamide affinity chromatography and kinetic studies for schiff base (**3a**, **3b**) were carried out by hydratase activity method using CO₂ as substrate. For two isoenzymes this was seen as a highly effective activator. The inhibitory effects on the activity of the isoenzymes were tested in vitro, the IC₅₀ values were calculated from the % Activity- [inhibitor] plot drawn for compound (**3a**, **3b**) and the results are given in (Figure 7), (Figure 8).

Figure 7. IC₅₀ values for the in vitro inhibition of hCA I and hCA II with synthetized *N*-Benzylidene-4-fluoroaniline (3a)

Chemical	CA-I	CA-II	
	IC_{50}	IC_{50}	
$C_{13}H_{10}N_2O_2$	1,85	2,07	
Acetazolamide (AAZ)	5.2	6.1	

Figure 8. IC₅₀ values for the in vitro inhibition of hCA I and hCA II with synthetized N-Benzylidene-4-nitroaniline (**3b**)

4. DISCUSSION

In vitro effects of $C_{13}H_{10}FN$, $C_{13}H_{10}N_2O_2$ on human activities of hCA-I and hCA-II enzyme purified from human erythrocytes revealed that inhibitory activity against acetazolamide reference compound was higher. The most effective effect on hCA1 and hCAII isoenzymes has IC₅₀ values of 1.82 mM and 1.85 mM, respectively, of the $C_{13}H_{10}FN$, compound. It is thought that the newly synthesized Schiff bases show strong inhibition effect on hCAI and hCAII isoenzymes and they will serve as a reference for the formation of new inhibitors of CA enzyme [26-27].

The pharmacological effects of synthesized compounds **3a** and **3b** can be improved clinically for hCA I and II. The novel therapeutic applications of these enzyme inhibitors or activators will be directed to the design of prodrugs and drugs in the health sector.

Acknowledgements

The authors thanks Prof. Dr Hasan SEÇEN and Prof. Dr. Hayrunnisa NADAROĞLU for her kind help on the biological activity studies of the compound.

Orcid

Hülya Çelik 🔟 https://orcid.org/0000-0003-0805-0523

5. REFERENCES

- [1]. Pfeiffer, P., Breith, E., Lubbe, E., Tsumaki, T. (1933). Tricyclische Orthokondenzierte Nebenvolenzringe. *Annalen Der Chemie.*, 503, 84-127.
- [2]. Oskay, E., (1990). Organik Kimya, Hacettepe Üniversitesi Yayınları A-42, 243, Ankara.
- [3]. Raman, N., Muthuraj, V., Ravichandran, S., Kulandaisamy, A. (2003). Synthesis, characterisation and electrochemical behaviour of Cu(II), Co(II), Ni(II) and Zn(II) complexes derived from acetylacetone and p-anisidine and their antimicrobial activity. *Proc Indian Acad. Sci.*, 115-161.
- [4]. Helmut, S. (1976). *Metal ions in biological systems*, Marcel Dekker Inc, New York, 2-50s.
- [5]. Metzler, CM., Cahill, A., Metzler DE. (1980). Equilibriums and absorption spectra of Schiff bases. *Journal of The American Chemical Society*, *102* (19), 6075- 6082.
- [6]. Şekeroğlu, Z.A., Şekeroğlu, V. (2011). The in vitro alkaline comet assay in genetic toxicology. *The Journal of Applied Behavioral Science*, *5*, 49-54.
- [7]. Kaya, B. (2003). Anti-genotoxic effect of ascorbic acid on mutagenic dose of three alkylating agents. *Turkish Journal of Biology*, 27, 241-246.
- [8]. Braconi, D., Sotgi, M., Millucci, L., Paffetti, A., Tasso, F., Alisi, C., Martini, S, Rappuoli, R., Lusini, P., Rosa, A., Rossi, C., Santucci, A. (2006). Comparative analysis of the effects of locally used herbicides and their active ingredients on a wild-type wine (Saccharomyces cerevisiae) strain. *Journal of Agricural and Food Chemistry*, 54, 3163-3172.
- [9]. Kappe, C.O. (2002). High-Speed Combinatorial Synthesis Utilizing Microwave Irradiation. *Curr Opin Chem Biol.*, *6*, 314-320.
- [10]. Kappe, C.O., Dallinger, D., Murphree, S.S. (2009). Practical Microwave Synthesis For Organic Chemists: Strategies, Instruments, And Protocols, Wiley-Vch, 299.
- [11]. Başarır, M. (2006). *N-Sübstitüe Pirol Türevlerinin Mikrodalga Etkisi Altında Sentezi*. Yüksek Lisans Tezi, Yıldız Teknik Üniversitesi Fen Bilimleri Enstitüsü, İstanbul, 82 S.
- [12]. Serdar, G., Demir, E., Bayrak, S., Sökmen, M. (2016). New Approaches for Effective Microwave Assisted Extraction of Caffeine and Catechins from Green Tea. *International Journal of Secondary Metabolite*, 3 (1), 3-13 <u>https://dx.doi.org/10.21448/ijsm.240697.</u>
- [13]. Özturk, B, Yiğitarslan, S. (2017). Microwave Extraction of Gallic Acid Equivalent of Total Flavonoids from Nigella sativa. *International Journal of Secondary Metabolite*, 4 (3, Special Issue 1), 117-124. <u>https://dx.doi.org/10.21448/ijsm.363838</u>.

- [14]. Shysh, S., Mazets, Z., Shutava, H., Sysha, O., Gorgun, Y. (2018). Electromagnetic radiation of low intensity as a factor of change of phenolic compounds content. *International Journal of Secondary Metabolite*, 5(3), 252 – 258 <u>https://dx.doi.org/10.214</u> <u>48/ijsm.459102</u>.
- [15]. Serdar, G., Demir, E., Sökmen, M. (2017). Recycling of Tea Waste: Simple and Effective Separation of Caffeine and Catechins by Microwave Assisted Extraction (MAE). *International Journal of Secondary Metabolite*, 4(2), 78 - 89. <u>https://dx.doi.org/10.21448</u> /ijsm.288226
- [16]. Gedye, R., Smith, F., Westaway, K., Ali, H., Baldisera, L., Laberge, L., Roussel, J. (1986). The use of microwave ovens for rapid organic synthesis. *Tetrahedron Letters* 27(3), 279-282.
- [17]. Loferer, M. J., Tautermann, C.S., Loeffler, H.H., Liedl, K.R. (2003). Influence of backbone conformations of human carbonic anhydrase II on carbon dioxide hydration: hydration pathways and binding of bicarbonate. J. Am. Chem. Soc., 125, 8921–8927.
- [18]. Lindskog, S., Coleman, J.E., (1973). The Catalytic Mechanism of Carbonic Anhydrase. *Proc. Nat. Acad. Sci.*, 70, (9), 2505-2508.
- [19]. Zimmerman, S.A., Ferry, J.G., Supuran, C.T. (2007). Inhibition of the archaeal beta-class (Cab) and gamma-class (Cam) carbonic anhydrases. *Curr Top Med Chem.*, 9, 901-8.
- [20]. Demir, Y., Nadaroğlu, H., Demir, N. (2007). Purification and some properties of carbonic anhydrase from *Elephas trogontherii*(Steppe elephant) bone. J. Biochem., 3, 252-256.
- [21]. Demir Y., Demir N., Nadaroğlu H., Bakan E. (2000).Purification and Characterization of Carbonic Anhydrase From Bovine Erythrocyte Plasma Membrane. *Prep. Biochem. Biothech.*, 30(1), 49-59, 2000.
- [22]. Demir N., Demir Y., Nadaroğlu H. (2001). Carbonic Anhydrase from Bovine Bone. *Prep. Biochem. & Biotech.*, *31*(1), 33-48.
- [23]. Sly, S.W., Peigi, Y. Hu., (1995). Human carbonic anhydrases and carbonic anhydrase deficiencies. Annu. *Rev. Biochem.*, 64, 375-401.
- [24]. Demir Y., Nadaroğlu H., Demir N. (2005). Purification and Characterization of Carbonic Anhydrase From Bovine Stomach And Inhibitory Effects Of Some Chemical Substances On Enzyme Activities. J. Enzym. Inhib. Med. Chem., 20, 75-80.
- [25]. Bradford, M.M. (1976). A Rapid and Sensitive Method for the Quantitation of Microgram Quantities of Protein Utilizing the Principle of Protein-Dye Binding. *Annal. Biochem.*, 72, 248-254.
- [26]. Altıntop M.D., Sever B., Çiftçi G.A., Kucukoglu K., Ozdemir A., Soleimani S.S., Nadaroglu H., Kaplancıklı Z.A. (2017). *Eur. J. Med. Chem.*, 125, 190-196.
- [27]. Celik, H., Maman, M., Babagil, A. (2018). hCA isoenzyme inhibitory activities of Schiff bases derived from cinnamaldehyde and p-methoxycinnamaldehyde, *Fresenius Environmental Bulletin*, 27, 4469-4482.