

PAPER DETAILS

TITLE: Phytochemistry, Ethnomedicinal Uses and Pharmacological Properties of Rhamnus
prinoides: A Review

AUTHORS: Gashaw NIGUSSIE, Mekdes ALEMU, Fozia IBRAHIM, Yemane WEREDE, Metasebia
TEGEGN, Sebsib NEWAY, Milkyas ENDALE ANNISA

PAGES: 136-151

ORIGINAL PDF URL: <https://dergipark.org.tr/tr/download/article-file/1422920>

Phytochemicals, Traditional Uses and Pharmacological Activity of *Rhamnus prinoides*: A Review

Gashaw Nigussie ^{1,2,*}, Mekdes Alemu ¹, Fozia Ibrahim ¹, Yemane Werde ³, Metasebia Tegegn ¹, Sebsib Neway ¹, Milkyas Endale ²

¹Armauer Hansen Research Institute, Addis Ababa, P.O. Box 1005, Addis Ababa, Ethiopia

²Adama Science and Technology University, Adama, P.O. Box 1888, Adama, Ethiopia

³Ministry of Industry, P.O. Box 704, Addis Ababa, Ethiopia

Abstract: *Rhamnus prinoides* L'Herit belongs to Rhamnaceae family widely distributed in India, Eastern, Central and Southern Africa. In Ethiopia it is commonly known as Gesho (Amharic, Tigrigna and Afan Oromo), Gishe (Guragegna) and Geshu (Agewgna). Decoction of the leaves is recommended as remedy for treatment of variety of diseases such as back pain, malaria, pneumonia, sexually transmitted disease, skin infections, wounds, blood purifiers, water borne diseases and as ethnoveterinary medicine. Various secondary metabolites such as flavonoids, alkaloids, tannins, terpenoids, saponins, steroids and anthraquinones have been reported from the genus of which polyphenols were abundant with tremedous antioxidant, wound healing and antiinflammatory activities. The plant also serve as hopping agent, making traditional alcoholic beverages like tella and tej (in Ethiopia), animal feed, medicine, nectar for bees, soil conservation, ornamental, shade and dyes in textiles. This review presents chemical profile as well as biological activities of the species which confirmed that the plant is a good source natural polyphenols and provided valuable information in support of its use as an alternative medicine for future healthcare practice.

ARTICLE HISTORY

Received: Nov. 30, 2020

Revised: Feb. 24, 2021

Accepted: Apr. 19, 2021

KEYWORDS

Traditional uses

Chemical constituents

Antiinflammatory activity

Antioxidant activity

Rhamnus prinoides.

1. INTRODUCTION

Traditional and complementary medicine is an essential and often underestimated health resource with many applications, especially in prevention and management of lifestyle-related chronic diseases. Many countries are expanding coverage of essential health services at a time when consumer expectations for care are rising, costs are soaring and most budgets are either stagnant or being reduced contributing to the revival of alternative traditional medicine. In developing countries, traditional medicine is the main source of healthcare and sometimes the only source of care, due to its closeness to the ordinary rural communities and its accessibility and affordability in view of the rising healthcare costs (Abebe *et al.*, 2003).

Rhamnus prinoides (Figure 1), known as Gesho (Amharic, Tigrigna and Afan Oromo), Gishe (Guragegna) and Geshu (Agewgna) in Ethiopia, belongs to *Rhamnaceae* family (Tesema *et al.*, 1993) widely distributed in India, East, Central and South African countries (Dlamini and Turner, 2002). The only two *Rhamnus* species that occur in Africa are *R. prinoides* and *R.*

*CONTACT: Gashaw Nigussie ✉ gashawnigussie20@gmail.com 📍 Armauer Hansen Research Institute, P.O. Box 1005, Addis Ababa, Ethiopia

Staddo (Abegaz et al., 1999). The genus name '*Rhamnus*' is derived from an ancient Greek word *rhamnos* for blackthorn meaning a 'tuft of branches' and the species name *prinoides* means 'resembling prion' which is an ever green oak. The genus *Rhamnus* comprises about 150 species of shrubs and small trees occurs both in temperate and tropical countries (Chen and Schirarend, 2007). *R. prinoides* also known as *R. pauciflorus* Hochst. ex A. Rich or *R. celifolius* Kallogg is a climbing shrub or small tree that grows up to 4 m high (Schmidt et al., 2002). In Ethiopia, *R. prinoides* (Gesho) is used to add flavour to the local drinks, tella and tej brewed from fermented barley, sorghum or finger millet (d'Avigdor et al., 2014; Tesema et al., 1993). In Angola, the bark is used to induce vomiting (Dlamini and Turner, 2002). In South Africa, a decoction of the root is taken to cleanse the blood by the Zulu and to treat pneumonia by the Sotho (Coates-Palgrave, 1988). The boiled soup of the root is used for the treatment of common cold, leaves are used for chest pain and leaves as well as stems are used for the treatment of tonsil in central Kenya (Njoroge & Bussmann, 2006). In a related study in Kenya, it was reported that the leaves are used in preparation of the "muteta" soup used as an appetizer (Mwangi, 2005).

The decoction of the leaves is recommended as a remedy for treatment of a variety of diseases such as back pain, malaria, pneumonia, sexually transmitted disease, skin infections, wounds, blood purifiers, water borne diseases and as ethnoveterinary medicine (Berhanu, 2014; Enyew et al., 2014; Fratkin, 1996; Megersa et al., 2013; Muthee et al., 2011; Njoroge & Bussmann, 2006; Prozesky et al., 2001; Sobiecki, 2002). The leaves are mainly used in traditional medicine followed by roots (Araya et al., 2015; Diallo et al., 2002; Giday et al., 2010; Seid & Aydagnehum, 2013). Various classes of secondary metabolites such as flavonoids, alkaloids, tannins, terpenoids, saponins, steroids and anthraquinones have been reported from the genus of which polyphenols were abundant with tremendous antioxidant and anti-inflammatory activities (Chen et al., 2020). The plant also serves as an important hopping agent, making traditional alcoholic beverages like tella and tej (in Ethiopia), animal feed, medicine, nectar for bees, soil conservation, ornamental, shade and dyes in textiles. This review presents a comprehensive overview of the chemical profile as well as biological activities of the plant underlining remarkable activities demonstrated by various parts of the plant supporting its potential use as an alternative medicine for future healthcare practice.

Figure 1. Photo of *Rhamnus prinoides* (Amabye, 2015; Molla, 2015, respectively).



1.1. Botanical Distribution

The plant grows up to 8 m in height and has globular, glossy and serrated leaves. The leaves size varies from 10-125 mm (Vetter, 1997). It is commonly cultivated between altitudes of 1400 to 3200 m above sea level (Nielsen, 1992). The flowers are light yellow-green, solitary up to 20 mm (Orwa et al., 2009). The plant is commonly distributed in African countries such as Tanzania, Uganda, exotic to Kenya, Ethiopia, Eritrea, Angola, Malawi, Mozambique, Zambia,

Zimbabwe, Cameron, Democratic Republic of Congo and South Africa (Abegaz, 1996; Dlamini & Turner, 2002; Edwards, 1991; Orwa *et al.*, 2009).

1.2. Ethnomedicinal uses

The leaves, fruits, roots, seeds, shoots and barks are reported to possess diverse traditional medicinal uses to heal various human as well as animal diseases (Table 1). In Tanzania, a root decoction of *R. prinoides* is mixed with bark of *Erythrina abyssinica* Lam. ex DC as remedy for colics (Chhabra *et al.*, 1984). In Ethiopia, the leaf often mixed with the root of *Rubus apetalus* Poir., is boiled, decocted and drunk before meal as remedy for sexually transmitted diseases (Tuasha *et al.*, 2018). The leaf decoction is used for treatment of animal diarrhea and intestinal parasites (Bekele & Musa, 2009), hepatitis (Yineger *et al.*, 2007) and leech infestation (Bekele *et al.*, 2012).

1.3. Economic Uses

R. prinoides has remarkable importance for nutrition, medicine, or other religious purposes in Africa. The leaves and stems are used to add flavor for the preparation of traditional alcoholic beverages like tella and tej (in Ethiopia) by mixing with other components responsible for the bitter taste. It maintains acidic pH during tella fermentation, so it is used to modify the nature of mesh growth of micro-flora and inhibits the growth of undesirable microorganisms (Kebede, 1994; Van Vuuren *et al.*, 1979). It is estimated that close to 5 million people consume it every day (Abegaz & Kebede, 1995; Ashenafi, 2006; Hayeshi *et al.*, 2004). The leaves and stems also serve as a commercial hopping agent in the brewery industry that can be used as an alternative substitute hop (Berhanu, 2014). In a related study, the leaves are also proved to serve as potential dye with direct affinity to cotton fabrics (Kechi *et al.*, 2013; Tewachew *et al.*, 2018). Farmers also use the plant to protect soil erosion and retaining soil on sloping land (Fernandes *et al.*, 1984).

1.4. Nutritional Content Analysis

Nutritional content analysis of *R. prinoides* showed that it has protein (8.5%), fiber (25.6%), ash (9.5%), carbohydrate (70.5%), moisture (9.5%) and fat (3.5%) in support of its nutritional significance (Amabye, 2015)

Table 1. Ethnomedicinal uses of *R. prinoides* in Africa.

Disease treated	Plant part	Mode of preparation and administration	Country/region
Sexually transmitted diseases, arthritis, flu/cold, back pains, stomachache, pneumonia, brucellosis, strength/nutrient supplement, enhancing/facilitating digestion	Tree	-	Kenya
Ear, nose and throat (ENT) diseases (chest pain, common cold and Tonsil)	Roots, Leaves, Stems	Roots, Leaves and Roots/ stems boiling taking orally respectively	Kenya
Tinea capitis, Itching/skin rash	Seeds, Leaves	Rubbed seeds on the affected part and Burn leaves in oven, crush, mix it with butter and apply on the skin	Ethiopia
Tonsillitis	Leaves	Leaves infusion Gargle to rinse the throat	Ethiopia
Skin infection	Leaves	Fresh leaves rubbed it the affected part	Ethiopia
Wound	Leaves	Crush the leaf and apply to the wound till the wound cures.	Ethiopia
Tonsillitis	Leaves	Mothers chewing the leaves and spit to mouth of their children whereas young ones chew it for themselves	Ethiopia
Eczema	Leaves	Leaves crushed, mixed with pure butter and dressing the affected part	
Scabies	Leaves	The leaf is crushed, powdered, mixed with butter and creamed Wounds and exposing the sunlight for few minutes.	Ethiopia
Tonsillitis	Fruits & Shoots	Three fruits and shoots are crushed with three fruits of malt barley, squeezed and droplets of juice are taken through the nose.	
Ring worm	Seed	Adding the grinded powder	Ethiopia

Table 1. *Continued.*

Fever in children	Leaves	Adding the leaf in to water, stay for a while and give the drop of this Concoction to the child	Ethiopia
Sedative, gonorrhea, blood purifier, colics	Leaves, roots	Decoction of leaves is sedative; decoction of roots drunk against gonorrhoea and rheumatism; its decoction mixed with that of the bark of <i>Erythrinaabyssinica</i> Lam. ex DC alleviates colics; decoction of the roots is considered blood purifier and plant liniment is used for sprains	Tanzania
Malaria	Bark, roots	-	South Africa
Blood cleaning, pneumonia, rheumatism, sprains, gargle, skin complaints, respiratory infections, sexually transmitted disease, arthritis, stomach ache, headache	Fruit, leaves, stems, seeds, roots	-	South Africa
Sexual Transmitted Infection	Root	A 3 finger thick pieces of root palm length are boiled in 2 liters of water. 200mls is taken 3 times daily for 7 days	Kenya
Malaria, backaches	Root bark	Root bark boiled all day and drunk with goat fat for malaria, backaches. Drink only one cup, one time	Kenya

Table 1. *Continued.*

Sexually transmitted diseases (STDs)	Leaves	The leaf, often with the root of <i>Rubus apetalus</i> Poir., is boiled, decocted, and drunk before meal	Ethiopia
Tuberculosis	Leaves	-	Uganda
Dandruff	Leaves	-	Ethiopia
Waterborne and related diseases	Leaves		Ethiopia
Animal diarrhea and intestinal parasites	Leaves	Chopped leaves mixed little water and then filtered then taken orally	Ethiopia
Animal hepatitis	Leaves	Crushed and powdered leaves of <i>Cheilanthes farinosa</i> Forsk. Kaulf. and <i>Rhamnus prinoides</i> ; and whole parts of <i>Crepis rueppellii</i> are brushed over the body	Ethiopia
Leech infestation	Leaves	Pounded leaves applied nasally	Ethiopia
Psychoactive agent	Bark	Ground bark that is administered as snuff for mental disorders	South Africa

1.5. Mineral Content

Mineral content analysis of leaves and stem revealed the presence of Ca, Mg, Cr, Mn, Fe, Co, Ni, Cu, Zn, Cd and Pb of which the concentration of Ca and Mg was reported to be high (Gebre and Chandravanshi, 2012, [Table 2](#)). In a related study potassium, sodium and magnesium were the most abundant elements from the leaves and stems samples collected from the low-altitude (1500-1670 m above sea level) and medium-altitude (1670-2000 m above sea level) areas (Nagari & Abebaw, 2013).

Table 2. The mineral content of *R. prinoides*.

Constituents (mg/Kg)	Leaf (mg/kg)	Stem (mg/kg)
Ca	6304-22236	3601- 5675
Mg	3202-5706	2635 - 5568
Cr	5.08 - 20.6	ND - 16.3
Mn	8.12 - 17.9	2.16 - 3.98
Fe	47.9 - 187	22 - 124
Co	22.2 -42.1	18.7 - 91.7
Ni	12.8 - 27.3	9.68 - 19.2
Cu	6.5 - 73.0	16.8 - 233
Zn	12.2 - 43	17.4 - 28.2
Cd	0.81 - 3.10	ND - 1.56
Pb	11.7 - 25	ND

ND – Not detected

1.6. Chemical Constituents

Various class of secondary metabolites including alkaloids, triterpenes, saponins, tannins, phenols, cardiac glycosides, anthraquinones, polyphenols and flavonoids were reported from different parts of *R. prinoides* (Amabye, 2015; Co *et al.*, 1990; Lin *et al.*, 1991; Lin *et al.*, 1990; Molla *et al.*, 2016; Ozipek *et al.*, 1994) as summerized here below ([Figure 2](#), [Table 3](#)).

Campbell and his co-workers (2019) reported the presence of numerous essential oils from the leaves of *R. prinoides* of which 4-hydroxy-4-methyl-2-pentanone (15, [Figure 3](#)) and ethyl 4-ethoxybenzoate (16, [Figure 3](#)) account for more than 85% and exhibited significant antibiofilm activity. In a related study, it has been reported that the essential oils have antibacterial, antifungal, antioxidant and anti-diabetic activities (Sut *et al.*, 2020).

Figure 2. Structures of compounds isolated from *R. prinoides*.





Table 3. Secondary metabolites isolated from *R. prinoides*.

Compound	Plant Part	Reference
Emodin (1)	Fruit, Leaves, stems	(Abegaz & Dagne, 1988; Abegaz & Kebede, 1995; Gebru, 2010)
Emodinanthrone (2)	Leaves, stems, Fruit	(Abegaz & Dagne, 1988; Abegaz & Peter, 1995)
Physcion (3)	Fruit, Leaves, stems	(Abegaz & Dagne, 1988; Abegaz & Kebede, 1995; Gebru, 2010)
Emodinbianthrone (4)	Fruit	(Abegaz & Dagne, 1988; Abegaz & Peter, 1995)
Musizin (5)	Leaves	(Abegaz & Kebede, 1995)
β -sorigenin (6)	Leaves, roots	(Abegaz & Kebede, 1995; Gebru, 2010)
Geshoidin (P-Sorigenin-8-O- β -D-glucoside (7)	Leaves, stems, roots	(Abegaz & Kebede, 1995; Gebru, 2010)
Rhamnocitrin (8)	Leaves	(Abegaz & Kebede, 1995)
Rhamnazin (9)	Leaves, roots	(Abegaz & Kebede, 1995; Gebru, 2010)
Prinoidin (10)	Fruits	(Abegaz & Kebede, 1995; Abegaz & Peter, 1995)
Rhamnetin (11)	Leaves, stems	(Abegaz & Kebede, 1995)
Quercetin (12)	Leaves	(Abegaz & Kebede, 1995)
Chrysophanol (13)	Leaves, stems, roots	(Abegaz & Kebede, 1995; Gebru, 2010)
3-O-Methylquercetin (14)	Leaves, Stems	(Abegaz & Kebede, 1995)

Figure 3. Structure of major constituents of essential oils from leaves of *R. prinoides*.

1.7. Biological Activity

1.7.1. Antioxidant activity

It was reported that *n*-hexane, chloroform, ethyl acetate, and methanolic extracts of the leaf and stem bark of *R. prinoids* possess antioxidant activities by DPPH assay with IC₅₀ values of >3000, >3000, >3000, 950.42, ~1500, 710.50, ~1000 and 902.78 $\mu\text{g mL}^{-1}$, respectively (Pillai *et al.*, 2019). The documented antioxidant activities of *R. prinoides* are probably due to flavonoids and polyphenols reported from leaves, stems and roots (Amabye, 2015; Molla *et al.*, 2016; Pillai *et al.*, 2019).

1.7.2. Antiinflammatory activity

Semi purified ethanolic stem and stem bark extracts of *R. prinoides* with higher contents of polyphenols and flavonoids displayed anti-inflammatory activity through reducing the Nitric

Oxide production at the dosage of 11.11-100 µg/mL and the COX-2 inhibitory activity with an IC₅₀ value at 20.61 ± 0.13 µg/mL (Chen *et al.*, 2020).

1.7.3. Antibiofilm activity

Leaf and stem ethanol extracts of *R. prinoides* showed significant inhibition of *Staphylococcus aureus* Rosenbach, *Bacillus subtilis* and *Streptococcus mutans* Clarke biofilm formation up to 99.9% and reduced planktonic cell growth up to 10 log units relative to untreated controls (Campbell *et al.*, 2019). In a related study, the stem ethanol extracts disrupted *S. mutans* and *C. albicans* co-culture synergism, with 98% less polymicrobial biofilm formation than the untreated control (Campbell *et al.*, 2020).

1.7.4. Antibacterial activity

Antibacterial activity of methanol and chloroform leaves extracts of *R. prinoides* against *Staphylococcus aureus*, *Streptococcus pyogenes* Rosenbach, *Streptococcus pneumoniae* revealed minimum inhibitory concentration ranging from 8.13-32.5 mg/mL and 8.13-16.25 mg/mL, respectively, compared to a positive control cefotaxin (0.03 mg/mL) and ampicillin (0.01 mg/mL) ranged from 23.67- 28.00 mg/mL in both fractions, respectively (Molla *et al.*, 2016).

1.7.5. Antimalarial activity

The chloroform root bark extracts of *R. prinoides* were active against chloroquine sensitive *Plasmodium falciparum* strain with IC₅₀ value of 3.53 µg/mL (Koch *et al.*, 2005). In a related study, *n*-hexane and dichloromethane extracts were demonstrated to have antiplasmodial activity with IC₅₀ values of 19.9 µg/mL and 30.3 µg/mL, respectively, with no toxicity in the brine shrimp test (Bosire, 2003). The *in-vitro* antiplasmodial activities of dichloromethane/methanol (1:1) crude extracts and the isolated compounds were performed against the chloroquine-sensitive (D6) and chloroquine-resistant (W2) strains of *Plasmodium falciparum*. The naphthalenic derivative Geshoidin (7) displayed IC₅₀ value of 4.0 ± 0.9 pM and 0.4 ± 0.2 pM against the chloroquine sensitive (D6) and chloroquine resistant (W2) strains of *Plasmodium falciparum* (Gebbru, 2010). *In vivo* antimalarial activity of aqueous extracts from leaves and root barks against a blood-induced CQ-resistant rodent parasite in mice showed high chemosuppression in the range of 51-75% (Muregi *et al.*, 2007) suggesting the promising application of leaves, stem bark and roots of *R. prinoides* to treat malaria.

1.7.6. Antimycobacterial activity

The acetone leaf extracts of *R. prinoides* demonstrated antimycobacterial activity with minimum inhibitory concentration (MIC) values ranging from 0.625 to >2.5 mg/mL against three fast-growing mycobacteria species *i.e.* *Mycobacterium smegmatis* Trevisan, *Mycobacterium aurum* Tsukamura and *M. fortuitum* Da Costa Cruz and one pathogenic *M. tuberculosis* strain (Dzoyem *et al.*, 2016).

1.7.7. Wound healing activity

In vivo study in mice revealed that the hydroalcoholic extracts of the leaves of *R. prinoides* possess wound healing activity established by a significant rate of wound contraction and shorter epithelization period. In this study, ten percent of 80% methanol leaves extract showed significant wound contraction against the control and rate increased in significant level with number of days $p < 0.05$, $p < 0.01$, and $p < 0.001$ on days 2 to 4, 8 to 10 and 12 to 14, respectively (Tessema *et al.*, 2021). Biological activities of different parts of *R. prinoides* are summerized here below (Table 4).

Table 4. Biological activity of different parts of *R. prinoides*

Plant part	Activity	Reference
Flowers, leaves, stems	Antimutagenic	(Wall <i>et al.</i> , 1988)
Leaves	Antibacterial	(Chhabra & Uiso, 1991; Chhabra <i>et al.</i> , 1984; Molla <i>et al.</i> , 2016)
	Insecticide	(Van Puyvelde <i>et al.</i> , 1985)
Root bark	Cytotoxic	(Koch <i>et al.</i> , 2005)
Leaves, roots, bark root	Antimalarial	(Muregi <i>et al.</i> , 2007)
Leaves	Wound healing	(Tessema <i>et al.</i> , 2021)

2. CONCLUSION

The present review presented chemical constituents and biological activity of *R. prinoides* native to Africa and India. Various secondary metabolites such as flavonoids, alkaloids, tannins, terpenoids, saponins, steroids and anthraquinones have been reported from the genus of which polyphenols were abundant with tremendous antioxidant, wound healing and anti-inflammatory activities. Economically, the leaves and stems have been used as hopping agent in the brewery industry, making traditional alcoholic beverages like tella and tej (Ethiopia), animal feed, medicine, nectar for bees, soil conservation, ornamental, and dyes in textiles. Considering diverse class of secondary metabolites as well as wide spectrum of biological activities of the plant, it is believed that plant is a good natural source of polyphenols and can be used as an alternative medicine for future healthcare practice.

Declaration of Conflicting Interests and Ethics

The authors declare no conflict of interest. This research study complies with research publishing ethics. The scientific and legal responsibility for manuscripts published in IJSM belongs to the authors.

Authorship Contribution Statement

GN, MA, SN, FI, YW, MT: literature compilation, review and analysis. **GN & ME:** manuscript write up and review. All authors have read and approved the manuscript.

ORCID

Gashaw Nigussie  <https://orcid.org/0000-0002-7758-6367>

Mekdes Alemu  <https://orcid.org/0000-0002-5860-7070>

Fozia Ibrahim  <https://orcid.org/0000-0002-4255-4285>

Yemane Werde  <https://orcid.org/0000-0002-5943-1847>

Metasebia Tegegn  <https://orcid.org/0000-0001-5118-0898>

Sebsib Neway  <https://orcid.org/0000-0002-2033-5874>

Milkyas Endale  <https://orcid.org/0000-00025301-9923>

3. REFERENCES

- Abdeta, D., Amante, M., & Tamiru, Y. (2020). Survey on Ethno Botany and Medicinal Animals at Sayo and Hawa Gelan Districts of Kelem Wollega Zone, Western Ethiopia. *Preventive Medicine*, 28(2), 21408-21420. <https://doi.org/10.26717/BJSTR.2020.28.004620>
- Abebe, D., Debella, A., & Urga, K. (2003). Medicinal plants and other useful plants of Ethiopia.
- Abegaz, B., & Dagne, E. (1988). Anthracene derivatives of *Rhamnus prinoides*. *Bulletin of the Chemical Society of Ethiopia*, 2(1), 15-20.

- Abegaz, B. M. (1996). Progress in the chemistry of marketed traditionally used plants of Ethiopia. *Bulletin of the Chemical Society of Ethiopia*, 10(1), 57-71.
- Abegaz, B. M., & Kebede, T. (1995). Geshoidin: A bitter principle of *Rhamnus prinoides* and other constituents of the leaves. *Bulletin of the Chemical Society of Ethiopia*, 9(2), 107-114.
- Abegaz, B. M., Ngadjui, B. T., Bezabih, M., & Mdee, L. K. (1999). Novel natural products from marketed plants of eastern and southern Africa. *Pure and Applied Chemistry*, 71(6), 919-926. <https://doi.org/10.1351/pac199971060919>
- Abegaz, B. M., & Peter, M. G. (1995). Emodin and emodinanthrone rhamnoside acetates from fruits of *Rhamnus prinoides*. *Phytochemistry*, 39(6), 1411-1414. [https://doi.org/10.1016/0031-9422\(95\)00093-M](https://doi.org/10.1016/0031-9422(95)00093-M)
- Amabye, T. G. (2015). Evaluation of phytochemical, chemical composition, antioxidant and antimicrobial screening parameters of *Rhamnus prinoides* (Gesho) available in the market of Mekelle, Tigray, Ethiopia. *Natural Products Chemistry & Research*, 4(1), 1-6.
- Araya, S., Abera, B., & Giday, M. (2015). Study of plants traditionally used in public and animal health management in Seharti Samre District, Southern Tigray, Ethiopia. *Journal of ethnobiology and ethnomedicine*, 11(1), 1-25. <https://doi.org/10.1186/s13002-015-0015-5>
- Ashenafi, M. (2006). A review on the microbiology of indigenous fermented foods and beverages of Ethiopia. *Ethiopian Journal of Biological Sciences*, 5(2), 189-245.
- Asmare, T., Yilkal, B., Mekuannint, T., & Yibeltal, A. (2018). Traditional medicinal plants used to treat maternal and child health illnesses in Ethiopia: an ethno-botanical approach. *Journal of Traditional Medicine and Clinical Naturopathy*, 7(277), 2.
- Bekele, A., & Musa, A. (2009). Ethnoveterinary practice in Chiro District, western Hararge, Ethiopia. *Pharmacologyon Line*, 1, 128-139.
- Bekele, D., Asfaw, Z., Petros, B., & Tekie, H. (2012). Ethnobotanical study of plants used for protection against insect bite and for the treatment of livestock health problems in rural areas of Akaki District, Eastern Shewa, Ethiopia. *Topclass Journal of Herbal Medicine*, 1(2), 12-24.
- Berhanu, A. (2014). Microbial profile of Tella and the role of gesho (*Rhamnus prinoides*) as bittering and antimicrobial agent in traditional Tella (Beer) production. *International Food Research Journal*, 21(1), 357-365.
- Bitew, H., Gebregergs, H., Tuem, K. B., & Yeshak, M. Y. (2019). Ethiopian medicinal plants traditionally used for wound treatment: a systematic review. *Ethiopian Journal of Health Development*, 33(2), 103-112.
- Bosire, K. O. (2003). *An investigation of a traditional herbal therapy used to treat malaria in Kisii* [MSc Thesis] University of Nairobi, Nairobi, Kenya.
- Campbell, M., Fathi, R., Cheng, S. Y., Ho, A., & Gilbert, E. S. (2020). *Rhamnus prinoides* (gesho) stem extract prevents co-culture biofilm formation by *Streptococcus mutans* and *Candida albicans*. *Letters in Applied Microbiology*, 71(3), 294-302.
- Campbell, M., Zhao, W., Fathi, R., Mihreteab, M., & Gilbert, E. S. (2019). *Rhamnus prinoides* (gesho): A source of diverse anti-biofilm activity. *Journal of Ethnopharmacology*, 241, 111955. <https://doi.org/10.1016/j.jep.2019.111955>
- Chen, G.-L., Munyao Mutie, F., Xu, Y.-B., Saleri, F. D., Hu, G.-W., & Guo, M.-Q. (2020). Antioxidant, anti-inflammatory activities and polyphenol profile of *Rhamnus prinoides*. *Pharmaceuticals*, 13(4), 55. <https://doi.org/10.3390/ph13040055>
- Chen, Y., Schirarend, C. (2007). *Rhamnus* L. In: Flora China Vol. 12. Beijing (China): Science Press; St. Louis (MO): Missouri Botanical Garden Press; p. 139-162.
- Chhabra, S., & Uiso, F. (1991). Antibacterial activity of some Tanzanian plants used in traditional medicine. *Fitoterapia*, 62(6), 499-503.

- Chhabra, S., Uiso, F., & Mshiu, E. (1984). Phytochemical screening of Tanzanian medicinal plants. I. *Journal of Ethnopharmacology*, 11(2), 157-179. [https://doi.org/10.1016/0378-8741\(84\)90037-0](https://doi.org/10.1016/0378-8741(84)90037-0)
- Co, M., Toshiko, S., Hori, K., Saiki, Y., & Tanker, M. (1990). Anthraquinone glycosides from *Rhamnus libanoticus*. *Phytochemistry*, 29(6), 2018-2020. [https://doi.org/10.1016/0031-9422\(90\)85060-S](https://doi.org/10.1016/0031-9422(90)85060-S)
- Coates-Palgrave, K. (1988). Trees of Southern Africa-Tamarindus indica. *LCS Striuk Publishers, Cape Town*, 10, 278-279.
- Cock, I., Selesho, M., & Van Vuuren, S. (2019). A review of the traditional use of southern African medicinal plants for the treatment of malaria. *Journal of Ethnopharmacology*, 245, 112176. <https://doi.org/10.1016/j.jep.2019.112176>
- d'Avigdor, E., Wohlmuth, H., Asfaw, Z., & Awas, T. (2014). The current status of knowledge of herbal medicine and medicinal plants in Fiche, Ethiopia. *Journal of ethnobiology and ethnomedicine*, 10(1), 1-33. <https://doi.org/10.1186/1746-4269-10-38>
- Diallo, D., Sogn, C., Samaké, F. B., Paulsen, B. S., Michaelsen, T. E., & Keita, A. (2002). Wound healing plants in Mali, the Bamako region. An ethnobotanical survey and complement fixation of water extracts from selected plants. *Pharmaceutical Biology*, 40(2), 117-128.
- Dlamini, M., & Turner, S. (2002). *Rhamnus prinoides* L'Hérit. *Witwatersran d National Botanical Garden: South African National Biodiversity Institute, South Africa*.
- Dzoyem, J. P., Aro, A. O., McGaw, L. J., & Eloff, J. N. (2016). Antimycobacterial activity against different pathogens and selectivity index of fourteen medicinal plants used in southern Africa to treat tuberculosis and respiratory ailments. *South African Journal of Botany*, 102, 70-74. <https://doi.org/10.1016/j.sajb.2015.08.002>
- Edwards, S. B. (1991). Crops with wild relatives found in Ethiopia. *Plant genetic resources of Ethiopia*, 42-47.
- Enyew, A., Asfaw, Z., Kelbessa, E., & Nagappan, R. (2014). Ethnobotanical study of traditional medicinal plants in and around Fiche District, Central Ethiopia. *Current Research Journal of Biological Sciences*, 6(4), 154-167.
- Fernandes, E. C., Oktingati, A., & Maghembe, J. (1984). The Chagga homegardens: a multistoried agroforestry cropping system on Mt. Kilimanjaro (Northern Tanzania). *Agroforestry Systems*, 2(2), 73-86.
- Fratkin, E. (1996). Traditional medicine and concepts of healing among Samburu pastoralists of Kenya. *Journal of Ethnobiology*, 16, 63-98.
- Gakuya, D., Kabasa, J. D., & Kiama, S. (2015). Medicinal plants used in the management of sexually transmitted infections by the Samburu community, Kenya. *International Journal of Pharmaceutical Research*, 7(2), 45.
- Gebeyehu, G. (2016). *An ethnobotanical study of traditional use of medicinal plants and their conservation status in Mecha Wereda, West Gojam Zone of Amhara Region, Ethiopia* [MScThesis]. Addis Ababa University, Addis Ababa, Ethiopia.
- Gebre, A., & Chandravanshi, B. S. (2012). Levels of essential and non-essential metals in *Rhamnus prinoides* (Gesho) cultivated in Ethiopia. *Bulletin of the Chemical Society of Ethiopia*, 26(3), 329-342.
- Geburu, M. (2010). *Phytochemical and antiplasmodial investigation of Rhamnus prinoides and Kniphofia foliosa* [MSc Thesis]. University of Nairobi, Nairobi, Kenya.
- Giday, M., Asfaw, Z., & Woldu, Z. (2010). Ethnomedicinal study of plants used by Sheko ethnic group of Ethiopia. *Journal of Ethnopharmacology*, 132(1), 75-85. <https://doi.org/10.1016/j.jep.2010.07.046>

- Hayeshi, R., Mukanganyama, S., Hazra, B., Abegaz, B., & Hasler, J. (2004). The interaction of selected natural products with human recombinant glutathione transferases. *Phytotherapy Research*, 18(11), 877-883.
- Hishe, M., & Asfaw, Z. (2014). Review paper review on ethnobotanical studies on traditional medicinal plants used to treat livestock and human ailments in tigray region, Ethiopia. *AJBSR*, 3, 8-36.
- Kebede, T. (1994). *The bitter constituent of Gesho (Rhamnus prinoides) leaves* [MSc thesis]. Addis Ababa University, Addis Ababa, Ethiopia.
- Kechi, A., Chavan, R., & Moeckel, R. (2013). Ethiopian Dye plants As a Source of Natural Dyes for Cotton Dyeing. *Universal Journal of Environmental Research & Technology*, 3(4), 501-510.
- Kidane, B., van Andel, T., van der Maesen, L. J. G., & Asfaw, Z. (2014). Use and management of traditional medicinal plants by Maale and Ari ethnic communities in southern Ethiopia. *Journal of Ethnobiology and Ethnomedicine*, 10(1), 1-15. <https://doi.org/10.1186/1746-4269-10-46>
- Kiringe, J. W. (2006). A survey of traditional health remedies used by the Maasai of Southern Kaijiado District, Kenya. *Ethnobotany Research and Applications*, 4, 061-074.
- Koch, A., Tamez, P., Pezzuto, J., & Soejarto, D. (2005). Evaluation of plants used for antimalarial treatment by the Maasai of Kenya. *Journal of Ethnopharmacology*, 101(1-3), 95-99. <https://doi.org/10.1016/j.jep.2005.03.011>
- Lin, C.-N., Chung, M.-I., Gan, K.-H., & Lu, C.-M. (1991). Flavonol and anthraquinone glycosides from *Rhamnus formosana*. *Phytochemistry*, 30(9), 3103-3106. [https://doi.org/10.1016/S0031-9422\(00\)98262-1](https://doi.org/10.1016/S0031-9422(00)98262-1)
- Lin, C.-N., Chung, M.-I., & Lu, C.-M. (1990). Anthraquinones from *Rhamnus formosana*. *Phytochemistry*, 29(12), 3903-3905. [https://doi.org/10.1016/0031-9422\(90\)85356-K](https://doi.org/10.1016/0031-9422(90)85356-K)
- Megersa, M., Asfaw, Z., Kelbessa, E., Beyene, A., & Woldeab, B. (2013). An ethnobotanical study of medicinal plants in Wayu Tuka district, east Welega zone of oromia regional state, West Ethiopia. *Journal of Ethnobiology and Ethnomedicine*, 9(1), 1-18. <https://doi.org/10.1186/1746-4269-9-68>
- Mesfin, K., Tekle, G., & Tesfay, T. (2013). Ethnobotanical study of traditional medicinal plants used by indigenous people of Gemad District, Northern Ethiopia. *Journal of Medicinal Plants Studies*, 1(4), 32-37.
- Molla, Y. (2015). *Evaluation of the antibacterial activity of the solvent fractions of the leaves of Rhamnus prinoides L'Herit (Rhamnaceae)* [Msc Thesis]. Addis Ababa University, Addis Ababa, Ethiopia.
- Molla, Y., Nedi, T., Tadesse, G., Alemayehu, H., & Shibeshi, W. (2016). Evaluation of the in vitro antibacterial activity of the solvent fractions of the leaves of *Rhamnus prinoides* L'Herit (Rhamnaceae) against pathogenic bacteria. *BMC Complementary and Alternative Medicine*, 16(1), 1-9. <https://doi.org/10.1186/s12906-016-1279-6>
- Muregi, F. W., Ishih, A., Suzuki, T., Kino, H., Amano, T., Mkoji, G. M., Miyase, T., & Terada, M. (2007). In Vivo antimalarial activity of aqueous extracts from Kenyan medicinal plants and their Chloroquine (CQ) potentiation effects against a blood-induced CQ-resistant rodent parasite in mice. *Phytotherapy Research*, 21(4), 337-343.
- Muthee, J., Gakuya, D., Mbaria, J., Kareru, P., Mulei, C. M., & Njonge, F. (2011). Ethnobotanical study of anthelmintic and other medicinal plants traditionally used in Loitokitok district of Kenya. *Journal of Ethnopharmacology*, 135(1), 15-21. <https://doi.org/10.1016/j.jep.2011.02.005>
- Mwangi, P. (2005). *The evaluation of the cognitive effects of Rhamnus prinoides; a single-blind randomized crossover trial* [Doctoral dissertation]. University of Nairobi, Nairobi, Kenya.

- Nagari, A., & Abebaw, A. (2013). Determination of Selected Essential and Non-essential Metals in the Stems and Leaves of *Rhamnus prinoides* (gesho). *Science, Technology and Arts Research Journal*, 2(4), 20-26.
- Nielsen, I. (1992). Hedberg, I. & Edwards, S.(eds), 1989. Flora Ethiopia vol. 3, Pittosporaceae to Araliaceae. *Nordic Journal of Botany*, 12(4), 402-402. <https://doi.org/10.1111/j.1756-1051.1992.tb01322.x>
- Njoroge, G. N., & Bussmann, R. W. (2006). Traditional management of ear, nose and throat (ENT) diseases in Central Kenya. *Journal of ethnobiology and ethnomedicine*, 2(1), 1-9. <https://doi.org/10.1186/1746-4269-2-54>
- Orwa, C., Mutua, A., Kindt, R., Jamnadass, R., & Simons, A. (2009). Agroforestry Database: a tree reference and selection guide. Version 4. *Agroforestry Database: a tree reference and selection guide. Version 4*.
- Ozipek, M., Calış, I., Ertan, M. t., & Rüedi, P. (1994). Rhamnetin 3-p-coumaroylrhamninoside from *Rhamnus petiolaris*. *Phytochemistry*, 37(1), 249-253. [https://doi.org/10.1016/0031-9422\(94\)85035-6](https://doi.org/10.1016/0031-9422(94)85035-6)
- Pillai, M. K., Santi, L. I., & Magama, S. (2019). DPPH radical scavenging activity of extracts from *Rhamnus prinoides*. *Journal of Medicinal Plants Research*, 13(15), 329-334.
- Prozesky, E., Meyer, J., & Louw, A. (2001). In vitro antiplasmodial activity and cytotoxicity of ethnobotanically selected South African plants. *Journal of Ethnopharmacology*, 76(3), 239-245.
- Schmidt, E., Lotter, M., & McClelland, W. (2002). *Trees and shrubs of Mpumalanga and Kruger national park* (J. E. Burrows, Ed.). Jacana Media.
- Seid, M. A., & Aydagnehum, S. G. (2013). Medicinal plants biodiversity and local Healthcare management system in Chenchu District; Gamo Gofa, Ethiopia. *Journal of Pharmacognosy and Phytochemistry*, 2(1), 284-293.
- Siyum, D., & Woyessa, D. (2013). Assessment of bacteriological quality and traditional treatment methods of water-borne diseases among well water users in Jimma Town, South West Ethiopia. *ARPJ Ag & Bio Sci*, 8, 477-486.
- Sobiecki, J.-F. (2002). A preliminary inventory of plants used for psychoactive purposes in southern African healing traditions. *Transactions of the Royal Society of South Africa*, 57(1-2), 1-24. <https://doi.org/10.1080/00359190209520523>
- Sut, S., Dall'Acqua, S., Sinan, K. I., Bene, K., Kumar, G., Mahomoodally, M. F., Picot-Allain, C., & Zengin, G. (2020). Cola caricifolia (G. Don) K. Schum and Crotalaria retusa L. from Ivory Coast as sources of bioactive constituents. *Industrial Crops and Products*, 147, 112246.
- Tabuti, J. R., Kukunda, C. B., & Waako, P. J. (2010). Medicinal plants used by traditional medicine practitioners in the treatment of tuberculosis and related ailments in Uganda. *Journal of Ethnopharmacology*, 127(1), 130-136. <https://doi.org/10.1016/j.jep.2009.09.035>
- Tamene, S. (2020). Ethnobotanical study of indigenous knowledge on medicinal plant uses and threatening factors around the Malga District, Southern Ethiopia. *International Journal of Biodiversity and Conservation*, 12(3), 215-226.
- Tegene, A. S. (2018). An Ethnobotanical Study of Traditionally Used Medicinal Plants for Treatment of Human Diseases in Goba District of Bale Zone, Southeast Ethiopia. *Advances in Life Science and Technology*, 68, 1-7.
- Teka, A., Asfaw, Z., Demissew, S., & Van Damme, P. (2020). Traditional uses of medicinal plants practiced by the indigenous communities in Gurage Zone, south central Ethiopia. *Ethnobotany Research and Applications*, 19, 1-31.

- Teklay, A., Abera, B., & Giday, M. (2013). An ethnobotanical study of medicinal plants used in Kilte Awulaelo District, Tigray Region of Ethiopia. *Journal of Ethnobiology and Ethnomedicine*, 9(1), 1-23. <https://doi.org/10.1016/j.jep.2009.09.035>
- Teklehaymanot, T., & Giday, M. (2007). Ethnobotanical study of medicinal plants used by people in Zegie Peninsula, Northwestern Ethiopia. *Journal of ethnobiology and ethnomedicine*, 3(1), 1-11. <https://doi.org/10.1186/1746-4269-3-12>
- Tesema, A. B., Ann, B., & Bo, T. (1993). Useful Trees and Shrubs for Ethiopia: Identification, Propagation and Management for Agricultural and Pastoral Communities. <http://hdl.handle.net/123456789/3008>
- Tessema, Z., Yibeltal, D., & Molla, Y. (2021). Evaluation of the wound healing activity of the crude extract of root bark of Brucea antidysentrica, the leaves of Dodonaea angustifolia and Rhamnus prinoides in mice. *Heliyon*, 7(1), 1-8. <https://doi.org/10.1016/j.heliyon.2021.e05901>
- Tewachew, A., Mitkie, H., & Gebeyehu, Y. (2018). *The production and characterization of dye from local Gesho for the application of textile industries* [Doctoral dissertation]. Bahir dar University, Bahir dar, Ethiopia.
- Tuasha, N., Petros, B., & Asfaw, Z. (2018). Medicinal plants used by traditional healers to treat malignancies and other human ailments in Dalle District, Sidama Zone, Ethiopia. *Journal of Ethnobiology and Ethnomedicine*, 14(1), 1-21. <https://doi.org/10.1186/s13002-018-0213-z>
- Van Puyvelde, L., Geysen, D., Ayobangira, F.-X., Hakizamungu, E., Nshimiyimana, A., & Kalisa, A. (1985). Screening of medicinal plants of Rwanda for acaricidal activity. *Journal of Ethnopharmacology*, 13(2), 209-215. [https://doi.org/10.1016/0378-8741\(85\)90008-X](https://doi.org/10.1016/0378-8741(85)90008-X)
- Van Vuuren, H., Loos, M., Louw, H., & Meisel, R. (1979). Distribution of bacterial contaminants in a South African lager brewery. *Journal of Applied Bacteriology*, 47(3), 421-424.
- Vetter, S. (1997). *Geisho: its uses, production potential and problems in Northern Tigray, Ethiopia*. <http://hdl.handle.net/123456789/3065>
- Wall, M. E., Wani, M. C., Hughes, T. J., & Taylor, H. (1988). Plant antimutagenic agents, 1. General bioassay and isolation procedures. *Journal of Natural Products*, 51(5), 866-873. <https://doi.org/10.1021/np50059a009>
- Yineger, H., Kelbessa, E., Bekele, T., & Lulekal, E. (2007). Ethnoveterinary medicinal plants at bale mountains national park, Ethiopia. *Journal of Ethnopharmacology*, 112(1), 55-70. <https://doi.org/10.1016/j.jep.2007.02.001>