# PAPER DETAILS

TITLE: Utilization of microalgae [Chlorella vulgaris Beyerinck (Beijerinck)] on plant growth and nutrient uptake of garden cress (Lepidium sativum L.) grown in different fertilizer applications AUTHORS: Aynur SADAK,Suat SENSOY PAGES: 240-245

ORIGINAL PDF URL: https://dergipark.org.tr/tr/download/article-file/2165621



**Research Article** 

International Journal of Agriculture, Environment and Food Sciences

e-ISSN: 2618-5946

Int J Agric Environ Food Sci 6 (2): 240-245 (2022)

www.iaefs.com

# Utilization of microalgae [*Chlorella vulgaris* Beyerinck (Beijerinck)] on plant growth and nutrient uptake of garden cress (*Lepidium sativum* L.) grown in different fertilizer applications

Aynur Sadak Turhan<sup>1,\*</sup> <sup>(D)</sup>

Abstract



<sup>1</sup>Van Yuzuncu Yil University, The Institute of Natural and Applied Sciences, Horticultural Sciences, Van, Türkiye <sup>2</sup>Van Yuzuncu Yil University, Agricultural Faculty, Horticulture Department, Van, Türkiye

\*Corresponding Author: aynuursadak@gmail.com

#### Citation

Sadak Turhan, A., Şensoy, S. (2022). Utilization of microalgae [Chlorella vulgaris Beyerinck (Beijerinck)] on plant growth and nutrient uptake of garden cress (*Lepidium sativum* L.) grown in different fertilizer applications. International Journal of Agriculture, Environment and Food Sciences, 6 (2), 240-245.

Doi: https://doi.org/10.31015/jaefs.2022.2.6

Received: 29 December 2021 Accepted: 30 May 2022 Published Online: 12 June 2022 Revised: 27 June 2022

Year: 2022 Volume: 6 Issue: 2 (June) Pages: 240-245



This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY-NC) license https://creativecommons.org/licenses/by-nc/4.0/

**Copyright © 2022** International Journal of Agriculture, Environment and Food Sciences; Edit Publishing, Diyarbakır, Türkiye.

Available online http://www.jaefs.com https://dergipark.org.tr/jaefs

#### Introduction

In response to the constantly increasing population in the world, the need for food is also increasing rapidly (Sencar, 1988). Due to the continuous increase in the world population, agricultural areas should also be increased, which is not so feasible. Therefore, the amount of product obtained from the unit area should be increased or the existing areas used should be sustainable (Midmore, 1993). The soil structure of Turkey is generally poor in terms of organic matter (Kaçar and İnal, 2008). It is also known that the intensive use of artificial chemical fertilizers in agricultural lands deteriorates the soil structure and pollutes the underground water resources, and the

This study was employed as a pot experiment in the controlled greenhouse conditions in order to investigate the effect of microalgae [Chlorella vulgaris Beyerinck (Beijerinck)] application on plant growth and nutrient uptake of garden cress (Lepidium sativum L.) grown in different doses of mineral fertilizer applications. Sieved soil in 3-liter pots was used as the growing medium. Equal amount of irrigation was applied to all pots during the period from seed sowing to the end of the experiment. Microalgae application was applied twice (100 ml and 150 ml per pot) to the seedling-growing medium. As chemical fertilizer, 0%, 50% and 100% of NPK (160 mg N kg<sup>-1</sup>, 80 mg  $P_2O_5$  kg<sup>-1</sup>, and 100 mg K<sub>2</sub>O kg<sup>-1</sup>) were applied. As the parameters in garden cress, shoot height, shoot fresh weight, total soluble content (TSS), chlorophyll amount (SPAD value) and some nutrients element (K, Ca, Fe, Zn, Cu and Mn) contents were examined. At the end of the study, microalgae applications were found to have a promising effect on plant growth and some nutrient uptake. It was observed that the values of the studied traits generally increased in the microalgae application compared to the control group.

#### Keywords

Bio-fertilizer, Garden cress, Microalgae, Nutrient content, Plant growth

excessive use of these chemicals disrupts the ecological balance (Turan, 2007). The sustainability of the solutions to be produced in response to the problems encountered in production is an important criterion (Koru and Cirik, 1999). Moreover, it is important to use sustainable resources correctly in terms of environment and human health (Kut et al., 2007).

Garden cress (*Lepidium sativum* L.), is included in the Brassicaceae family. It is a rather fast-growing, edible herb; it is an annual herbaceous plant; it is used as an appetizer with its pleasant peppery, tangy flavor and aroma (Vural et al., 2000). It is also prized as a medicinal herb especially for diabetes (Tahraoui et al., 2007).

*Chlorella*, a genus of single-celled green algae and defined as the oldest living thing known for 2.5 billion years, is an important food source. *Chlorella* is dark green. Its greenery is high in chlorophyll. It is known as green freshwater algae (Kut et al., 2007). In recent years, different microalgae have been used as plant nutrients in agricultural production (Okur et al., 2001). Today, the widespread use of microalgae fertilizers is getting prominent (Eşiyok et al., 2001). It is known that microalgae have positive effects on plant growth and development. Moreover, it increases the yield and quality of the agricultural crops and is used as a biofertilizer in agriculture (Engin et al., 2019). Schreiber et al. (2018) reported that algal biomass could also support crop growth on marginal soils.

This study aimed to investigate the effects of microalgae [*Chlorella vulgaris* Beyerinck (Beijerinck)] use on plant growth and nutrient uptake in garden cress (*Lepidium sativum* L.) in different doses of mineral

fertilizer applications to see if it can compensate for the use of artificial fertilizers to some extent.

### Materials and methods

In the present study, the effect of microalgae [*Chlorella vulgaris* Beyerinck (Beijerinck)] use on plant growth in garden cress [*Lepidium sativum* L. cv. Helen (Sim Arzuman Seeds)] was investigated. The study was carried out as a pot experiment in controlled greenhouse conditions  $(15\pm 4 \text{ C}^{\circ} \text{ night} \text{ and } 28\pm 4 \text{ C}^{\circ} \text{ day})$ . Microalgae application was applied twice (100 ml and 150 per pot) to the seedling growing medium and as mineral fertilizer, 0%, 50% and 100% of NPK (160 mg N kg<sup>-1</sup>, 80 mg P<sub>2</sub>O<sub>5</sub> kg<sup>-1</sup>, and 100 mg K<sub>2</sub>O kg<sup>-1</sup> as 20:20:20 NPK, Potassium sulfate, and urea) were applied as base fertilizer and 40 mg N kg<sup>-1</sup> as urea two weeks after seedling emergence (Table 1). The soil properties used in the experiment are given in Table 1.

Sieved soil in 3-liter pots was used as the growing medium, and 40 seeds were sown in each pot. Equal amount of irrigation was applied to all pots during the period from seed sowing to the end of the experiment, is about 6.5 weeks.

Table 1. Applications and soil properties

Application		
Control (no microalgae and no fertilizer)		
%50 NPK fertilizer		
%100 NPK fertilizer		
Microalgae		
%50 NPK fertilizer + Microalgae		
%100 NPK fertilizer + Microalgae		
Soil properties		
580.2	High	
31.2	Little	
11.025	Medium	
1.98	Little	
0.036	No salt	
7.21	Neutral	
	ApplicationControl (no microalgae and no fertilizer)% 50 NPK fertilizer% 100 NPK fertilizerMicroalgae% 50 NPK fertilizer + Microalgae% 100 NPK fertilizer + MicroalgaeSoil properties580.231.211.0251.980.0367.21	

#### Cultivation and application of microalgae

The microalgae [Chlorella vulgaris Beyerinck (Beijerinck)] used in the present study were obtained from Cukurova University, Faculty of Fisheries and cultured in a tissue culture laboratory at Van Yuzuncu Yil University. Bold Wynne nutrient medium (NaNO3-0.250 g; MgSO<sub>4</sub>.7H<sub>2</sub>O-0.075 g; K<sub>2</sub>HPO<sub>4</sub>-0.075 g; KH<sub>2</sub>PO<sub>4</sub>-0.0175 g; NaCl-0.025 g; CaCl<sub>2</sub>.H<sub>2</sub>O-0.025 g; 1000 ml of Distilled Water) was employed in the production of bulk cultures (Duygu et al., 2017). The nutrient medium prepared in 1000 ml was equally separated into two 500 ml flasks, autoclaved at 121°C for 20 minutes for sterilization. Microalgae were cultured with 9 ml of medium +1 ml of suspended culture. The sowing of the bulk cultures was initiated with 500 ml and then transferred to 1000 ml nutrient medium. To prevent contamination and for air circulation, the mouth of the flasks was not closed, but the sterilized cotton was placed (Ağırman, 2015). After the first two days, the cultures were aired with an aquarium pump to keep the bulk cultures in suspension. The light needs of the cultures were met artificially with a light source (150 µmol m<sup>-2</sup>.s<sup>-1</sup>) placed horizontally at a distance of 22 cm from the cultures. Cultures were treated with a 16:8 light/dark period and grown at 22-25

°C. Cell counts were performed using a Thoma slide. *Chlorella vulgaris* Beyerinck (Beijerinck) algae were applied at the rate of  $2x10^7$  algae l<sup>-1</sup> to the half of the application twice (at seed sowing and 20 days after as 100 ml and 150 ml, respectively).

### The studied parameters are as follows: Shoot fresh weight (g)

The shoot weight was determined with a weighing scale with a precision of 0.1 g.

#### Shoot height (cm)

The shoot length was measured with a ruler with a precision of 1 mm.

# Total Soluble Solid Content (TSS brix<sup>o</sup>)

Garden cress samples were extracted with a blender, then TSS was determined with a hand refractometer (Atago, Tokyo, Japan).

#### Chlorophyll amount (SPAD value)

Chlorophyll content in garden cress leaves was determined with a SPAD meter (Minolta SPAD-502, Osaka, Japan).

#### **Plant Nutrient analysis**

The plant samples were put in an oven (65 °C) for constant weight. The 0.5 grams of dried and grinded samples were pre-burned with 1 ml of ethyl alcohol in crucibles and the samples were burned in a muffle furnace at 500 °C for 9-12 hours. 4 ml of 3 N HCl was mixed with the obtained ash. The samples were left on the hot plate and kept on it until they turned yellow. When the yellow color was formed, the crucibles taken from the hot plate were transferred to the volumetric flask with the help of the filtering set and the samples were made ready for reading (Kaçar and İnal, 2008).

#### Statistical analysis

IBM SPSS 21.0 package program was employed for variance analysis to compare the data obtained from the average of different doses of fertilizer applications and microalgae applications. T-test was applied for the differences between microalgae applications, and Duncan multiple range test was used for fertilizer applications (Duncan, 1955).

#### **Results and Discussion**

The effect of different doses of NPK fertilizer and microalgae [*Chlorella vulgaris* Beyerinck (Beijerinck)] application on the growth of garden cress (*Lepidium sativum* L.) was investigated in the present study. It has been determined that microalgae applications have a positive effect on plant growth and some mineral matter contents in general and some noticeable parameters are discussed below. The data about shoot fresh weight,

shoot height, Total Soluble Solid Content (TSS) and chlorophyll amount (SPAD value) are given in Table 2; the values of some nutrients (K, Ca, Fe, Zn, Cu and Mn) were also indicated in Table 3.

NPK fertilizer application had mostly higher and more significant values than the control application on plant growth parameter of garden cress (Table 2). Microalgae application had also largely higher and significant values than the control application on plant growth parameter of garden cress (Table 2). Microalgae application caused a significant increase in the shoot fresh weight and shoot height values compared to control microalgae application. Microalgae application caused significant (p≤0.01) increases (about 129% and 36%) in the shoot fresh weight and shoot height values in garden cress, respectively (Table 2). Similarly, chlorophyll amount (SPAD value) was affected and increased significantly by NPK fertilizer (p≤0.01) and microalgae  $(p \le 0.05)$  applications. Microalgae application had about 27 % more chlorophyll amount (SPAD value) than the control application. On the other hand, the increases caused by NPK fertilizer and microalgae on TSS content of garden cress were insignificant (Table 2).

Table 2. Effect of microalgae application on plant growth of garden cress grown with different doses of NPK fertilizer

	Shoot fresh weight (g pot <sup>-1</sup> )					
NPK	Microalgae		Mean	Microalgae		Mean
Fertilizer	-	+		-	+	
0% (Control)	2.21	6.13	4.17 B**	9.5	11.7	10.6 C**
50%	3.84	8.53	6.18 A	12.0	16.2	14.1 B
100%	4.39	9.23	6.81 A	13.0	18.2	15.6 A
Mean	3.48 B**	7.96 A		11.5 B**	15.3 A	

	Chlorophyll amo	nt (SPAD value) Total Soluble Solid Content (TSS				S brix <sup>0</sup> )
NPK	Microalgae		Mean	Microalgae		Mean
Fertilizer	-	+		-	+	
0% (Control)	28.65	39.02	33.83 C**	3.40	3.50	3.45 <sup>ns</sup>
50%	34.55	42.70	38.62 B	3.45	4.45	3.95
100%	35.32	43.55	39.43 A	4.10	4.52	4.31
Mean	32.80 B*	41.75 A		3.65 ns	4.15	

<sup>ns</sup>: not significant, \*: Significant at p≤0.05, \*\*: Significant at p≤0.01

In the present study, some nutrient (K, Ca, Fe, Zn, Cu and Mn) contents of garden cress grown with different doses of NPK fertilizer and microalgae application were determined and listed in Table 3. The effect of microalgae use on nutrient content was especially positive for potassium ((K), zinc (Zn), and copper (Cu). Microalgae application caused a significant ( $p\leq0.01$ ) increase (about 33%) in K content of garden cress (Table 3). Similarly, microalgae application caused a significant ( $p\leq0.01$ ) increase (about 17%) in Zn content of the plants (Table 3). Likewise, in Cu content, there was about 24% significant ( $p\leq0.01$ ) increase due

to microalgae application (Table 3). Moreover, fertilizer applications significantly increased the contents of K, Cu, and Zn. However, both applications caused no significant differences for calcium (Ca) and manganese (Mn) contents. Bio-fertilizers could possess a significant place in the development of plant growth, soil fertility and environmental factors in sustainable agriculture (Vernieri et al., 2006; Singh et al., 2016; Godlewska et al., 2019). It has been reported that microalgae application in agriculture might play an important role in the amelioration of nutrient content and plant growth (Jardin, 2015).

K (ppm)				Ca (ppm)			
NPK	Microalgae			Microalga			
Fertilizer	-	+	Mean	-	+	Mean	
0% (Control)	5098	6486	5792 C**	4244	4300	4277 <sup>ns</sup>	
50%	5873	7491	6797 B	4325	4408	4366	
100%	7407	9741	8574 A	4506	4696	4601	
Mean	6023 B **	8018 A		4358 ns	4468		
	Fe (ppm	ı)		Zn (pp	m)		
NPK	Microalgae			Microalga	ie		
Fertilizer	-	+	Mean	-	+	Mean	
0% (Control)	441.0	462.9	451.9 <sup>ns</sup>	5.56 e**	9.79 d	7.67 C*	
50%	427.3	475.7	451.5	10.47 c	10.57 c	10.52 B	
100%	449.8	489.6	469.7	12.66 b	13.26 a	12.96 A	
Mean	439.3 <sup>ns</sup>	476.0		9.56 B**	11.20 A		
	Cu (ppm	)		Mn (pp	om)		
NPK	Microalgae			Microalga	e		
fertilizer	-	+	Mean	-	+	Mean	
0% (Control)	2.89	3.30	3.12 B**	25.29	25.66	25.47 <sup>ns</sup>	
50%	3.01	3.94	3.57 AB	26.94	26.08	26.49	
100%	3.21	4.26	3.66 A	25.13	24.82	24.97	
Mean	3.06 B**	3.78 A		25.78 <sup>ns</sup>	25.52		

Table 3. Effect of microalgae application on some nutrient contents of garden cress grown with different doses of NPK fertilizer

ns: not significant, \*: Significant at p≤0.05, \*\*: Significant at p≤0.01

Microalgae, as a bio-fertilizer or bio-stimulant, could especially play important role in particular infertile soils. Organic matter produced by microalgae might also enrich the soil structure. The results of Schreiber et al. (2018) studied microalgae on wheat and confirmed that algal biomass can support crop growth on marginal soils. These researchers determined that microalgae fertilization increased the shoot size to 87 and 107% of that of sand supplied with mineral fertilizer. Michalat et al. (2015) suggested that algal extracts might induce stronger seed germination and plant growth parameters in garden cress because they are rich in nutrients and polyphenols. Michalat et al. (2016) also reported that algal extracts enhanced chlorophyll and carotenoid content in the shoots, as well as root thickness and aboveground biomass in garden cress and wheat. It was also indicated that Chlorella vulgaris as a bio-fertilizer is effective and cost-effective in improving soil nutrients for greater yield in okra (Agwa et al., 2021). Anitha et al. (2016) studied the effect of microalgae use on the nutrient content in tomato seedlings and determined that microalgae ameliorated the zinc, potassium and calcium contents significantly. Zodape et al. (2011) also examined the effect of microalgae on tomatoes and determined that nutrient contents of the microalgae applied plants improved significantly compared to the control plants.

Similarly, Grouch et al. (1990) investigated the microalgae effect on the lettuce seedlings and reported that microalgae usually improved the yield and the

contents of Ca and K. Sayed Ahmed et al. (2021) accounted that *Cholorella* might raise mineral matter content in garden rocket by upregulation of key genes in their biosynthetic pathway.

# Conclusion

Parallel to the expanding populace in the world, those require to nourishment also increases. An aftereffect of expanded pressures, the nonstop-escalated utilization of artificial chemical fertilizers might cause pollution in nature and could deteriorate the ecological balance. In the face of this situation, agricultural lands would be destroyed, and therefore, the amount and quality of agricultural crops declines. With the utilization of claiming bio-fertilizers such as microalgae, the employment of artificial chemical fertilizers could be decreased. In the present study, in which the impact microalgae [Chlorella vulgaris Beyerinck of (Beijerinck)] utilization on plant growth in garden cress (Lepidium sativum L.) was investigated, the effect of microalgae, defined as bio-fertilizer, was determined to be by and large encouraging for plant development and nutrient uptake. It was seen that there were important ameliorations of microalgae applications in many examined traits, and microalgae might compensate for the use of artificial fertilizers to some extent. In the future, it is thought that as the number of investigations on the association of microalgae and agricultural plants increases, newer knowledge will be revealed for more sustainable agriculture.

# Compliance with Ethical Standards

### Conflict of interest

The authors declared that for this research article, they have no actual, potential or perceived conflict of interest.

#### Author contribution

The contribution of the authors to the present study is equal. All the authors read and approved the final manuscript. All the authors verify that the Text, Figures, and Tables are original and that they have not been published before.

### References

Agwa, O. K., Ogugbue, C. J., & Williams, E. E. (2017). Field evidence of Chlorella vulgaris potentials as a biofertilizer for *Hibiscus esculentus*. Int. J. Agric. Res, 12(4), 181-189. DOI: 10.3923/ijar.2017.181.189.

- Ağırman, N. (2015). *Chlorella vulgaris* ve Scenedesmus acutus'un Gelişimi, Pigment Oluşumu, Lipit ve Protein İçeriği Üzerine Farklı Stres Faktörlerinin Etkileri. Fırat Üniversitesi Fen Bilimleri Enstitüsü. Elazığ. (in Turkish).
- Anitha, L, Sai Bramari, G, & Kalpana, P. (2016). Effect of supplementation of Spirulina platensis to enhance the zinc status in plants of Amaranthus gangeticus, Phaseolus aureus and tomato. Advances in Bioscience and Biotechnology 7:289–299. DOI: 10.4236/abb.2016.76027

Duncan, D. B. (1955). Multiple range and multiple F test. Biometrichs, 11: 1-42. DOI: https://doi.org/10.2307/3001478

- Duygu, Y., Udoh, A. U., Özer, T., & Erkaya, I. A. (2017). The characteristics and importance of microalgae culture collections. Süleyman Demirel Üniversitesi Eğirdir Su Ürünleri Fakültesi Dergisi, 13(1), 80-87 (in Turkish)
- Engin, Y. Ö., Yağmur, B., Cirik, S., Okur, B., Eşiyok, D., & Gökpınar, Ş. (2019). The potential use of macroalgae Ulva rigida (C. Agardh) as organic matter source in common bean production. Acta Aquatica Turcica, 15(2), 151-162. DOI: https://doi.org/10.22392/actaquatr.577506 (in Turkish)
- Eşiyok, D., Yağmur, B., & Okur, B. (2001). The effects of some naturel and mineral fertilizers on yield and mineral content of parsley (*Petroselinum crispum* Mill.). 37th Croatian symposium on agriculture with an int. participation. 19-23 February. Opatija, 179-181. DOI: https://doi.org/10.20289/zfdergi.687824
- Godlewska, K., Michalak, I., Pacyga, P., Basladynska, S., & Chojnacka, K. (2019). Potential applications of cyanobacteria: Spirulina platensis filtrates and homogenates in agriculture. World Journal of Microbiology and Biotechnology 35:80. https://doi.org/10.1007/s11274-019-2653-6

Grouch, I.J., Beckett, R.P., & Staden, J.V. (1990). Effect of Seaweed Concentrate on the Growth and Mineral Nutrition of Nutrient- Stressed Lettuce. Journal of Applied Phycology 2: 269-272. https://doi.org/10.1007/BF02179784

- IBM, SPSS. Retrieved from http://www.ibm.com/tr-tr/analytics/spss-statistics-software.
- Jardin, P. (2015). Plant biostimulants: Definition, concept, main categories and regulation. Scientia Horticulturae 196:3–14.DOI: https://doi.org/10.1016/j.scienta.2015.09.021
- Kaçar, B., & İnal, A. (2008). Plant analysis. Nobel Publication Distribution. http://www. Nobel yayın no: 1241 (in Turkish).
- Koru, E., & Cirik, S. (1999). Use of algae in agriculture and industry. Turkey 1st Ecological Agriculture Symposium, 21-23 June 1999, İzmir, 344. DOI: https://doi.org/10.24925/turjaf.v10i2.323-329.4909 (in Turkish).
- Kut, G. B., Cirik, Ş., Güroy, D., Sanver, F., & Tekinay, A. A. (2007). Effects of *Ulva rigida* or *Cystoseira barbata* meals as a feed additive on growth performance, feed utilization, and body composition in Nile tilapia, *Oreochromis niloticus*. Turkish Journal of Veterinary and Animal Science, 31(2): 91-97 (in Turkish).
- Michalak, I., Górka, B., Wieczorek, P. P., Rój, E., Lipok, J., Łęska, B., & Chojnacka, K. (2016). Supercritical fluid extraction of algae enhances levels of biologically active compounds promoting plant growth. European Journal of Phycology, 51(3), 243-252. DOI: https://doi.org/10.1080/09670262.2015.1134813
- Michalak, I., Tuhy, Ł., & Chojnacka, K. (2015). Seaweed extract by microwave assisted extraction as plant growth biostimulant. Open Chemistry, 13(1): 1183–1195. DOI: https://doi.org/10.1515/chem-2015-0132
- Midmore, D. J. (1993). Agronomic modification of resource use and intercrop productivity. Field Crops Research, 34, 357–380. DOI: https://doi.org/10.1016/0378-4290(93)90122-4
- Okur, B., Eşiyok, D., & Anaç, D. (2001). Effect of mineral and organic fertilizers on leaf nitrogen compounds of rocket (*Eruca vesicaria subsp.* sativa Mill). 37<sup>th</sup> Croatian Symposium on Agriculture. 19-23 February, Opatija-Croatia, 188-189. DOI: https://doi.org/10.20289/zfdergi.687824
- Sayed Ahmed, H. I., Elsherif, D. E., El-Shanshory, A. R., Haider, A. S., & Gaafar, R. M. (2021). Silver nanoparticles and Chlorella treatments induced glucosinolates and kaempferol key biosynthetic genes in *Eruca sativa*. Beni-Suef University Journal of Basic and Applied Sciences, 10(1), 1-15. DOI: https://doi.org/10.1186/s43088-021-00139-2
- Schreiber, C., Schiedung, H., Harrison, L., Briese, C., Ackermann, B., Kant, J., & Nedbal, L. (2018). Evaluating potential of green alga Chlorella vulgaris to accumulate phosphorus and to fertilize nutrient-poor soil substrates for crop plants. Journal of applied phycology, 30(5), 2827-2836. DOI: https://doi.org/10.1007/s10811-018-1390-9
- Sencar, Ö. (1988). The Effects of Sowing Frequency and Nitrogen in Corn Cultivation. C.Ü. Tokat Ziraat Fakültesi Yayınları No: 6, Bilimsel Araştırma ve İncelemeler No: 3, Tokat. (in Turkish).

Ethical approval Ethics committee approval is not required. Funding No financial support was received for this study. Data availability Not applicable. Consent form publication Not applicable.

- Singh, J. S., Kumar, A., Rai, A. N., & Singh, D. P. (2016). Cyanobacteria: a precious bio-resource in agriculture, ecosystem, and environmental sustainability. Frontiers in Microbiology 7:529. DOI: https://doi.org/10.3389/fmicb.2016.00529
- Tahraoui, A., El-Hilaly, J., Israili, Z. H., & Lyoussi, B. (2007). Ethnopharmacological survey of plants used in the traditional treatment of hypertension and diabetes in south-eastern Morocco (Errachidia province). Journal of Ethnopharmacology, 110(1), 105-117. DOI: https://doi.org/10.1016/j.jep.2006.09.011
- Turan, G. (2007). Use of algae in thalassotherapy. E.U. Institute of Science Department of Aquaculture, Ph.D. Thesis, 125-129. DOI: https://doi.org/10.24925/turjaf.v10i2.323-329.4909 (in Turkish).
- Vernieri, P., Borghesi, E., Tognoni, F., Serra, G., Ferrante, A., & Piagessi, A. (2006). Use of biostimulants for reducing nutrient solution concentration in floating system. In III International Symposium on Models for Plant Growth, Environmental Control and Farm Management in Protected Cultivation 718 (pp. 477-484). DOI: https://doi.org/10.17660/ActaHortic.2006.718.55
- Vural H, Eşiyok D, & Duman İ. (2000). Cultured Vegetables (Vegetable Cultivation), Ege University Faculty of Agriculture, Department of Horticulture, İzmir. https://doi.org/10.20289/zfdergi.687824 (in Turkish)
- Zodape, S.T., Gupta, A., Bhandarı, S.C., Rawat, U.S., Chaudhary, D.R., Eswaran, K., & Chıkara, J. (2011). Foliar application of seaweed sap as biositimulant for enhancement of yield and quality of tomato. Journal of Scientific & Industrial Research, Vol. 70, pp. 215- 219.