# PAPER DETAILS

TITLE: Spectroscopic Properties Determination of Flavonol by Quantum Anaysis

AUTHORS: Ibrahim ISAH NASIDI

PAGES: 36-40

ORIGINAL PDF URL: https://dergipark.org.tr/tr/download/article-file/1377560

#### Journal Physical Chemistry of and **Functional Materials**

Home Page of Journal: https://dergipark.org.tr/jphcfum

# **Specroscopic Properties Determination of Flavonol by Quantum Analysis**

Ibrahim Isah Nasidi a,\* . Mansur said<sup>c</sup> and Yau Datti<sup>b</sup>

<sup>a</sup>Department of Physics, Faculty of Science, Firat University, 23169 Elazig Turkey

<sup>b</sup> Department of Chemistry, Yusuf Maitama Sule University, Kano. Nigeria

<sup>c</sup>Department of Physics, Yusuf maitama Sule, Kano. Nigeria

## \* Corresponding author: E-mail: sinonasidi@yahoo.com

#### ABSTRACT

Computational quantum theory methods especially Hartree-Fock (HF) and Density functional theory (DFT) plays an important role in physical chemistry and theoretical physics. Flavonoids are a class of polyphenolic secondary metabolites found in plants, and thus commonly consumed in diets, that is why it is important to understand its stability, reactivity and structure analysis. In this study, we optimize the energy band gaps using different basis sets for both DFT and HF methods. LanL2MB at DFT method was selected for the molecule. IR, NMR and UV were also investigated during the research ..

#### **1. INTRODUCTION**

Flavonoids are widely distributed in plants, fulfilling many functions [1]. Flavonoids are the most important plant pigments for flower coloration, producing yellow or red/blue pigmentation in petals designed to attract pollinator animals. In higher plants, flavonoids are involved in UV filtration, symbiotic nitrogen fixation and floral pigmentation. They may also act as chemical messengers, physiological regulators, and cell cycle inhibitors. Flavonoids secreted by the root of their host plant help Rhizobia in the infection stage of their symbiotic relationship with legumes like peas, beans, clover, and soy. Rhizobia living in soil are able to sense the flavonoids and this triggers the secretion of Nod factors, which in turn are recognized by the host plant and can lead to root hair deformation and several cellular responses such as ion fluxes and the formation of a root nodule. In addition, some flavonoids have inhibitory

# **ARTICLE INFO**

Keywords: Flavonols. Band gap energy. DFT. Hartree-Fock Theory.

Received: 3-November-2020. Accepted: 20-November-2020 ISSN: 2651-3080

activity against organisms that cause plant diseases, e.g. Fusarium oxysporum [4].

Many researchers have focused on the development of a sensitive method to measurement flavonoids which is important class of natural product chemistry [12-14]. The resulting of nuclear magnetic resonance (NMR) and DFT was provided information of electrochemical properties of flavonoids and understanding its activity in aqueous solutions. DFT theoretical calculation with the basis set of LanL2MB was analyzed various hydrogen bond of flavonoids. In this study, we will explain and find the energy band gaps by using Gaussian program software. The different basis sets (3-21G, 6-31G, SDD, 6-311G, LanL2DZ, LanL2MB) for two methods (DFT and HF) performed on flavonol geometry. LanL2MB was chosen at the DFT method to show the IR and NMR spectra for the geometry structure.



# 2. COMPUTATIONAL METHODS

The flavonol structure was drawn by Chem. Bio Draw Ultra 14.0 (Fig. 1). The structure of flavonol was optimized by both density functional theory (DFT) and Hartree-Fock (HF) [9]. Different basis sets have been applied to find the energy band gaps of flavonol. The reason why different basis sets was used is that each basis sets involves different functionalities and characteristics. Table 1 shows the energy band gaps for the two methods of different basis sets. The geometry system of flavonol at the B3LYP (DFT), LanL2MB basis set has been chosen [9, 18]. The frequency and NMR were calculated to this basis set [19] The Gaussian 09 program has been used for all the calculations. Also, the origin program was performed for drawing the IR as in figure 2.



Figure 1. Structure of Flavonol

#### **3. RESULT AND DISCUSSION**

### 3.1. Energy bandgap

The first task of the computational work is to detct the optimized molecular structure using Gaussian 09 program. The energy band gaps associated with basis sets (3-21G, 6-31G, SDD, 6-311G, LanL2DZ, LanL2MB) were listed in Table 1. The energy band gaps for the Hartree-Fock (HF) method have higher values compared with the density functional theory (DFT). The optimized structural parameter of Flavonol with the LanL2MB was selected by the DFT method [9].

**Table 1.** The energy bandgap for different basis sets forboth HF and DFT methods.

	HF method	DFT method
Basis sets	Energy gaps	Energy gaps
	(ev)	(ev)
3-21G	14.9100038	9.3089228
6-31G	15.1266606	9.73979604
SDD	15.474831	10.2230032
6-311G	15.349283	10.2457806
LanL2DZ	15.4626278	10.2289687
LanL2MB	10.6438435	4.43536412

# 3.2. Molecular Structure:

The molecular structure and numbering of atom in the flavonol molecule was shown in Figure 1. The 72 vibration modes are present in the molecule. The first task of computational work was optimization of the molecule. The Gauss programs viewed the number of atoms scheme. To calculate the geometric structure DFT was used.

## 3.3. Vibrational Assignments:

The vibrational frequency was calculated by the DFT method at LanL2MB basis sets, FT-IR Figure 3 was observed with various vibration have been present in Table 2. The fundamental functional group present in the molecule vibrated at different frequencies as discussed below.

# 3.3.1. C-C /C=C vibrations:

The vibration carbon-carbon stretching in a ring usually occurs in a region 1400-1600 cm<sup>-1</sup> [21]. In aromatic benzene ring the vibration carbon-carbon was occurring in the region 1420-1625 cm<sup>-1</sup>. In the aromatic like benzene has six-member rings, there are two or more vibration are occur in a region due to the vibration of the skeletal, the strong vibration was started at 1500 cm<sup>-1</sup>. Moreover, where the ring is conjugated by the groups the further vibration observed at 1580 cm<sup>-1</sup>. In the present work, the carbon-carbon stretching vibration occurred in a region 1768.02-1839.39 cm<sup>-1</sup>.The strong starching vibration of C=C in a ring occurred in a region 1579.42-1675.25 cm<sup>-1</sup>. The vibration carbon-carbon single bond occurred in 1417.40-1488.18 cm<sup>-1</sup> it is symmetrical vibration, but the stronger vibration for C-C in benzene ring was observed in region 1488.18 cm<sup>-1</sup>. In 595.44, 649.6, 723.66 and 731.78 cm<sup>-1</sup> C-C and C=C in a plane (in a ring) was rocking but in a region 884.88, 896.45 and 964.37 cm<sup>-1</sup> was symmetrical Stretching.

#### **3.3.2.** C-H vibrations:

Commonly aromatic compounds in a region 3473-3468 cm<sup>-1</sup> were observed multiple weak bonds due to the C-H stretching vibration of the aromatic rings [14]. The C-H bending vibration in a plan was observed in a region 990-1390 cm<sup>-1</sup> and the peak was weak intensity [5]. This bond of C-H in a plane was very useful to found the characterization of compounds [26]. When the carbon and hydrogen are in a plane interaction was vibration above 1200 cm<sup>-1</sup> usually moves in the opposite direction. [7, 8]. The C-H out of the plane usually occurs in a region 700-1000 cm<sup>-1</sup> [19, 10]. In our flavonol molecule the C-H symmetrical stretching vibration out of the plane was observed between the regions 3027.75-3094.4 cm<sup>-1</sup>, but the C-H in a plane was symmetrically vibrated between the regions 3181.27-3236.82 cm<sup>-1</sup> this is an agreement with literature survey. In a region 3236.82- 526.53 cm<sup>-1</sup> all C-H

out of the plan was rocking vibrations. Also, al hydrogen was rocking vibration in flavonol molecules in a region between 1181.68- 1256.21cm<sup>-1</sup>.

Table 2.	Various	vibration	of atoms	calculated	by	DFT	(LanL2MB)
----------	---------	-----------	----------	------------	----	-----	-----------

St. NO	vibration assignments	observed frequency (cm <sup>-1</sup> )	St. NO	vibration assignments	observed frequency (cm <sup>-1</sup> )
1	(C-H) out	41.77	32	(all H) ro	1181.68
2	(C-H) out	60.72	33	(all H) ro	1200.46
3	(C-H) out	148.26	34	(all H) ro	1227.25
4	(C-H) out	181.71	35	(all H) out,ro	1256.21
5	(C-H) out	246.98	36	(C1-O8)Sy,s	1281.71
6	(O7-H17)ro	271.63	37	(C4-C12)Sy,s	1326.77
7	(N14-H15,16)ro	286.72	38	(C12-C13)Sy,s	1358.67
8	(N14-H15,16)ro (C-O)ro	307.55	39	(C-C)in,Sy	1278.63
9	(C-H)out,ro	312.8	40	(all H)out, ro	1322.75
10	(C-H) out,ro	369.13	41	(C-C)in,Sy	1393.75
11	(C-H) out,ro	428.62	42	(C-C)in,Sy	1501.59
12	(C-H) out,ro	458.24	43	(C-C)in,Sy	1567.00
13	(C-H) out,ro	473.85	44	(C13-H21,22)sc	1596.03
14	(C-H) out,ro	490.4	45	(C12-H19,20)sc	1705.81
15	(C-H) out,ro	526.53	46	(C=C)in,Sy	1746.85
16	(N14-H15,16)ro	569.12	47	(C=C)in,Sy	2048.16
17	(C-C,C=C)in,ro	595.44	48	(C=C)in,Sy	3712.96
18	(C-C,C=C)in,ro	649.6	49	(N14-H15,16)sc	3724.18
19	(C-C,C=C)in,ro	723.66	50	(C12-H19,20)sy	3027.75
20	(C-C,C=C)in,ro	731.78	51	(C13-H21,22)sy	4123.97
21	(C-H) ro	794.48	52	(C12-H19,20)asy	4261.73
22	(C-H) ro	844.21	53	(C13-H21,22)asy	4352.29
23	(C-H) ro	857.95	54	(C5-H9)sy	4463.35
24	(C12-C13)sy	884.88	55	(C3-H10)sy	3199.5
25	(C12-C13)sy	896.45	56	(C2-H11)sy	3236.82
26	(C12-C13)sy	964.37	57	(N14-H15,16)sy	3551.73
27	(C-H)in,sc	976	58	(O8-H18)sy	3626.92
28	(C12-H19,20)ro	1020.82	59	(N14- H15,16)asy	3672.03
29	(C13-N14)sy	1103.82	60		
30	(O7-H17)ro	1126.49			
31	(C13-N14)sc				

Abbreviation: out: out of the plane, ro: rocking, sy: symmetrical, asy: anti-symmetrical, in: in the plan, sc: Scissoring.

### 3.3.3. C-N vibration

In this study of flavonol structure was shown the vibration of C-N in a region 996.46 cm<sup>-1</sup> was symmetrical and in a region, 1152.66 cm<sup>-1</sup> was scissoring.

# 3.4. NMR analysis

The <sup>13</sup>C NMR chemical shifts used to identify chemical compounds [35]. GIAO (Gauge Including Atomic Orbital) is the faster procedure to calculation upon the basis set used. A GIAO method was preferable for many aspects state of the subject [19]. NMR was used to calculation chemical shift for the title compound best on TMS HF/LanL2MB(d) GIAO, TMS B3LYP/LanL2MB(2d, P)GIAO, and CH4 HF/LanL2MB(d) GIAO for carbon atoms. Hydrogen best HF/LanLMB(d) TMS GIAO, TMS on B3LYP/LanLMB(2d, P)GIAO and oxygen best on the H<sub>2</sub>O B3LYP/LanL2MB(2d, P) GIAO. H<sub>2</sub>O HF/LanL2MB(d) GIAO. Table 3 showed the 13C NMR for all references. According to TMS HF/LanL2MB(d) ordered reference. the carbon was from C3>C1>C4>C6>C5>C2>C12>C13 that means carbon number C3 was higher ppm lower filed but C13 was lower ppm higher filed. However, for TMS b3LYP/LanL2MB(2d, P) reference to carbon atoms was order from C6>C1>C3>C4 >C2>C5>C13>12 in this reference C6 was observed at higher ppm lower filed but C12 was lower ppm higher filed. For the reference, CH4 HF/6-31 G(d), C1>C6>C4>C3>C2>C5>C13>C12, carbon number one (C1) was found in higher ppm and carbon number twelve found in lower ppm higher filed.



Figure 2. FT-IR Spectrum of Flavonol

For H-NMR Table 4 used two references TMS HF/LanL2MB G(d)GIAO and TMS B3LYP/LanL2MB+G(2d, P) GIAO. According to the reference TMS HF/LanLMB (d) GIAO the observation hydrogen peak started from higher filed (H15) to the end lower of filed (H11).H11>H10>H9>H18>H22>H17>H21>H19>H20>H 16>H15. Besides for TMS reference B3LYP/LanL2MBG(2d, P) GIAO the order of hydrogen peak is the same as the previse reference with a little be different in a lower ppm for all hydrogen atoms. The NMR for nitrogen shown in Table 5, the peak was observed at 19.98 ppm for TMS HF/6-31 G(d)GIAO, but for TMS B3LYP/LanL2MB(2d, P) GIAO was observed at 17.5ppm.

**Table 3.** The observed C NMR chemical shifts in ppm for

 Flavonol

Journal of Physical Chemistry and Functional Materials

C. NO	NONE ppm	TMS HF/LanL 2MB(d) GIAO Ppm	TMS B3LYP/LanL 2MB(2d,P) GIAO ppm	CH4 HF/LanL2 MB(d) GIAO ppm
C1	54	146	151.9	145
C2	80.9	120.7	102	118.3
C3	75	155	132.7	124.1
C4	55.7	145.8	123	133.9
C5	84	124	98.3	115
C6	56	144	153.9	143.1
C12	151	71	31.9	48.1
C13	149	69	33.8	50.1

**Table 4.** The observed H NMR chemical shifts in ppm for

 Flavonol

H. NO	NONE ppm	TMS HF/LanL2MB(d) GIAO ppm	TMS B3LYP/LanL2MB(2d,P) GIAO ppm
H9	26.79	5.81	5.1
H10	26.1	6.5	5.78
H11	25.83	6.7	6.0
H15	33.2	- 0.6	- 1.31
H16	32.41	0.9	- 0.58
H17	29.85	2.75	2.05
H18	27.6	5.0	4.28
H19	30.25	2.31	1.61
H20	30.8	1.8	1.1
H21	30.05	2.55	1.58
H22	29.78	2.82	2.13

## 4. CONCLUSION

In this work, all basis sets with both HF and DTF methods investigated to calculation bandgap energy for flavonol molecule. Proposed the better basis set is LanL2MB according to the literature previous. The IR shows the vibration for all atoms that is good agreement with the kinds of literature. NMR has determined the identity of the molecular structure. The peaks of atoms in NMR were the relation with the types of references which was used on it. For example, C1 has different shifted in a ppm for all three references. The reference H2O HF/LanL2MB(d)GIAO and H2O B3LYP/LanL2MB(2d, P)GIAO show the oxygen's ppm and it is closed to each other.

# Nasidi et al.

# REFERENCES

- [1] Cermak R, Wolffram S (October 2006). "The potential of flavonoids to influence drug metabolism and pharmacokinetics by local gastrointestinal mechanisms". Curr. Drug Metab. 7 (7): 729–44. doi:10.2174/138920006778520570.
  PMID 17073577. Archived from the original on 2012-07-20.
- [2] Smith, Gerald J.; Markham, Kenneth R. (1998).
  "Tautomerism of flavonol glucosides: relevance to plant UV protection and flower colour". Journal of Photochemistry and Photobiology A: Chemistry. 118 (2): 99–105. doi:10.1016/s1010-6030(98)00354-2.
- [3] "Juicy news about cranberries". medicalxpress.com. Retrieved 13 April 2018.
- [4] Nakabayashi R1, Yonekura-Sakakibara K, Urano K, Suzuki M, Yamada Y, Nishizawa T, Matsuda F, Kojima M, Sakakibara H, Shinozaki K, Michael AJ, Tohge T, Yamazaki M, Saito K. (2014).
  "Enhancement of oxidative and drought tolerance in Arabidopsis by overaccumulation of antioxidant flavonoids". Plant J. 77 (3): 367–79. doi:10.1111/tpj.12388. PMC 4282528. PMID 24274116.
- [5] Si D, Wang Y, Zhou YH, et al. (March 2009).
  "Mechanism of CYP2C9 inhibition by flavones and flavonols". Drug Metab. Dispos. 37 (3): 629–34. doi:10.1124/dmd.108.023416. PMID 19074529.
- [6] McNaught, Alan D; Wilkinson, Andrew; IUPAC (1997), IUPAC Compendium of Chemical Terminology (2 ed.), Oxford: Blackwell Scientific, doi:10.1351/goldbook.F02424, ISBN 978-0-9678550-9-7
- [7] Nič, Miloslav; Jirát, Jiří; Košata, Bedřich; Jenkins, Aubrey; McNaught, Alan, eds. (2009). "Flavonoids (isoflavonoids and neoflavonoids)". The Gold Book. doi:10.1351/goldbook. ISBN 978-0-9678550-9-7. Retrieved 16 September 2012.
- [8] Galeotti, F; Barile, E; Curir, P; Dolci, M; Lanzotti, V (2008). "Flavonoids from carnation (Dianthus caryophyllus) and their antifungal activity". Phytochemistry Letters. 1: 44–48. doi:10.1016/j.phytol.2007.10.001.
- [9] Ververidis F, Trantas E, Douglas C, Vollmer G, Kretzschmar G, Panopoulos N (October 2007).
   "Biotechnology of flavonoids and other phenylpropanoid-derived natural products. Part I:

Chemical diversity, impacts on plant biology and human health". Biotechnology Journal. 2 (10): 1214– 34. doi:10.1002/biot.200700084. PMID 17935117.

- [10] Isolation of a UDP-glucose: Flavonoid 5-Oglucosyltransferase gene and expression analysis of anthocyanin biosynthetic genes in herbaceous peony (Paeonia lactiflora Pall.). Da Qiu Zhao, Chen Xia Han, Jin Tao Ge and Jun Tao, *Electronic Journal of Biotechnology*, 15 November 2012, Volume 15, Number 6, doi:10.2225/vol15-issue6-fulltext-7
- [11] Spencer JP (2008). "Flavonoids: modulators of brain function?". British Journal of Nutrition. 99 (E-S1): ES60–77. doi:10.1017/S0007114508965776. PMID 18503736.
- [12] USDA's Database on the Flavonoid Content
- [13] Oomah, B. Dave; Mazza, Giuseppe (1996).
  "Flavonoids and Antioxidative Activities in Buckwheat". Journal of Agricultural and Food Chemistry. 44 (7): 1746–1750. doi:10.1021/jf9508357.
- [14] Ayoub M, de Camargo AC, Shahidi F (2016).
  "Antioxidants and bioactivities of free, esterified and insoluble-bound phenolics from berry seed meals".
  Food Chemistry. 197 (Part A): 221–232. doi:10.1016/j.foodchem.2015.10.107.
  PMID 26616944.
- [15] [1] p. 223 Table 1
- [16] The Lancet (2007). "The devil in the dark chocolate". Lancet. 370 (9605): 2070. doi:10.1016/S0140-6736(07)61873-X. PMID 18156011.
- [17] Serafini M, Bugianesi R, Maiani G, Valtuena S, De Santis S, Crozier A (2003). "Plasma antioxidants from chocolate" (PDF). Nature. 424 (6952): 1013. Bibcode:2003Natur.424.1013S. doi:10.1038/4241013a. PMID 12944955.
- [18] Serafini M, Bugianesi R, Maiani G, Valtuena S, De Santis S, Crozier A (2003). "Nutrition: milk and absorption of dietary flavanols" (PDF). Nature. 424 (6952): 1013. Bibcode:2003Natur.424.1013S. doi:10.1038/4241013a. PMID 12944955.
- [19] Roura E, et al. (2007). "Milk Does Not Affect the Bioavailability of Cocoa Powder Flavonoid in Healthy Human" (PDF). Ann Nutr Metab. 51 (6): 493–498. doi:10.1159/000111473. PMID 18032884. [permanent dead link]