


PAPER DETAILS

TITLE: Weakly Prime Radical of Submodules

AUTHORS: Sibel Cansu

PAGES: 105-108

ORIGINAL PDF URL: <https://dergipark.org.tr/tr/download/article-file/3129887>



Weakly Prime Radical of Submodules

Sibel Cansu¹

¹Department of Mathematics, Faculty of Science and Arts, Bolu Abant İzzet Baysal University, Bolu, Türkiye

Abstract

In this paper, some properties of weakly prime radical are stated. The characterization of weakly prime radical for finitely generated modules is given. Also, the relationship between the weakly prime radical of a submodule and the ideals of the ring T is considered.

Keywords: Prime radical; Prime submodule; weakly prime radical; weakly prime submodule.

2010 Mathematics Subject Classification: 13C99, 13A99

1. Introduction

All rings will be commutative and all modules will be unitary throughout this work. Let T be a ring and H be a T -module and K be a proper submodule. Then K is called prime if $rm \in K$ implies $m \in K$ or $rH \subseteq K$ where $r \in T, m \in H$. Also K is called weakly prime submodule whenever $rs m \in K$ either $rm \in K$ or $sm \in K$ for some $r, s \in T, m \in H$. Weakly prime submodules were introduced in [4]. It can be easily shown that every prime submodule is weakly prime. But not every weakly prime submodule is prime as the following example shows [4].

Let $T = \mathbb{Q}[x, y]$, $P = \langle x \rangle$ be a non-zero prime ideal of T , H be a free T -module $T \oplus T$. Then $Q = 0 \oplus P$ is a weakly prime submodule of H but it is not a prime submodule.

The concept of the radical of an ideal was generalized to modules over commutative rings [5]. As a result of this generalization, the definition of a prime submodule has shown up. If K is a proper submodule of a T -module H , then the prime radical of K , $rad_H(K)$, is the intersection of all prime submodules containing K . If there is no prime submodule contains K , then $rad_H(K) = H$. In 2006, Behboodi gave the definition of weakly prime radical of submodule [3]. The weakly prime radical of K in H , $wrad_H(K)$, is the intersection of all weakly prime submodules of H containing K . If there is no weakly prime submodule containing K , then $wrad_H(K) = H$. By the definitions, $wrad_H(K) \subseteq rad_H(K)$.

In this work, we dealt with the weakly prime radical of submodules, its properties, and its relationship with the ideals of the ring T .

2. Weakly Prime Submodules

It is well-known that if K is a prime submodule of H , then $(K : H) = \{r \in T : rH \subseteq K\}$ is a prime ideal. If K is weakly prime, then $(K : H)$ is a radical ideal [6]. Also, the following can easily be shown if I is an ideal of T .

Proposition 2.1. Let W be any weakly prime submodule of H and I be an ideal of T . Then $(W : I) = \{m \in H : Im \subseteq W\}$ is a weakly prime submodule of H .

Proof. Let $r, s \in T$ and $m \in H$ such that $rs m \in (W : I)$. Then $(ars)m \in W$ for every $a \in I$. Since W is weakly prime submodule, $(ar)m \in W$ or $sm \in W$. If $(ar)m \in W$, then either $am \in W$ or $rm \in W$. $am \in W$ implies that $m \in (W : I)$. Since $W \subseteq (W : I)$, $sm \in W$ or $rm \in W$ also gives that $sm \in (W : I)$ or $rm \in (W : I)$. Hence in each cases, either $rm \in (W : I)$ or $sm \in (W : I)$. \square

The radical of an ideal I of a ring is known as the intersection of all minimal prime ideals of I . We can consider a similar characterization for the weakly prime radical of a submodule.

Definition 2.2. A weakly prime submodule W of H is called a minimal weakly prime submodule of a submodule K if $K \subseteq W$ and there is no weakly prime submodule of H containing K which is smaller than W . A minimal weakly prime submodule of $\langle 0 \rangle$ is minimal weakly prime submodule of H .

Proposition 2.3. Let $\{W_i : i \in I\}$ be a non-empty family of weakly prime submodules of a T -module H . If the family is totally ordered by inclusion, then $\bigcap_{i \in I} W_i$ is a weakly prime submodule of H .

Proof. Let $xym \in \cap_{i \in I} W_i$ for $x, y \in T$ and $m \in H$. Then $xym \in W_i$ for all $i \in I$. Since W_i is weakly prime, $xm \in W_i$ or $ym \in W_i$. Since the family is totally ordered, $xm \in \cap_{i \in I} W_i$ or $ym \in \cap_{i \in I} W_i$. Thus $\cap_{i \in I} W_i$ is a weakly prime submodule. \square

Proposition 2.4. *If a submodule K of a T -module H is contained in a weakly prime submodule N , then N contains a minimal weakly prime submodule of K .*

Proof. Let $M = \{L : L \text{ is weakly prime submodule of } H \text{ and } K \subseteq L \subseteq N\}$. It is clear that M is non-empty. If $L_1, L_2 \in M$, define a relation \leq on M as $L_1 \leq L_2$ if $L_2 \subseteq L_1$. It is a partial order on M . For any non-empty totally ordered subset M_1 of M , if $\bar{L} = \cap_{L_i \in M_1} L_i$, then \bar{L} is a weakly prime submodule since M_1 is totally ordered and also by the above proposition. Then $K \subseteq \bar{L} \subseteq N$. Hence $\bar{L} \in M$ and thus it is an upper bound for M_1 . M contains a maximal element Y by Zorn's Lemma. Since $Y \in M$, it is a weakly prime submodule of H .

To complete the proof it is enough to show that Y is a minimal prime submodule of K . Suppose that \bar{Y} is a weakly prime submodule of H where $K \subseteq \bar{Y} \subseteq N$ and $\bar{Y} \subsetneq Y$. Then $\bar{Y} \in M$ and hence $Y \leq \bar{Y}$. Thus $Y = \bar{Y}$. Therefore Y is a minimal weakly prime submodule of K . \square

Corollary 2.5. *Every proper submodule of a finitely generated module possesses at least one minimal weakly prime submodule.*

Proof. Let H be a finitely generated module, and Q be a proper submodule of H . Then there exists a submodule K of H such that $Q \subseteq K$ and K is maximal. Since K is maximal, it is a prime submodule [7]. Hence K is a weakly prime submodule of H . Then by Proposition 2.4, K will contain a minimal weakly prime submodule of Q . Thus Q has at least one minimal weakly prime submodule. \square

If H is a finitely generated module, then we can redefine the weakly prime radical of submodules of H by the above corollary.

Theorem 2.6. *Let H be a finitely generated T -module and K be a proper submodule of H . Then weakly prime radical of K is the intersection of its minimal weakly prime submodules.*

Proof. Let K be a proper submodule of H . By Corollary 2.5, K has at least one minimal weakly prime submodule, say W_i . Let L be the intersection of all minimal weakly prime submodules of H containing K . By the definition of weakly prime radical, $\text{wrad}_H(K) \subseteq \cap_{i \in I} W_i = L$. On the other hand, if W is any weakly prime submodule containing K , then W contains some minimal weakly prime submodule Q_i of K by Proposition 2.4. Hence $L = \cap_{i \in I} W_i \subseteq \text{wrad}_H(K)$. \square

Some properties of weakly prime radical is given in the following proposition.

Proposition 2.7. *Let H be a T -module, J be an index set and let N, N_j be submodules of H for $j \in J$ and I be an ideal of T . Then*

- (i) $N \subseteq \text{wrad}_H(N)$,
- (ii) $\text{wrad}_H(\text{wrad}_H(N)) = \text{wrad}_H(N)$,
- (iii) $\text{wrad}_H(\bigcap_{j \in J} N_j) \subseteq \bigcap_{j \in J} \text{wrad}_H(N_j) = \text{wrad}_H(\bigcap_{j \in J} \text{wrad}_H(N_j))$,
- (iv) $\sum_{j \in J} \text{wrad}_H(N_j) \subseteq \text{wrad}_H(\sum_{j \in J} N_j) = \text{wrad}_H(\sum_{j \in J} \text{wrad}_H(N_j))$,
- (v) $\text{wrad}_H(IH) = \text{wrad}_H(\sqrt{I}H) = \text{wrad}_H(I^n H)$ for every positive integer n ,
- (vi) $\sqrt{(N : H)} \subseteq (\text{wrad}_H(N) : H)$

Proof. (i) If $x \in N$, then $x \in W$ for every weakly prime submodule of H containing N . Thus $x \in \text{wrad}_H(N)$.

(ii) Since $N \subseteq \text{wrad}_H(N) \subseteq W_k$ for any weakly prime submodule W_k containing $\text{wrad}_H(N)$, $\bigcap_{k \in K} W_k \subseteq P_j$ for all weakly prime submodules P_j containing N . Hence $\text{wrad}_H(\text{wrad}_H(N)) = \bigcap_{k \in K} W_k \subseteq \text{wrad}_H(N)$. The other side is clear by part (i).

(iii) Let $\text{wrad}_H(\bigcap_{j \in J} N_j) = \bigcap_{k \in K} W_k$ where W_k is weakly prime submodule containing $\bigcap_{j \in J} N_j$ for every $k \in K$, and let $\{Q_{kj}\}$ be the set of weakly prime submodules containing N_j . Since $\bigcap_{j \in J} N_j \subseteq N_j \subseteq Q_{kj}$ for all k and j , $\text{wrad}_H(\bigcap_{j \in J} N_j) \subseteq \text{wrad}_H(N_j)$.

Since $\bigcap_{j \in J} \text{wrad}_H(N_j) \subseteq \text{wrad}_H(N_j)$ for all $j \in J$,

$$\text{wrad}_H(\bigcap_{j \in J} \text{wrad}_H(N_j)) \subseteq \text{wrad}_H(\text{wrad}_H(N_j)) = \text{wrad}_H(N_j) \text{ for all } j \in J.$$

Hence $\text{wrad}_H(\bigcap_{j \in J} \text{wrad}_H(N_j)) \subseteq \bigcap_{j \in J} \text{wrad}_H(N_j)$. By part (i), $\bigcap_{j \in J} \text{wrad}_H(N_j) = \text{wrad}_H(\bigcap_{j \in J} \text{wrad}_H(N_j))$ is clear.

(iv) Since $N_j \subseteq \sum_{j \in J} N_j$ for all j , $\text{wrad}_H(N_j) \subseteq \text{wrad}_H(\sum_{j \in J} N_j)$. Then $\sum_{j \in J} \text{wrad}_H(N_j) \subseteq \text{wrad}_H(\sum_{j \in J} N_j)$. Since $\sum_{j \in J} N_j \subseteq \sum_{j \in J} \text{wrad}_H(N_j)$, it can easily be shown that $\text{wrad}_H(\sum_{j \in J} N_j) = \text{wrad}_H(\sum_{j \in J} \text{wrad}_H(N_j))$.

(v) This is trivially true if $\text{wrad}_H(IH) = H$. If $\text{wrad}_H(IH) \neq H$, then there exists a weakly prime submodule W of H where $IH \subseteq W$. Since $I \subseteq (IH : H) \subseteq (W : H)$, $\sqrt{I} \subseteq \sqrt{(W : H)} = (W : H)$. So that $\sqrt{I}H \subseteq (W : H)H \subseteq W$. Thus $\text{wrad}_H(\sqrt{I}H) \subseteq \text{wrad}_H(IH)$. The other side is clear since $I \subseteq \sqrt{I}$ is always true.

Since $\sqrt{I^n} = \sqrt{I}$ for any positive integer n ,

$$\text{wrad}_H(I^n H) = \text{wrad}_H(\sqrt{I^n} H) = \text{wrad}_H(\sqrt{I} H) = \text{wrad}_H(IH).$$

(vi) Let $0 \neq a \in \sqrt{(N : H)}$. Then there exists $k \in \mathbb{Z}^+$ such that $a^k H \subseteq N \subseteq W$ for every weakly prime submodules W which contains N . Therefore $a \in (W : H)$ since $(W : H)$ is a prime ideal. So $aH \subseteq \text{wrad}_H(N)$ and thus $a \in (\text{wrad}_H(N) : H)$. \square

Lemma 2.8. If H is finitely generated T -module and N is a submodule of H , then $\text{wrad}_H(N) = H$ if and only if $N = H$.

Proof. Assume that $\text{wrad}_H(N) = H$ and $N \neq H$. By Corollary 2.5, N has at least one minimal weakly prime submodule W . Hence $H = \text{wrad}_H(N) \subseteq W$. Since W is weakly prime, $N = H$. The other side is clear if $N = H$. \square

Corollary 2.9. Let H be a finitely generated module, K , and L be submodules of H . Then $\text{wrad}_H(K) + \text{wrad}_H(L) = H$ if and only if $K + L = H$.

Proof. Assume that $\text{wrad}_H(K) + \text{wrad}_H(L) = H$. Then by Lemma 2.8 and Proposition 2.7, $\text{wrad}_H(K + L) = H$ and hence $K + L = H$. Conversely if $K + L = H$, then $H = \text{wrad}_H(K + L) = \text{wrad}_H(\text{wrad}_H(K) + \text{wrad}_H(L))$. Hence $\text{wrad}_H(K) + \text{wrad}_H(L) = H$ by Lemma 2.8. \square

Proposition 2.10. If K is a proper submodule of a T -module H , then $\text{wrad}_H(K) = \text{wrad}_H(K + pH)$ where $p = \sqrt{(K : H)}$ is a prime ideal.

Proof. It is clear that $\text{wrad}_H(K) \subseteq \text{wrad}_H(K + pH)$, since $K \subseteq K + pH$. Assume $\text{wrad}_H(K) = \bigcap_{i \in I} W_i$ where W_i is weakly prime submodule of H containing K . Since $(W_i : H)$ is a prime ideal, $p \subseteq (W_i : H)$ which implies that $K + pH \subseteq W_i$. So, $\text{wrad}_H(K + pH) \subseteq \bigcap_{i \in I} W_i = \text{wrad}_H(K)$. \square

Corollary 2.11. Let H be a finitely generated module, K be a proper submodule of H , and $p = (K : H)$ be a maximal ideal of T . Then $\text{wrad}_H(K)$ is weakly prime submodule and $\text{wrad}_H(K) = K + pH$.

Proof. $K + pH \subseteq \text{wrad}_H(K + pH) = \text{wrad}_H(K)$ by Proposition 2.10. Since $p = (K : H) \subseteq ((K + pH) : H)$, $((K + pH) : H) = p$ or $((K + pH) : H) = T$. If $((K + pH) : H) = T$, then $TH = H \subseteq K + pH \subseteq \text{wrad}_H(K)$ which implies that $H = \text{wrad}_H(K)$. Since H is finitely generated, $H = K$ by Lemma 2.8. So $((K + pH) : H) = p$. Hence $K + pH$ is a prime submodule by [7] and thus it is a weakly prime submodule containing K . Therefore $\text{wrad}_H(K) = K + pH$. \square

For the rest of this section, the relationship between the weakly prime radical of a submodule and the ideals of the ring T will be examined.

Lemma 2.12. Let N be a proper submodule of a T -module H and I be an ideal of T . Then $\text{wrad}_H(\text{wrad}_H(N) : I) = (\text{wrad}_H(N) : I)$.

Proof. Assume that $\text{wrad}_H(N) = \bigcap_{j \in J} W_j$ for all weakly prime submodules W_j of H containing N . Then

$$\text{wrad}_H(\text{wrad}_H(N) : I) = \text{wrad}_H(\bigcap_{j \in J} W_j : I) = \text{wrad}_H(\bigcap_{j \in J} (W_j : I)) \subseteq \bigcap_{j \in J} \text{wrad}_H(W_j : I)$$

by Proposition 2.7. Also Proposition 2.1 implies that

$$\bigcap_{j \in J} \text{wrad}_H(W_j : I) = \bigcap_{j \in J} (W_j : I) \subseteq (W_j : I)$$

for all weakly prime submodules W_j of H containing N . So,

$$I(\text{wrad}_H(\text{wrad}_H(N) : I)) \subseteq I(W_j : I) \subseteq W_j$$

and it implies that $I(\text{wrad}_H(\text{wrad}_H(N) : I)) \subseteq \text{wrad}_H(N)$. Hence $\text{wrad}_H(\text{wrad}_H(N) : I) \subseteq (\text{wrad}_H(N) : I)$. The other side is clear by Proposition 2.7. \square

Corollary 2.13. Let N be a proper submodule of a T -module H and I be an ideal of T . Then $\text{wrad}_H(N : I) = (\text{wrad}_H(N) : I)$.

Proof. Since $N \subseteq \text{wrad}_H(N)$, $(N : I) \subseteq (\text{wrad}_H(N) : I)$ which means that $\text{wrad}_H(N : I) \subseteq \text{wrad}_H(\text{wrad}_H(N) : I) = (\text{wrad}_H(N) : I)$ by Lemma 2.12. \square

Proposition 2.14. Let N be a proper submodule of a T -module H , I and J be ideals of T . Then

$$\text{wrad}_H(IJN) = \text{wrad}_H(IN) \cap \text{wrad}_H(JN).$$

Proof. Consider the sets $X_1 = \{W : IJN \subseteq W\}$, $X_2 = \{W' : IN \subseteq W'\}$, and $X_3 = \{\bar{W} : JN \subseteq \bar{W}\}$ where W, W' and \bar{W} are weakly prime submodules of H . Since each $W \in X_1$ is weakly prime, $IJN \subseteq W$ implies $IN \subseteq W$ or $IJ \subseteq W$. Hence $X_1 = X_2 \cup X_3$ and

$$\text{wrad}_H(IJN) = \bigcap_{W \in X_1} W = (\bigcap_{W' \in X_2} W') \cap (\bigcap_{\bar{W} \in X_3} \bar{W}).$$

Therefore $\text{wrad}_H(IJN) = \text{wrad}_H(IN) \cap \text{wrad}_H(JN)$. \square

Article Information

Acknowledgements: The authors would like to express their sincere thanks to the editor and the anonymous reviewers for their helpful comments and suggestions.

Author's contributions: All authors contributed equally to the writing of this paper. All authors read and approved the final manuscript.

Conflict of Interest Disclosure: No potential conflict of interest was declared by the author.

Copyright Statement: Authors own the copyright of their work published in the journal and their work is published under the CC BY-NC 4.0 license.

Supporting/Supporting Organizations: No grants were received from any public, private or non-profit organizations for this research.

Ethical Approval and Participant Consent: It is declared that during the preparation process of this study, scientific and ethical principles were followed and all the studies benefited from are stated in the bibliography.

Plagiarism Statement: This article was scanned by the plagiarism program. No plagiarism detected.

Availability of data and materials: Not applicable.

References

- [1] A. Azizi, Radical formula and prime submodules, *Journal of Algebra* Vol:307, (2007), 454-460.
- [2] A. Azizi, Radical formula and weakly prime submodules, *Glasgow Mathematical Journal of Trust* Vol:51, (2009), 405-412.
- [3] M. Behboodi, On weakly prime radical of modules and semi-compatible modules, *Acta Math. Hungar.* Vol:113, No:3, (2006), 243-254.
- [4] M. Behboodi and H. Koohy, Weakly prime modules, *Vietnam Journal of Mathematics* Vol:32, No:2 (2004), 185-195.
- [5] R.L. McCasland and M.E. Moore, On radical of submodules, *Communications in Algebra* Vol:19, No:5, (1991), 1327-1341.
- [6] E. Yılmaz and S. K. Cansu, Baer's lower nilradical and classical prime submodules, *Bulletin of the Iranian Mathematical Society* Vol:40, No:5, (2014), 1263-1274.
- [7] C-P. Lu, Prime submodules of modules, *Comment. Math. University Sancti Pauli* Vol:33, No:1, (1984), 61-69.