

PAPER DETAILS

TITLE: Integrated Weed Management in Canola (*Brassica napus* L.)

AUTHORS: Ghodratollah Fathi

PAGES: 57-63

ORIGINAL PDF URL: <https://dergipark.org.tr/tr/download/article-file/158832>

INTEGRATED WEED MANAGEMENT IN CANOLA

(*Brassica napus L.*)

Ghodratollah Fathi

Associate Professor, Department of Agronomy, Agricultural Sciences and Technology of
Khoozestan University, Ahwaz, IRAN

ABSTRACT

A field experiment was conducted at Agricultural Sciences and Technology of Khoozestan University, Ahwaz in South of Iran during 2001-2 and 2002-3 to find out most effective control measure for weeds in canola (*Brassica napu L.*). Terflan (2.5 kg/ha), terflan+ one cultivation, terflan+ two cultivations, terflan+ nabo-s (2 L/ha), terflan+ nabo-s +one cultivation, terflan+ nabo-s + two cultivation, nabo-s + two cultivations, nabo-s+one cultivation, nabo-s, one cultivation, two cultivations were compared with weed-free and weedy check treatments. Treatments were replicated 4 times and were compared in a complete randomized block design. *Malva sylvestris*, *Convolvulus arvensis* and *Atriplex patulum* were the predominant weeds. Combining pre and post emergence herbicides with two cultivations effectively controlled weeds. Pre-emergence herbicide also provided desired control of weeds. With treatments dry weight and number of weeds was sharply reduced particularly *C. arvensis* and *A. patulum*. Inclusion of post-emergence weed-control measures with pre-emergence herbicide markedly improved WCE, yield attributes and grain yield. However, cultivations of weed proved most effective, followed by hand-weeding. The highest grain yield (3015.6 kg/ha), superior yield attributes, were recorded under terflan+ nabo-s+two cultivations. It can be concluded that cultivations was completed the effects of herbicides.

Keywords: Canola, integrated control, chemical control, mechanical control, weeds