

PAPER DETAILS

TITLE: EFFECT OF RADIATION ON VIABILITY AND INFECTIVITY OF NEMATODIRUS SP.
EGGS

AUTHORS: Harun ÇERÇİ, Metin ALABAY, Zisan EMRE, Ali DÜZGÜN

PAGES: 27-33

ORIGINAL PDF URL: <https://dergipark.org.tr/tr/download/article-file/456411>

EFFECT OR RADIATION ON VIABILITY AND INFECTIVITY OF NEMATODIRUS SP. EGGS

H. ÇERÇİ, M. ALABAY, Z. EMRE, A. DÜZGÜN

Turkish Atomic Energy Authority, Ankara Nuclear Research Center for Agriculture and Animal Sciences, Saray – Kazan, 06983, Ankara-Turkey

ABSTRACT

In this study, effects of different levels of gamma irradiation on the hatching ability of *Nematodirus* sp. Eggs and infectivity of irradiated larvae in Angora goats were investigated. *Nematodirus* sp. Eggs were exposed to radiation doses ranging from 1 to 300 Gy. It was found that there was no significant differences with respect to hatching ability between the control and irradiated groups of eggs in low doses such as 1, 5, 10 and 20 Gy when larval development in eggs were examined. The results show that there is an inverse relationship between the gamma radiation dose used and the hatching ability of eggs after the dose of 30 Gy. The more the radiation dose level, the lesser the hatching was observed and there was no hatchings after the dose of 225 Gy. Moreover, degenerative changes in the developments and multiplications of blastomeres relating to increases of radiation doses were observed. The differences between the control group and irradiated groups (starting from 30 Gy) were found to be statistically significant ($P<0.05$). Twelve Angora goats were infected with each receiving 500 *Nematodirus* sp. Larvae irradiated with doses of 0, 30, 50 and 100 Gy. All animals were sacrificed and worm counts were performed 28 days after infection. It was determined that there were decreases in the egg and worm counts and delayed egg sheddings with the increases of irradiation doses and no worms were obtained at a dose of 100 Gy. Also, the survival of more females than males in the collected parasites was found.

Keywords: Radiation, *Nematodirus* sp., Viability, Infectivity

ÖZET

Bu çalışmada, değişik dozlarda ışınlamanın *Nematodirus* sp. Yumurtalarının açılmasına ve ışınlamanın larvaların enfektivitelerine etkileri ankara keçilerinde araştırılmıştır. *Nematodirus* sp. Yumurtaları 1-300 Gy arasında değişen dozlarda ışınlanmıştır. Yumurtalardaki larval gelişme incelendiğinde, 1, 5, 10 ve 20 Gy gibi düşük dozlarda ışınlanan gruplarla kontrol grubu arasında yumurtaların açılmasına ilişkin önemli farklar bulunmamıştır. Otuz Gy'lik ışınlamadan sonra, yumurtaların açılma yeteneği ve kullanılan gama radyasyon dozu arasında ters bir ilişki olduğu görülmüştür. Radyasyon dozu arttıkça, daha az sayıda açılma gözlenmiş ve 225 Gy'lik dozdan sonra açılma olmamıştır. Bundan başka, radyasyon dozunun artışına bağlı olarak blastomerlerin çoğalmasında ve gelişmesinde dejeneratif değişiklikler gözlenmiştir. Kontrol ve ışınlama grupları arasındaki fark (30 Gy'den başlayarak) istatistiksel olarak önemli bulunmuştur. Oniki Ankara keçisi, 0, 30, 50 ve 100 Gy dozda ışınlanmış 500'er *Nematodirus* sp. Larvası ile enfekte edilmişlerdir. Enfeksiyondan 28 gün sonra tüm hayvanlar kesilmiş ve parazit sayımları yapılmıştır. Radyasyon dozunun artışına bağlı olarak yumurta ve parazit sayımlarında bir azalma olduğu belirlenmiş ve 100 Gy dozda ise hiçbir parazit elde edilememiştir. Ayrıca, diğer gruplarda da toplanan parazitlerin çoğunun dişi olduğu görülmüştür.

Anahtar kelimeler: Radyasyon, Gama ışınlaması, *Nematodirus*, Canlılık, Enfektivite

INTRODUCTION

Nematodirus sp. are the parasites of ruminants and sited in small intestine. Its distribution is almost world wide but commoner in temperate regions. There are four species of *nematodirus* named *N. Filicollis*, *N. Spathiger*, *N. Battus*, *N. Helvetianus*. The first three species are commoner in sheep and goats and the last one is the species of cattle and is less commonly found in sheep, goat and camel. The adults of *Nematodirus* sp. Are slender worms about 2 cm. Long. *Nematodiriasis* is an example of a parasitic disease where the principal pathogenic effect is attributable to the larval stages. Following ingestion of large numbers of infective larvae, there is disruption of the intestinal mucosa particularly in the ileum, although the majority of developing stages are found on the mucosal surface.

Development through L4 to L5 is complete by 10-12 days from infection and this coincides with severe damage to the intestinal villi and erosion of the mucosa leading to villous atrophy. The ability of the intestine to exchange fluids and nutrients is grossly reduced and with the onset of diarrhoea the lamb rapidly becomes dehydrated. At necropsy the carcass has a dehydrated appearance and there is an enteritis in the ileum. In severe infections, diarrhoea is the most prominent clinical sign. As dehydration proceeds the affected animals become thirsty, inappetent and congregate round drinking places.

Nematodirus sp. Infections are wide spread all over Turkey, especially young lambs are more affected by the parasites. Its prevalence in sheep and goats was studied by a number of workers and following figures have been reported; 32% in İstanbul, 67.4-91.3% in Samsun, 41-78% in Ankara (2, 3, 4, 5, 6).

MATERIALS AND METHODS

In this study, effect of different levels of gamma irradiation on the hatching ability of *Nematodirus sp.* Eggs and infectivity of irradiated larvae in angora goats were investigated. For this purpose fresh faecal samples were collected from the villages of Ankara province. These samples were processed at the laboratory by floatation technique using saturated NaCl solutions and reasonably clean egg suspensions were obtained. Then, *Nematodirus sp.* eggs were picked up one by one from these suspensions under a stereo microscope. The collected *Nematodirus* eggs had been incubated at 25°C until the larvae hatched. A total of 3000 larvae were administered to a goat and were placed in a metabolism cage to collect faeces. This monospecifically infected goat produced enough egg materials for irradiation studies. The collection and cleaning methods of faecal samples were as reported by Alabay et al. (1).

To study the effect of different levels of gamma irradiation on the hatching ability of *Nematodirus sp.* eggs; egg materials were exposed to radiation doses ranging from 1-300 Gy by a ^{137}Cs source of 10.000 Ci (Mark I-22 irradiator) with a dose rate of 1.81 kGy/h as determined by Fricke dosimetry. To study the infectivity of irradiated larvae; goats were into four groups. First three groups were given irradiated larvae and the forth one left

as control given nonirradiated larvae. Irradiation doses were chosen according to vitality percentages of larvae. Those were 75 % (30 Gy), 50% (50 Gy) and 25 % (100 Gy). Experimental animals including controls were infected with 500 *Nematodirus* larvae each. Animals were examined for infections by McMaster egg counting technique starting from day 16 th. and necropsied on day 28 th post infection. At the necropsy, small intestines were removed and examined for the presence of parasites.

RESULTS AND DISCUSSION

It was found that there was not significant differences with respect to hatching ability between the control and irradiated groups of eggs in low doses such as 1, 5, 10 and 20 Gy when larval development in eggs were examined. The results show that there is an inverse relationship between the gamma irradiation dose used and the hatching ability of eggs after the dose of 30 Gy. The more the radiation dose level, the lesser the hatching of eggs after the dose of 225 Gy (Table 1). Moreover, degenerative changes in the developments and multiplications of blastomers relating to increases of radiation doses were observed. The differences between the control groups and the irradiated groups (starting from 30 Gy) were found to be statistically significant ($P<0.05$). Twelve Angora goats were divided into four groups with 3 animals in each and infected with *Nematodirus sp.* larvae. Each animal was given 500 *Nematodirus sp.* larvae irradiated with doses of 0, 30, 50 and 100 Gy. All animals were sacrificed and worm counts were performed 28 days after infection. It was determined that there were decreases in the egg and worm counts and delayed egg sheddings with the increases of irradiation doses and no worms were obtained at a dose of 100 Gy. Also, the survival of more females than males in the collected parasites was found (Table 2). Although there are a number of reports on the effects of radiation on different parasite species, no studies were found on *Nematodirus sp.*

Table 1. Numbers of hatching larvae from irradiated eggs.

Irradiation Doses	Number of larvae	%
Control	431	86.2
1	426	85.2
5	428	85.6
10	419	83.8
20	401	80.2
30	367	73.4
40	311	62.2
50	246	49.2
100	113	22.6
125	66	13.2
150	2	0.4
175	1	0.2
200	1	0.2
225	1	0.2
250	-	-
300	-	-

Table 2. Numbers of parasites recovered from Angora goats.

Irradiation Doses (Gy)	Number of animals	Parasites		Total number of Parasites
		Female	Male	
Control	1	113	96	209
	1	151	107	258
	1	78	63	141
30	1	12	5	17
	1	23	14	37
	1	42	17	59
50	1	-	-	-
	1	8	2	10
	1	-	-	-
100	1	-	-	-
	1	-	-	-
	1	-	-	-

REFERENCES

[1] Alabay. M., Çerçi, H., Aykol, F. (1985). An improved method of obtaining clean infective larvae of *Trichostrongylus vitrinus*. Doğa Bilim Derg., 9: 229-231.

[2] Celep, A. (1987). Samsun yöresi kuzu ve toklularında paraziter fauna tebsiti ile kontrol ve tedavi gruplarında aylık ortalama ağırlık artışlarının belirlenmesine dair araştırmalar. Türk Vet. Hek. Dern. Derg., 57: 69-79

[3] Çetindağ, M., Bıyıkoglu, G. (1997). İç Anadolu bölgesi tiftik keçilerinde mide barsak nematodlarının yayılışı. 10. Ulusal Parazitoloji Kongresi, Bildiri Özetleri, 2: 79 Ankara, 8-12 Eylül.

[4] Umur, T. (1991). Ankara yöresi tiftik keçilerinde sindirim sistemi helmintleri. Ankara Üniv. Vet. Fak. Derg., 38: 322-338.

[5] Vural, A., Onar, E., Özkoç, Ü., Everett, G. (1968). İstanbul iline bağlı köylerde koyunlarda gastrointestinal nematod, moniezia ve karaciğer trematodlarının yayılış durumu. Pendik Vet. Mikrobiyol. Derg., 1: 146-159

[6] Zeybek, H. (1980). Samsun yöresi koyun ve kuzularında paraziter fauna saptama çalışmaları. Ankara üniv. Vet. Fak. Derg., 27: 215-236.

INSTRUCTION FOR AUTHORS

General Requirements:

Contributions to the Turkish Journal of Nuclear Sciences may be:

- a) Original research findings which are the result of investigations or experimentations. They may have an objective of investigations or experimentations. They may be on new or improved techniques in every field of nuclear science;
 - b) Critical comments or articles published in the Turkish Journal of Nuclear Sciences;
 - c) Short, critical reviews or essays on timely subjects.
- Only contributions that have not been published previously, will be considered for publication.

Manuscript Requirements:

- 1) Three copies of double-spaced, typewritten manuscript in English should be submitted to the Managing Editor, TAEK (Eskişehir Yolu 06530, ANKARA)
- 2) Manuscripts should include Title, Abstracts (both in English and Turkish), Introduction, Materials and Methods, Results and Discussion, Acknowledgements (if applicable) and References.
- 3) A short title which must accurately identify and describe manuscript contents is also required.
- 4) References should be numbered in square brackets throughout the text and at the end of the article. Every references should include the whole title of the reference.

The method of citation should be as follows:

- [1] Nagy, S., Csikai, J, "Nuclear Techniques". Nucl.Instr.Meth.91 (1971) 345.
 - [2] Covell, D.F. "Phosphorus Adsorption Isotherms". Anal. Chem. 31 (1959) 185.
 - [3] Lederer, L.M., Perlman, I. "Table of Isotopes", 6th ed., John Wiley, New York (1967).
- 6) Figures and tables should accompany the manuscript separately and legends should also be typed on a separate sheet.
 - 7) If contributions is supported by an organization, the name of the sponsor should be indicated at the bottom of the first page.
 - 8) Footnotes should be indicated by asterisk, and they should appear at the bottom of the page and be separated from the main text by a line.
 - 9) Photographs must be glossy prints and figures must be drawn on tracing paper.