PAPER DETAILS

TITLE: Periodic Korovkin Theorem via \$P_{p}^{2}-Statistical \$\\mathcal{A}}-Summation Process

AUTHORS: Sevda YILDIZ, Fadime DIRIK, Kamil DEMIRCI

PAGES: 156-162

ORIGINAL PDF URL: https://dergipark.org.tr/tr/download/article-file/2775306

UJMA

Universal Journal of Mathematics and Applications

Journal Homepage: www.dergipark.gov.tr/ujma ISSN 2619-9653 DOI: https://doi.org/10.32323/ujma.1205420



Periodic Korovkin Theorem via P_p^2 -Statistical \mathscr{A} -Summation Process

Sevda Yıldız¹, Fadime Dirik^{1*} and Kamil Demirci¹

¹Department of Mathematics, Faculty of Arts and Sciences, Sinop University, 57000, Sinop, Türkiye *Corresponding author

Article Info	Abstract
Keywords: Korovkin theorem, periodic functions, power series method, rates of convergence, statistical convergence. 2010 AMS: 40C05, 40G10, 41A25, 41A36. Received: 15 November 2022 Accepted: 27 December 2022 Available online: 29 December 2022	In the current research, we investigate and establish Korovkin-type approximation theorems for linear operators defined on the space of all 2π -periodic and real valued continuous functions on \mathbb{R}^2 by means of \mathscr{A} -summation process via statistical convergence with respect to power series method. We demonstrate with an example how our theory is more strong than previously studied. Additionally, we research the rate of convergence of positive linear operators defined on this space.

1. Introduction and Preliminaries Notations

Before starting with the presentation of the definitions which will be used to prove approximation theorems, we recall the well-known notions.

A double sequence $x = (x_{ij})$ is convergent to *L* in Pringsheim's sense if, for every $\varepsilon > 0$, there exists $N = N(\varepsilon) \in \mathbb{N}$ such that $|x_{ij} - L| < \varepsilon$ whenever i, j > N and denoted by $P - \lim_{i,j} x_{ij} = L$ (see [1]). A double sequence is bounded if there exists a positive number *M* such that

 $|x_{ij}| \le M$ for all $(i, j) \in \mathbb{N}^2 = \mathbb{N} \times \mathbb{N}$. As it is known that every single convergent sequence (in the usual sense) is bounded, while a convergent double sequence need not to be bounded.

Let us turn our attention to statistical convergence and power series method for double sequences.

Moricz [2] proposed and investigated the idea of statistical convergence for double sequences, which may be restated in terms of natural density. Let $E \subset \mathbb{N}^2$ be a two-dimensional subset of positive integers and let $E_{m,n} = \{(i, j) \in E : i \leq m, j \leq n\}$. Then the two-dimensional analogue of natural density can be defined as follows:

$$\delta_2(E) := P - \lim_{m,n} \frac{1}{mn} |E_{mn}|$$

if it exists. The number sequence $x = (x_{ij})$ is statistically convergent to *L* provided that for every $\varepsilon > 0$, the set $E := E_{mn}(\varepsilon) := \{i \le m, j \le n : |x_{ij} - L| \ge \varepsilon\}$ has natural density zero; in that case we write $st_2 - \lim_{i,j} x_{ij} = L$. For all that a statistically convergent sequence need not be convergent in light of the above.

It is obvious that a double sequence that is *P*-convergent statistically converges to the same value, but the opposite is not always true. Additionally, Moricz [2] characterized the statistical convergence for double sequences as follows:

A double sequence $x = (x_{ij})$ is statistically convergent to *L* if and only if there exists a set $E \subset \mathbb{N}^2$ such that the natural density of *E* is 1 and $P = \lim_{x \to \infty} x_{ij} = L$

$$P - \lim_{\substack{i,j \to \infty \\ \text{and } (i,j) \in E}} x_{ij} = I$$

Let (p_{ij}) be a double sequence of nonnegative numbers with $p_{00} > 0$ and such that the following power series

$$p(t,s) := \sum_{i,j=0}^{\infty} p_{ij} t^i s^j$$

Email addresses and ORCID numbers: sevdaorhan@sinop.edu.tr, 0000-0002-4730-2271 (S. Yıldız), fdirik@sinop.edu.tr, 0000-0002-9316-9037 (F. Dirik), kamild@sinop.edu.tr, 0000-0002-5976-9768 (K. Demirci)



has radius of convergence *R* with $R \in (0,\infty]$ and $t, s \in (0,R)$. If for all $t, s \in (0,R)$,

$$\lim_{t,s\to R^{-}}\frac{1}{p(t,s)}\sum_{i,j=0}^{\infty}p_{ij}t^{i}s^{j}x_{ij}=L$$

then we say that the double sequence $x = (x_{ij})$ is convergent to *L* in the sense of power series method and denoted by $P_p^2 - \lim x_{ij} = L$ ([3]). Keep in mind that the method is regular if and only if

$$\lim_{s \to R^{-}} \frac{\sum_{i=0}^{\infty} p_{iv} t^{i}}{p(t,s)} = 0 \text{ and } \lim_{t,s \to R^{-}} \frac{\sum_{j=0}^{\infty} p_{\mu j} s^{j}}{p(t,s)} = 0, \text{ for any } \mu, \nu,$$
(1.1)

hold [3].

t

Remark 1.1. In case of R = 1, if $p_{ij} = 1$ and $p_{ij} = \frac{1}{(i+1)(j+1)}$, the power series methods coincide with Abel summability method and logarithmic summability method, respectively. In the case of $R = \infty$ and $p_{ij} = \frac{1}{i!j!}$, the power series method coincides with Borel summability method.

Here and throughout the paper power series method is always assumed to be regular.

Ünver and Orhan [4] have recently introduced P_p -density of $E \subset \mathbb{N}_0$ and the definition of P_p -statistical convergence for single sequences. A natural question is what about statistical convergence or P_p -statistical convergence of the sequence. Hence, they showed that statistical convergence and P_p -statistical convergence are incompatible. In view of their work, Yıldız, Demirci and Dirik [5] have introduced the definitions of P_p^2 -density of $F \subset \mathbb{N}_0^2 = \mathbb{N}_0 \times \mathbb{N}_0$ and P_p^2 -statistical convergence for double sequences:

Definition 1.2 ([5]). Let $F \subset \mathbb{N}_0^2$. If the limit

$$\delta_{P_p}^2(F) := \lim_{t,s \to R^-} \frac{1}{p(t,s)} \sum_{(i,j) \in F} p_{ij} t^i s^j$$

exists, then $\delta_{P_p}^2(F)$ is called the P_p^2 -density of F. Note that, from the definition of a power series method and P_p^2 -density it can be established that $0 \le \delta_{P_p}^2(F) \le 1$ whenever it exists.

Definition 1.3 ([5]). Let $x = (x_{ij})$ be a double sequence. Then x is said to be statistically convergent with respect to power series method $(P_p^2$ -statistically convergent) to L if for any $\varepsilon > 0$

$$\lim_{t,s\to R^{-}}\frac{1}{p(t,s)}\sum_{(i,j)\in F_{\varepsilon}}p_{ij}t^{i}s^{j}=0$$

where $F_{\varepsilon} = \{(i, j) \in \mathbb{N}_0^2 : |x_{ij} - L| \ge \varepsilon\}$, that is $\delta_{P_p}^2(F_{\varepsilon}) = 0$ for any $\varepsilon > 0$. In this case we write $st_{P_p}^2 - \lim x_{ij} = L$.

Let $A = [a_{klmn}], k, l, m, n \in \mathbb{N}$, be a four-dimensional infinite matrix. The A-transform of a given double sequence $x = (x_{mn})$ is given by

$$(Ax)_{kl} = \sum_{(m,n)\in\mathbb{N}^2} a_{klmn} x_{mn}, \quad k,l\in\mathbb{N}$$

provided the double series converges in Pringsheim's sense for every $(k,l) \in \mathbb{N}^2$ and denoted by $Ax := ((Ax)_{kl})$. If the *A*-transform of *x* exists for all $k, l \in \mathbb{N}$ and convergent in the Pringsheim's sense i.e.,

$$P - \lim_{p,q} \sum_{m=1}^{p} \sum_{n=1}^{q} a_{klmn} x_{mn} = y_{kl} \text{ and } P - \lim_{k,l} y_{kl} = L$$

then we say that a sequence x is A-summable to L. A two-dimensional matrix transformation is referred to as regular in summability theory if it converts each convergent sequence into one with the same limit.

Now consider a sequence of four-dimensional infinite matrices with non-negative real elements $\mathscr{A} := (A^{(i,j)}) = (a_{klmn}^{(i,j)})$. For a given double sequence of real numbers, $x = (x_{mn})$ is said to be \mathscr{A} -summable to L if

$$P - \lim_{k,l} \sum_{(m,n)\in\mathbb{N}^2} a_{klmn}^{(i,j)} x_{mn} = L$$

uniformly in *i* and *j*.

 \mathscr{A} -summability is the *A*-summability for four-dimensional infinite matrix if $A^{(i,j)} = A$, four-dimensional infinite matrix. Some results regarding matrix summability method for double sequences may be found in the papers [6,7]. $C^*(\mathbb{R}^2)$ stands for the space of all continuous functions on \mathbb{R}^2 that are real valued and have a period of 2π . If a function $h \in C^*(\mathbb{R}^2)$, then

$$h(x,y) = h(x+2k\pi, y) = h(x, y+2k\pi)$$
, for all $(x, y) \in \mathbb{R}^2$,

holds for $k = 0, \pm 1, \pm 2, \dots$ In what follows, this space is equipped with the supremum norm

$$||f||_{*} = \sup_{(x,y)\in\mathbb{R}^{2}} |h(x,y)|, (h \in C^{*}(\mathbb{R}^{2})).$$

A sequence $\mathbb{L} := (L_{mn})$ of positive linear operators from $C^*(\mathbb{R}^2)$ into itself is referred to as an \mathscr{A} -summation process on $C^*(\mathbb{R}^2)$ if $(L_{mn}h)$ is \mathscr{A} -summable to h for every $h \in C^*(\mathbb{R}^2)$, i.e.,

$$P - \lim_{k,l} \left\| S_{klij}^{\mathbb{L}} h - h \right\|_* = 0, \text{ uniformly in } i, j$$

where for all $k, l, i, j \in \mathbb{N}, h \in C^*(\mathbb{R}^2)$ the series

$$S_{klij}^{\mathbb{L}}h := \sum_{(m,n)\in\mathbb{N}^2} a_{klmn}^{(i,j)} L_{mn}h$$

$$\tag{1.2}$$

and it is assumed that the series in (1.2) absolutely convergent for each $i, j, k, l \in \mathbb{N}$ and h. For the rate of convergence, we need to recall the following modulus of continuity of h. Let $h \in C^*(\mathbb{R}^2)$, then

$$w(h;\gamma) = \sup\left\{ |h(u,v) - h(x,y)| : (u,v), (x,y) \in \mathbb{R}^2 \text{ and } \sqrt{(u-x)^2 + (v-y)^2} \le \gamma \right\}$$

for $\gamma > 0$. This definition yields the following basic property for $h \in C^*(\mathbb{R}^2)$. For any a > 0,

$$w(h;a\gamma) \leq (1+[a])w(h;\gamma)$$

where [a] is defined to be the greatest integer less than or equal to a.

The paper of Korovkin [8] is an important issue. It can help us to understand the nature of approximation of sequences. This approximation problem has a rich history associated with the names of the different convergence methods on some spaces in the theory. For some recent research works in this direction, see [9–21]. In this paper, we investigate and establish Korovkin-type approximation theorems for linear operators defined on the space of all 2π -periodic and real valued continuous functions on \mathbb{R}^2 by means of \mathscr{A} -summation process via statistical convergence with respect to power series method. We demonstrate with an example how our theory is more strong than previously studied. Additionally, we research the rate of convergence of positive linear operators defined on this space.

2. The Second Theorem of Korovkin Type

The aim of this section is to deal with approximation of all 2π -periodic and real valued continuous functions on \mathbb{R}^2 by means of \mathscr{A} -summation process via statistical convergence with respect to power series method. Our main result is the following.

Theorem 2.1. Let $\mathscr{A} = (A^{(i,j)})$ be a sequence of four-dimensional infinite matrices. Let $\mathbb{L} = (L_{mn})$ be a sequence of positive linear operators acting from $C^*(\mathbb{R}^2)$ into itself. Assume that (1.2) holds. Then, for all $h \in C^*(\mathbb{R}^2)$

$$st_{P_p}^2 - \lim \left\| S_{klij}^{\mathbb{L}} h - h \right\|_* = 0 \quad uniformly \text{ in } i \text{ and } j$$

$$(2.1)$$

if and only if

$$st_{P_{p}}^{2} - \lim \left\| S_{klij}^{\mathbb{L}} h_{r} - h_{r} \right\|_{*} = 0 \quad uniformly \text{ in } i \text{ and } j \quad (r = 0, 1, 2, 3, 4)$$
(2.2)

where $h_0(x,y) = 1$, $h_1(x,y) = \sin x$, $h_2(x,y) = \sin y$, $h_3(x,y) = \cos x$ and $h_4(x,y) = \cos y$.

Proof. Since 1, sin *x*, sin *y*, cos *x* and cos *y* belong to $C^*(\mathbb{R}^2)$, the necessity is clear. Suppose now that (2.2) holds. Let $h \in C^*(\mathbb{R}^2)$ and *I*, *J* be closed subinterval of length 2π of \mathbb{R} . Fix $(x, y) \in I \times J$. As in the proof of Theorem 2.1 in [22], it follows from the continuity of *h* that

$$|h(u,v) - h(x,y)| < \varepsilon + \frac{2M_h}{\sin^2 \frac{\delta}{2}} \varphi(u,v)$$

which gives,

$$\begin{split} \left| \sum_{(m,n)\in\mathbb{N}^{2}} a_{klmn}^{(i,j)} L_{mn}(h;x,y) - h(x,y) \right| &\leq \sum_{(m,n)\in\mathbb{N}^{2}} a_{klmn}^{(i,j)} L_{mn}(|h(u,v) - h(x,y)|;x,y) + |h(x,y)| \left| \sum_{(m,n)\in\mathbb{N}^{2}} a_{klmn}^{(i,j)} L_{mn}(h_{0};x) - h_{0}(x,y) \right| \\ &\leq \sum_{(m,n)\in\mathbb{N}^{2}} a_{klmn}^{(i,j)} L_{mn}\left(\varepsilon + \frac{2M_{h}}{\sin^{2}\frac{v}{2}}\varphi(u,v);x,y\right) + M_{h} \left| \sum_{(m,n)\in\mathbb{N}^{2}} a_{klmn}^{(i,j)} L_{mn}(h_{0};x) - h_{0}(x,y) \right| \\ &\leq \varepsilon + (\varepsilon + M_{h}) \left| \sum_{(m,n)\in\mathbb{N}^{2}} a_{klmn}^{(i,j)} L_{mn}(h_{0};x) - h_{0}(x,y) \right| \\ &+ \left| \sin x \right| \left| \sum_{(m,n)\in\mathbb{N}^{2}} a_{klmn}^{(i,j)} L_{mn}(h_{0};x) - h_{0}(x,y) \right| \\ &+ \left| \sin x \right| \left| \sum_{(m,n)\in\mathbb{N}^{2}} a_{klmn}^{(i,j)} L_{mn}(h_{1};x,y) - h_{1}(x,y) \right| \\ &+ \left| \sin x \right| \left| \sum_{(m,n)\in\mathbb{N}^{2}} a_{klmn}^{(i,j)} L_{mn}(h_{2};x,y) - h_{2}(x,y) \right| \\ &+ \left| \cos x \right| \left| \sum_{(m,n)\in\mathbb{N}^{2}} a_{klmn}^{(i,j)} L_{mn}(h_{3};x,y) - h_{3}(x,y) \right| \\ &+ \left| \cos y \right| \left| \sum_{(m,n)\in\mathbb{N}^{2}} a_{klmn}^{(i,j)} L_{mn}(h_{4};x,y) - h_{4}(x,y) \right| \\ &\leq \varepsilon + N \sum_{r=0}^{4} \left| \sum_{(m,n)\in\mathbb{N}^{2}} a_{klmn}^{(i,j)} L_{mn}(h_{r};x) - h_{r}(x,y) \right| \end{aligned}$$

where $M_h = \|f\|_*$, $\varphi(u, v) = \sin^2 \frac{u-x}{2} + \sin^2 \frac{v-y}{2}$ and $N := \varepsilon + M_h + \frac{2M_h}{\sin^2 \frac{\delta}{2}}$. Then, taking supremum over $(x, y) \in \mathbb{R}^2$, we obtain

$$\left\|S_{klij}^{\mathbb{L}}h - h\right\|_{*} \leq \varepsilon + N \sum_{r=0}^{4} \left\|S_{klij}^{\mathbb{L}}h_r - h_r\right\|_{*}.$$
(2.3)

Now given r > 0, choose $\varepsilon > 0$ such that $\varepsilon < r$, and define

$$D := \left\{ (k,l) : \left\| S_{klij}^{\mathbb{L}} h - h \right\|_{*} \ge r \right\},$$

$$D_{r} := \left\{ (k,l) : \left\| S_{klij}^{\mathbb{L}} h_{r} - h_{r} \right\|_{*} \ge \frac{r - \varepsilon}{5N} \right\}, r = 0, 1, 2, 3, 4$$

It is easy see that from (2.3)

$$D \subseteq \bigcup_{r=0}^{4} D_r.$$

Hence, we may write

$$\delta_{P_p}^2\left(D
ight)\leq\sum_{r=0}^4\delta_{P_p}^2\left(D_r
ight).$$

Then, according to (2.2), we have

$$\delta_{P_p}^2\left(D\right)=0,$$

and hence

$$st_{P_p}^2 - \lim \left\| S_{klij}^{\mathbb{L}} h - h \right\|_* = 0$$
 uniformly in *i* and *j*

which is the desired result.

3. An example

Now, we give an example that our theorem (Theorem 2.1) is stronger than Theorem 9 in [23].

Example 3.1. Now assume that $\mathscr{A} = (A^{(i,j)})$ is a sequence of four-dimensional infinite matrices defined by $a_{klmn}^{(i,j)} = \frac{1}{kl}$ if $i \le m \le k+i-1$, $j \le n \le l+j-1$ and $a_{klmn}^{(i,j)} = 0$ otherwise. Let us consider the double sequence of Fejer operators on $C^*(\mathbb{R}^2)$ where

$$L_{mn}(h;x,y) = \frac{1}{(m\pi)(n\pi)} \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} h(u,v) F_m(u) F_n(v) dudv$$
(3.1)

where $F_m(u) = \frac{\sin^2 \frac{m(u-x)}{2}}{2\sin^2 \frac{u-x}{2}}$ and $\frac{1}{\pi} \int_{-\pi}^{\pi} F_m(u) du = 1$. Let (p_{ij}) be defined as follows

$$p_{ij} = \begin{cases} 0, & i \text{ and } j \text{ even} \\ 1, & i \text{ or } j \text{ odd} \end{cases}$$

and take the sequence (x_{ij}) defined by

$$x_{ij} = \begin{cases} ij, & i \text{ and } j \text{ even} \\ 0, & i \text{ or } j \text{ odd} \end{cases}$$
(3.2)

It is easy to see that

$$st_{P_p}^2 - \lim x_{ij} = 0. ag{3.3}$$

However, the sequence (x_{ij}) neither statistically convergent to 0 nor Pringsheim convergent. Now using (3.1) and (3.2), we define the following double positive linear operators $\mathbb{T} = (T_{mn})$ on $C^*(\mathbb{R}^2)$ as follows:

$$T_{mn}(f;x,y) = (1+x_{mn})L_{mn}(f;x,y).$$
(3.4)

We now claim that

$$st_{P_p}^2 - \lim \left\| S_{klij}^{\mathbb{T}} h_r - h_r \right\|_* = 0 \quad uniformly \text{ in } i \text{ and } j, \ (r = 0, 1, 2, 3, 4).$$
(3.5)

Observe that $L_{mn}(h_0;x,y) = h_0(x,y)$, $L_{mn}(h_1;x,y) = \frac{m-1}{m}h_1(x,y)$, $L_{mn}(h_2;x,y) = \frac{n-1}{n}h_2(x,y)$, $L_{mn}(h_3;x,y) = \frac{m-1}{m}h_3(x,y)$, $L_{mn}(h_4;x,y) = \frac{n-1}{n}h_4(x,y)$. So, we can see,

$$\left\|S_{klij}^{\mathbb{T}}h_{0}-h_{0}\right\|_{*}=\left\|\sum_{m=i}^{i+k-1}\sum_{n=j}^{j+l-1}\frac{1}{kl}(1+x_{mn})-1\right\|_{*}\leq\frac{1}{kl}\sum_{m=i}^{i+k-1}\sum_{n=j}^{j+l-1}x_{mn}.$$

It is well known that if a sequence is convergent, its arithmetic mean will also converge to the same value. Thus, by virtue of P_p^2 -statistical convergence and thanks to (3.3) it is clear that

$$st_{P_p}^2 - \lim\left(\sup_{i,j} \frac{1}{kl} \sum_{m=i}^{i+k-1} \sum_{n=j}^{j+l-1} x_{mn}\right) = 0,$$
(3.6)

and hence

$$st_{P_p}^2 - \lim \left\| S_{klij}^{\mathbb{T}} h_0 - h_0 \right\|_* = 0, \text{ uniformly in } i \text{ and } j,$$

which guarantees that (3.5) holds true for r = 0. Also, we compute

$$\left\|S_{klij}^{\mathbb{T}}h_{1}-h_{1}\right\|_{*} = \left\|\sum_{m=i}^{i+k-1}\sum_{n=j}^{j+l-1}\frac{1}{kl}(1+x_{mn})\frac{m-1}{m}h_{1}-h_{1}\right\|_{*} \le \left|\frac{1}{kl}\sum_{m=i}^{i+k-1}\sum_{n=j}^{j+l-1}\frac{m-1}{m}-1\right| + \frac{1}{kl}\sum_{m=i}^{i+k-1}\sum_{n=j}^{j+l-1}\frac{x_{mn}(m-1)}{m}+\frac{1}{kl}\sum_{m=i}^{i+k-1}\sum_{n=j}^{j+l-1}\frac{x_{mn}(m-1)}{m}+\frac{1}{kl}\sum_{m=i}^{i+k-1}\sum_{n=j}^{j+l-1}\frac{x_{mn}(m-1)}{m}+\frac{1}{kl}\sum_{m=i}^{i+k-1}\sum_{n=j}^{j+l-1}\frac{x_{mn}(m-1)}{m}+\frac{1}{kl}\sum_{m=i}^{i+k-1}\sum_{n=j}^{j+l-1}\frac{x_{mn}(m-1)}{m}+\frac{1}{kl}\sum_{m=i}^{i+k-1}\sum_{n=j}^{j+l-1}\frac{x_{mn}(m-1)}{m}+\frac{1}{kl}\sum_{m=i}^{i+k-1}\sum_{m=i}^{j+l-1}\frac{x_{mn}(m-1)}{m}+\frac{1}{kl}\sum_{m=i}^{i+k-1}\sum_{m=i}^{j+l-1}\frac{x_{mn}(m-1)}{m}+\frac{1}{kl}\sum_{m=i}^{i+k-1}\sum_{m=i}^{j+l-1}\frac{x_{mn}(m-1)}{m}+\frac{1}{kl}\sum_{m=i}^{i+k-1}\sum_{m=i}^{j+l-1}\frac{x_{mn}(m-1)}{m}+\frac{1}{kl}\sum_{m=i}^{i+k-1}\sum_{m=i}^{j+l-1}\frac{x_{mn}(m-1)}{m}+\frac{1}{kl}\sum_{m=i}^{i+k-1}\sum_{m=i}^{j+l-1}\frac{x_{mn}(m-1)}{m}+\frac{1}{kl}\sum_{m=i}^{i+k-1}\sum_{m=i}^{j+l-1}\frac{x_{mn}(m-1)}{m}+\frac{1}{kl}\sum_{m=i}^{i+k-1}\sum_{m=i}^{j+l-1}\frac{x_{mn}(m-1)}{m}+\frac{1}{kl}\sum_{m=i}^{i+k-1}\sum_{m=i}^{j+l-1}\frac{x_{mn}(m-1)}{m}+\frac{1}{kl}\sum_{m=i}^{i+k-1}\sum_{m=i}^{j+l-1}\frac{x_{mn}(m-1)}{m}+\frac{1}{kl}\sum_{m=i}^{i+k-1}\sum_{m=i}^{j+l-1}\frac{x_{mn}(m-1)}{m}+\frac{1}{kl}\sum_{m=i}^{i+k-1}\sum_{m=i}^{i+k$$

Since $st_{P_p}^2 - \lim\left(\sup_{i,j}\left(\frac{1}{kl}\sum_{m=i}^{i+k-1}\sum_{n=j}^{j+l-1}\frac{m-1}{m}-1\right)\right) = 0$ and from (3.6) we have,

$$st_{P_p}^2 - \lim \left\| S_{klij}^{\mathbb{T}} h_1 - h_1 \right\|_* = 0, \text{ uniformly in } i \text{ and } j.$$

So (3.5) valid for r = 1. Likewise, we have

$$\begin{split} st_{P_{p}}^{2} &- \lim \left\| S_{klij}^{\mathbb{T}} h_{2} - h_{2} \right\|_{*} = 0, \ \text{uniformly in } i \text{ and } j, \\ st_{P_{p}}^{2} &- \lim \left\| S_{klij}^{\mathbb{T}} h_{3} - h_{3} \right\|_{*} = 0, \ \text{uniformly in } i \text{ and } j, \\ st_{P_{p}}^{2} &- \lim \left\| S_{klij}^{\mathbb{T}} h_{4} - h_{4} \right\|_{*} = 0, \ \text{uniformly in } i \text{ and } j. \end{split}$$

So, our claim (3.5) is valid for each r = 0, 1, 2, 3, 4. Then, observe that the double sequence $\mathbb{T} = (T_{mn})$ defined by (3.4) satisfy all hypotheses of Theorem 2.1. Hence, we have, for all $f \in C^*(\mathbb{R}^2)$,

$$st_{P_p}^2 - \lim \left\| S_{klij}^{\mathbb{T}} h - h \right\|_* = 0.$$

Also, since (x_{ij}) is not statistically convergent to 0, (T_{mn}) does not satisfy Theorem 9 in [23].

4. Rates of Convergence

In this section, via \mathscr{A} -summation process via statistical convergence with respect to power series method, we study the rates of convergence of a double sequence of positive linear operators mapping acting from $C^*(\mathbb{R}^2)$ into $C^*(\mathbb{R}^2)$ by means of the modulus of continuity. We have the following result.

Theorem 4.1. Let $\mathscr{A} = (A^{(i,j)})$ be a sequence of four-dimensional infinite matrices. Let $\mathbb{L} = (L_{mn})$ be a double sequence of positive linear operators moving from $C^*(\mathbb{R}^2)$ into $C^*(\mathbb{R}^2)$. Suppose that (1.2) and the following conditions provided: (i) $st_{P_p}^2 - \lim \left\|S_{klij}^{\mathbb{L}}h_0 - h_0\right\|_* = 0$, uniformly in i and j, (ii) $st_{P_p}^2 - \lim w(h; \gamma) = 0$, uniformly in i and j, where $\gamma := \gamma_{(j,k)}^{(i,l)} := \sqrt{\left\|S_{klij}^{\mathbb{L}}\varphi\right\|_*}$ with $\varphi(u, v) = \sin^2 \frac{u-x}{2} + \sin^2 \frac{v-y}{2}$. Then we have, for all $h \in C^*(\mathbb{R}^2)$, $st_{P_p}^2 - \lim \left\|S_{klij}^{\mathbb{L}}h - h\right\|_* = 0$, uniformly in i and j.

Proof. To prove this, we firstly suppose that $(x, y) \in [-\pi, \pi] \times [-\pi, \pi]$ and $h \in C^*(\mathbb{R}^2)$ be fixed, and that Let (*i*) and (*ii*) be provided. Let γ be a positive number. As in the proof of Theorem 9 in [23], since the function *h* is continious, the following inequality is obtained:

$$|h(u,v) - h(x,y)| \le \left(1 + \pi^2 \frac{\sin^2 \frac{u-x}{2} + \sin^2 \frac{v-y}{2}}{\gamma^2}\right) w(h;\gamma)$$

Using the definition of modulus of continuity and since the operators L_{mn} is linear and the positive, we have

$$\begin{split} \left| S_{klij}^{\mathbb{L}}(h;x,y) - h(x,y) \right| &= \left| \sum_{(m,n) \in \mathbb{N}^2} a_{klmn}^{(i,j)} L_{mn}(h;x,y) - h(x,y) \right| \\ &\leq \sum_{(m,n) \in \mathbb{N}^2} a_{klmn}^{(i,j)} L_{mn}\left(|h(u,v) - h(x,y)| ; x, y \right) + |h(x,y)| \left| \sum_{(m,n) \in \mathbb{N}^2} a_{klmn}^{(i,j)} L_{mn}\left(h_0; x, y\right) - h_0(x,y) \right| \\ &\leq w(h;\gamma) \sum_{(m,n) \in \mathbb{N}^2} a_{klmn}^{(i,j)} L_{mn}\left(h_0; x, y\right) + \pi^2 \frac{w(h;\gamma)}{\gamma^2} \sum_{(m,n) \in \mathbb{N}^2} a_{klmn}^{(i,j)} L_{mn}\left(\varphi; x, y\right) \\ &+ |h(x,y)| \left| \sum_{(m,n) \in \mathbb{N}^2} a_{klmn}^{(i,j)} L_{mn}\left(h_0; x, y\right) - h_0(x, y) \right| \end{split}$$

where $\varphi(u,v) = \sin^2 \frac{u-x}{2} + \sin^2 \frac{v-y}{2}$. If supremum over (x,y) is taken on both sides of the above inequality and is chosen $\gamma := \gamma_{(j,k)}^{(i,l)} := \sqrt{\left\|S_{klij}^{\mathbb{L}}\varphi\right\|_{*}}$, then we obtain

$$\left\|S_{klij}^{\mathbb{L}}h - h\right\|_{*} \leq w\left(h;\gamma_{(j,k)}^{(i,l)}\right) \left\|S_{klij}^{\mathbb{L}}h_{0} - h_{0}\right\|_{*} + \left(1 + \pi^{2}\right) w\left(h;\gamma_{(j,k)}^{(i,l)}\right) + M_{h}\left\|S_{klij}^{\mathbb{L}}h_{0} - h_{0}\right\|_{*}$$
(4.1)

where $M_h := \|h\|_*$. Now, given $\varepsilon > 0$, define the following sets:

$$D := \left\{ (k,l) : \left\| S_{klij}^{\mathbb{L}} h - h \right\|_{*} \ge \varepsilon \right\},$$

$$D_{1} := \left\{ (k,l) : w \left(h; \gamma_{(j,k)}^{(i,l)} \right) \left\| S_{klij}^{\mathbb{L}} h_{0} - h_{0} \right\|_{*} \ge \frac{\varepsilon}{3} \right\}$$

$$D_{2} := \left\{ (k,l) : w \left(h; \gamma_{(j,k)}^{(i,l)} \right) \ge \frac{\varepsilon}{3\left(1 + \pi^{2}\right)} \right\},$$

$$D_{3} := \left\{ (k,l) : \left\| S_{klij}^{\mathbb{L}} h_{0} - h_{0} \right\|_{*} \ge \frac{\varepsilon}{3M_{h}} \right\}.$$

Then, it follows from (4.1) that $D \subset D_1 \cup D_2 \cup D_3$. Also, defining

$$D_4 := \left\{ (k,l) : w\left(h; \gamma_{(j,k)}^{(i,l)}\right) \ge \sqrt{\frac{\varepsilon}{3}} \right\},$$

$$D_5 := \left\{ (k,l) : \left\| S_{klij}^{\mathbb{L}} h_0 - h_0 \right\|_* \ge \sqrt{\frac{\varepsilon}{3}} \right\},$$

we have $D_1 \subset D_4 \cup D_5$, which yields

$$D\subseteq \bigcup_{i=2}^5 D_i.$$

Hence, we may write

$$\delta_{P_p}^2(D) \leq \sum_{r=0}^5 \delta_{P_p}^2(D_r).$$

Using the hypothesis (i) and (ii), we get

$$\delta_{P_n}^2(D) = 0,$$

and hence

$$st_{P_p}^2 - \lim \left\| S_{klij}^{\mathbb{L}} h - h \right\|_* = 0$$
, uniformly in *i* and *j*.

Therefore, the proof is completed.

5. Conclusion

The paper contains Korovkin-type approximation theorem and the rate of convergence for linear operators defined on the space of all 2π -periodic and real valued continuous functions on \mathbb{R}^2 by means of \mathscr{A} -summation process via statistical convergence with respect to power series method. Also, it is demonstrated with an example how the new theory is more stronger than previously studied.

Acknowledgements

The authors would like to express their sincere thanks to the editor and the anonymous reviewers for their helpful comments and suggestions.

Funding

There is no funding for this work.

Availability of data and materials

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author's contributions

All authors contributed equally to the writing of this paper. All authors read and approved the final manuscript.

References

- [1] A. Pringsheim, Zur theorie der zweifach unendlichen zahlenfolgen, Math. Ann., 53 (1900), 289-321.
- [2] F. Moricz, Statistical convergence of multiple sequences, Arch. Math. (Basel), 81 (2004), 82-89.
- [3] S. Baron, U. Stadtmüller, Tauberian theorems for power series methods applied to double sequences, J. Math. Anal. Appl., 211(2) (1997), 574-589.
- [4] M. Ünver, C. Orhan, Statistical convergence with respect to power series methods and applications to approximation theory, Numerical Functional
- Analysis and Optimization, **40**(5) (2019), 535-547.
- [5] S. Yıldız, K. Demirci, F. Dirik, Korovkin theory via P_p-statistical relative modular convergence for double sequences, Rend. Circ. Mat. Palermo, II. Ser, (2022), 1-17.
- [6] R. F. Patterson, E. Savaş, Uniformly summable double sequences, Studia Scientiarum Mathematicarum Hungarica, 44 (2007), 147-158.
- [7] E. Savaş, B. E. Rhoades, *Double summability factor theorems and applications*, Math. Inequal. Appl., **10** (2007), 125-149.
- [8] P. P. Korovkin, On convergence of linear positive operators in the space of continuous functions, Doklady Akademii Nauk, 90 (1953), 961-964 (Russian).
 [9] Ö.G. Atlıhan, C. Orhan, Summation process of positive linear operators, Computers and Mathematics with Applications, 56 (2008), 1188-1195.
- [10] N. S. Bayram, Strong summation process in locally integrable function spaces, Hacettepe Journal of Mathematics and Statistics, 45(3) (2016), 683-694.
- [11] N. S. Bayram, C. Orhan, A-Summation process in the space of locally integrable functions, Stud. Univ. Babes-Bolyai Math., 65 (2020), 255-268.
- [12] S. Çınar, S. Yıldız, *P-statistical summation process of sequences of convolution operators*, Indian Journal of Pure and Applied Mathematics, **53**(3) (2022), 648-659.
- [13] F. Dirik, K. Demirci, Korovkin type approximation theorem for functions of two variables in statistical sense, Turkish Journal of Mathematics, **34**(1) (2010), 73-84.
- [14] F. Dirik, K. Demirci, B-statistical approximation for periodic functions, Studia Scientiarum Mathematicarum Hungarica, 47(3) (2010), 321-332.
- [15] K. Demirci, F. Dirik, Approximation for periodic functions via statistical σ-convergence, Mathematical Communications, 16(1) (2011), 77-84.
- [16] K. Demirci, S. Orhan, Statistical relative approximation on modular spaces, Results in Mathematics, 71(3) (2017), 1167-1184.
- [17] K. Demirci, S. Yıldız, F. Dirik, Approximation via power series method in two-dimensional weighted spaces, Bulletin of the Malaysian Mathematical Sciences Society, **43**(6) (2020), 3871-3883.
- [18] O. Duman, *Statistical approximation for periodic functions*, Demonstratio Mathematica, **36**(4) (2003), 873-878.
- [19] A. D. Gadjiev, C. Orhan, Some approximation theorems via statistical convergence, The Rocky Mountain Journal of Mathematics, (2002), 129-138.
- [20] S. Orhan, K. Demirci, Statistical A-summation process and Korovkin type approximation theorem on modular spaces, Positivity, 18(4) (2014), 669-686.
 [21] V. I. Volkov, On the convergence of sequences of positive linear operators in the space of two variables, Dokl. Akad. Nauk. SSSR (N.S.) 115 (1957),
- 17-19.
 [22] O. Duman, E. Erkuş, Approximation of continuous periodic functions via statistical convergence, Comput. Math. Appl., 52 (2006) 967-974.
- [23] K. Demirci, F. Dirik, Four-dimensional matrix transformation and rate of A-statistical convergence of periodic functions, Mathematical and Computer Modelling, 52 (2010), 1858-1866.