# PAPER DETAILS

TITLE: PCR detection of Anaplasma phagocytophilum in stray dogs in Batman, Turkey

AUTHORS: Bekir OGUZ, Serdar DEGER

PAGES: 183-185

ORIGINAL PDF URL: https://dergipark.org.tr/tr/download/article-file/860537

Van Vet J, 2019, 30 (3) 183-185



ISSN: 2149-3359

**Original Article** 

e-ISSN: 2149-8644

# PCR detection of Anaplasma phagocytophilum in stray dogs in Batman, Turkey

# Bekir OĞUZ Serdar DEĞER

Van Yuzuncu Yil University, Faculty of Veterinary Medicine, Department of Parasitology, Van, Turkey

Received: 18.09.2019

Accepted: 16.10.2019

**ABSTRACT** Anaplasma phagocytophilum causes high-grade fever in humans, ruminants, horses, cats, and dogs. Anaplasma phagocytophilum, which causes granulocytic anaplasmosis in dogs, has a wide geographical distribution in the world, but there are not many studies on the Anaplasma species in dogs in Turkey. Anaplasma phagocytophilum, which infects leukocytes in dog, might also be zoonotic. This study aimed to investigate the *A. phagocytophilum* prevalence in stray dogs in Batman province of Turkey using Polymerase Chain Reaction (PCR) technique. Blood samples were drawn into EDTA tubes from the venae cephalica antebrachii of randomly selected 97 asymptomatic dogs in total. Identification of *A. phagocytophilum* was carried out using the conventional PCR method that was conducted by using the DNAs obtained from the samples. Anaplasma phagocytophilum prevalence was found to be 3.1% in the examined blood samples. In conclusion, infection of dogs in Batman province with *A. phagocytophilum* has been revealed molecularly for the first time with this study. It is considered that detection of this species, which can cause disease (Human granulocytic anaplasmosis) in humans as well, has contributed epidemiologically to the data on the spread of the disease throughout Turkey. Moreover, investigating the tick species that transmit the disease in the region where the study was conducted is important for developing effective control strategies against the disease.

Keywords: Anaplasma phagocytophilum, Dog, Pathogen, PCR

# öz Batman İli Sokak Köpeklerinde *Anaplasma phagocytophilum*'un PCR ile Belirlenmesi

*Anaplasma phagocytophilum* insan, ruminant, at, kedi ve köpeklerde şiddetli ateşli hastalıklara neden olmaktadır. Köpeklerde granülositik anaplasmosise neden olan *A. phagocytophilum* dünyada geniş bir coğrafik dağılım göstermektedir. Ancak Türkiye'de köpeklerde *Anaplasma* türleri üzerine çok fazla çalışma bulunmamaktadır. Bu çalışma, Batman ili sokak köpeklerinde *A .phagocytophilum* yaygınlığının Polimeraz Zincir Reaksiyonu (PZR) yöntemi ile araştırılması amacıyla yapılmıştır. Rastgele seçilen toplam 97 asemptomatik köpeğin vena cephalica antebrachii'lerinden EDTA'lı tüplere kan örnekleri alınmıştır. Alınan örneklerden elde edilen DNA'lar kullanılarak yapılan konvensiyonel PCR yöntemi ile A. *phagocytophilum* identifikasyonu gerçekleştirilmiştir. İncelenen kan örneklerinde *A. phagocytophilum* prevalansı %3.1 olarak bulunmuştur. Sonuç olarak, bu çalışma ile Batman ili sokak köpeklerinin *A. phagocytophilum* ile enfeksiyonu moleküler olarak ilk kez ortaya konmuştur. İnsanlarda da hastalık oluşturabilen (Human granulocytic anaplasmosis) bu türün tespit edilmesinin, hastalığın Türkiye genelinde yayılışı konusundaki verilere epidemiyolojik anlamda katkı sağladığı düşünülmektedir. Ayrıca, çalışmanın yapıldığı bölgede enfeksiyonu nakleden kene türlerinin araştırılması hastalığa karşı etkili kontrol stratejilerinin geliştirilmesi bakımından önem taşımaktadır.

Anahtar Kelimeler: Anaplasma phagocytophilum, Köpek, Pathojen, PCR

## INTRODUCTION

*Anaplasma phagocytophilum* is an obligate intracellular bacterium which causes tick-borne zoonotic disease and can infect primarily neutrophils and also eosinophils (Carrade et al. 2009). This species, which causes granulocytic anaplasmosis in dogs, has been identified also in cats (Bjo<sup>°</sup>ersdorff et al. 1990), horses (Passamonti et al.

2010), dogs (Bjo<sup>°</sup>ersdorff et al. 1990, Jennifer et al. 2009), rodents (Liz et al. 2000, Christova and Gladnishka 2005), deer (Jenkins et al. 2001) and humans (Petrovec et al. 1997) besides domestic ruminants. It is considered that dogs, horses, and humans are accidental hosts. Granulocytic anaplasmosis in dogs was first reported in California in 1982 (Madewell and Gribble 1982), later its existence has been revealed at increasing rates year by year (Carrade et al. 2009). The vector transmitting the disease is the tick species of Ixodes ricinus (I. ricinus) in Europe, Ixodes persulcatus (I. persulcatus) in Asia, and Ixodes pacificus (I. pasificus) and Ixodes scapularis (I. scapularis) in North America (Woldehiwet 2010). In the regions where the disease is endemic, any clinical findings were not observed in many of the dogs that were detected to be 60% seropositive. However, lethargy, inappetence, and fever are observed in the dogs infected with A. phagocytophilum (Beall et al. 2008; Eberts et al. 2011). Splenomegaly and lymphadenopathy have been reported in anaplasmosis and it was concluded that they were caused by lymphoid hyperplasia. In a great portion (90%) of anaplasmosis cases that are caused bv phagocytophilum, thrombocytopenia Α. develops depending on the destruction of thrombocytes (Eberts et al. 2011). Diagnosis of A. phagocytophilum infections in dogs is usually made by the serological methods such as immunofluorescence antibody technique (IFAT) and rapid diagnostic ELISA test. However, cross-reactivities may be observed in the dogs infected with A. phagocytophilum and A. platvs. It has been reported that this creates a serious disadvantage in regions where both two species progress together (Carrade et al. 2009). In recent years, PCR method, which has a higher specificity and sensitivity than microscopic and serological diagnosis methods, has become a reason for preference in the diagnosis of anaplasmosis in dogs. By means of this technique, the infection can be detected even in the animals having very low parasitemia and thus reservoir animals can be determined (Massung and Slater 2003; Santos et al. 2011; Ebani et al. 2013; Vargas-Hernandez et al. 2016; Huber et al. 2017). In Turkey, the number of studies on the Anaplasma species in dogs is limited. Blood samples of 371 dogs in the Aegean Region were examined by nested PCR in terms of A. platys and A. phagocytophilum, and A. platys was detected in 146 dogs (39.4%), and A. phagocytophilum was detected in 193 (52%) of them (Karagenc et al. 2005). Moreover, a case related to A. platys was reported (Ulutas et al. 2007). However, the existence of A. phagocytophilum was shown both in possible vectors and in cattle, sheep, and wild mice using both serological and molecular methods (Aktas et al. 2010, 2011; Sen et al. 2011).

This study aimed to investigate the *A. phagocytophilum* prevalence in the peripheral whole blood samples of stray dogs in Batman province by using the PCR technique.

#### **MATERIALS and METHODS**

In the Veterinary Technical Health Center of the Municipality of Batman in various periods in 2018, blood samples were drawn into the EDTA tubes from the venae cephalica antebrachii of 97 asymptomatic dogs in total. The animals from which the blood samples were collected were the stray dogs that were brought to the Technical Health Center for sterilization or treatment. Animal ethics, method and sampling approval was obtained from the University of Van Yuzuncu Yil Animal Ethics Committee (approval number: 05/04/2019-27762). All samples were stored at -20°C until the laboratory studies were conducted. Genomic DNA was obtained from the blood

## DISCUSSION

Currently, it is important to know the presence and spread of the tick-borne parasites in dogs, and to reveal their molecular epizootiologies for necessary treatment, control and protection measures. *Anaplasma phagocytophilum*, one of these blood parasites, is a zoonotic species which is samples collected from the stray dogs by using commercial blood kit (GeneAll® ExgeneTM Tissue SV (plus!), Korea). The obtained DNA samples were stored at -20°C until the PCR examinations were carried out. In analyzing the obtained DNA samples in terms of A. phagocytophilum, they were processed by KO97 numbered commercial PCR kit (Genekam Biotechnology, Germany) in compliance with the procedure that the producing company had reported. The amplification products were electrophoresed on 1.5% agarose gel and were evaluated at the gel documentation system (Avegene, Taiwan).

#### RESULTS

In the examinations conducted using PCR method, 3 (3.1%) of the 97 asymptomatic dogs were found infected with *A. phagocytophilum*. The information on sex and gender of the examined animals were summarized in Table 1.

**Table 1.** Relation to age and gender of the canine study population

		Study population No. = 97	PCR positive dogs No. = 3.1 (%)
Gender			
Age	Female	50	2
(year)	Male	47	1
	<1	14	0
	1-2	50	2
	>2	33	1

As a result of the amplification and the agarose gel electrophoresis of the DNA that was extracted from the blood samples in the PCR, positive bands of about 444 bp long were obtained (Figure 1).



**Figure 1.** Appearance on agarose gel electrophoresis (1.5%) stained with ethidium bromide of *A. phagocytophilum* DNA extracted from the blood samples. M: 50 bp DNA ladder, 87, 97, 77: Positive samples, NC: Negative control, PC: *A. phagocytophilum* Positive control.

transtadially transmitted to humans, dogs, and other animals through ticks of family Ixodidae. Several methods were used in the diagnosis of this parasite in dogs. Although it is the oldest among these models, microscopic examination is still a reliable method (Sainz et al. 2015). This method facilitates diagnosis of the disease mostly in its acute periods. Besides, serological methods, which are based on the detection of antibodies that develop against the parasite, are also used (Potkonjak et al. 2015). False positives due to the cross-reactivities that emerge during serological diagnosis (especially positives that develop against A. platys species) are tried to be removed by recombinant antigens. Molecular diagnostic methods that are based on the detection of the parasite DNA, which has been widely used in recent years, are also used in the diagnosis of the Anaplasma species, as in the diagnosis of several infectious factors. PCR is used as a sensitive and specific method for detecting the Anaplasma spp. DNA from the clinical samples taken from ticks and animals. Ebani et al. (2013) detected the parasite with the specific primers that they designed from A. phagocytophilum 16S rRNA gene by using Nested-PCR method. Besides, Santos (2011) conducted the identification of et al. A. phagocytophilum in dogs with Real-Time PCR using different gene region and primers. The same researchers revealed partial sequences of the isolates they detected according to msp2 gene region. Melter et al. (2007) conducted the identification of A. phagocytophilum with a PCR in a dog with tick infestation, acute onset of fever, ataxia, and lethargy. The commercial PCR kit protocols which were prepared specifically for A. phagocytophilum were used in this study. The bands with 444 bp long were obtained as a result of the amplification of the positive control and positive samples in the kit protocol. Three (3.1%) of 97 stray dogs in total were found to be infective. In Turkey, a few molecular studies, which were conducted on A. phagocytophilumin dogs by using the conventional and Real-Time PCR techniques, were reported. Pasa et al. (2009) reported that 4 of 10 dogs had been infected with A. phagocytophilum in a study conducted using conventional PCR in the Aegean Region. Duzlu et al. (2014) detected A. phagocytophilum at 7.8% in a molecular study conducted on dogs in the Kayseri region by using Real Time PCR technique. In this study, the molecular prevalence of A. phagocytophilum in the dogs of Batman province was detected to be 3.1%. This result was found to be in parallel with the prevalence rates reported from the Aegean Region and the Kayseri region.

# CONCLUSION

Granulocytic anaplasmosis has been investigated for the first time in the stray dogs living in Batman province and found to be low. However, it is considered that the study has contributed epidemiologically to the data on the spread of the disease throughout Turkey. More extensive studies, in which vectors are investigated and serological methods are included, are needed in order to reveal this parasite, which has a zoonotic potential for humans, in more detail.

#### REFERENCES

- Aktas M, Altay K, Dumanli N (2011). Molecular detection and identification of Anaplasma and Ehrlichia species in cattle from Turkey. Ticks Tick Borne Dis, 2, 62-65.
- Aktas M, Vatansever Z, Altay K, Aydin MF, Dumanli N (2010). Molecular evidence for *Anaplasma phagocytophilum* in Ixodes ricinus from Turkey. Trans R Soc Trop Med Hyg, 104, 10-15.
- Beall MJ, Chandrashekar R, Eberts MD, Cyr KE, Diniz PP, Mainville C, Hegarty BC, Crawford JM, Breitschwerdt EB (2008). Serological and molecular prevalence of *Borrelia burgdorferi*, *Anaplasma phagocytophilum*, and Ehrlichia species in dogs from Minnesota. Vector Borne Zoonotic Dis, 8, 455-464.
- Bjo"ersdorff A, Christenson D, Johnsson A, Sjo" stro"m AC, Madigan JE (1990). Ehrlichia equi-infektion diagnostiserat hos ha"st (In Swedish). Svensk Veterina"rtidning. 42, 357-360.

- Carrade D, Foley JE, Borjesson DL, Sykes JE (2009). Canine Granulocytic Anaplasmosis: A Review. J Vet Intern Med, 23(6), 1129-1141.
- **Christova I, Gladnishka T (2005).** Prevalence of infection with Francisella tularensis, *Borrelia burgdorferi* sensulato and *Anaplasma phagocytophilum* in rodenst from an endemic focus of tularaemia in Bulgaria. Ann Agric Environ Med, 12, 149-152.
- Duzlu O, Inci A, Yildirim A, Onder Z, Ciloglu A (2014). Köpeklerde kene kaynaklı bazı protozoon ve rickettsial enfeksiyonların Real Time PCR ile araştırılması ve saptanan izolatların molekuler karakterizasyonları. Ankara Univ Vet Fak Derg, 61, 275-282.
- Ebani VV, Bertelloni F, Turchi B, Cerri D (2013). Serological and molecular survey of *Anaplasma phagocytophilum* in Italian hunting dogs. Ann Agric Environ Med, 20(2), 289-292.
- Eberts MD, Diniz PP, Beall MJ, Stillman BA, Chandrashekar R, Breitschwerdt EB (2011). Typical and atypical manifestations of *Anaplasma phagocytophilum* infection in dogs. J Am Anim Hosp Assoc, 47, 88-94.
- Huber D, Rei I, Duvnjank S, Jurković D, Lukačevic D, Pilat M, Beck A, Mihaljević Z, Vojta L, Polkinghorne A, Beck R (2017). Molecular detection of Anaplasma platys, Anaplasma phagocythophilum and Wolbachia sp. but not Ehrlichia canis in Croatian dogs. Parasitol Res, 116, 3019-3026
- Jenkins A, Kristiansen BE, Allum AG, Aakre RK, Strand L, Kleveland EJ, et al. (2001). *Borrelia burgdorferi* sensu lato and Ehrlichia spp. in Ixodes ticks from southern Norway. J Clin Microbiol 39, 3666-3671.
- Jennifer L Granick, P Jane Armstrong, Jeff BB (2009). Anaplasma phagocytophilum infection in dogs: 34 cases (2000-2007). J Am Vet Med Assoc, 234(12), 1559-1565.
- Karagenc T, Hosgor M, Bilgic HB, Pasa S, Kirli G, Eren H (2005). The determination of the prevalence of *E. canis, A. phagocytophila, A. platys* with nested PCR in dogs in the Aegean Region. İzmir, Turkey: 14th National Parasitology Congress, p. 18–25.
- Liz JS, Anderes L, Sumner JW, Massung RF, Gern L, Rutti B, Brossard M (2000). PCR detection of granulocytic ehrlichia in Ixodes ricinus ticks and wild small mammals in western Switzerland. J Clin Microbiol, 39(2), 828.
- Madewell BR, Gribble DH (1982). Infection in two dogs with an agent resembling Ehrlichiaequi. J Am Vet Med Assoc, 180, 512–514.
- Massung RF, Slater KG (2003). Comparison of PCR assays for detection of the agent of human granulocytic ehrlichiosis, Anaplasma phagocytophilum. J Clin Microbiol, 41, 717-722.
- Melter O, Stehlik I, Kinska H, Volfova I, Ticha V, Hulinska D (2007). Infection with *Anaplasma phagocytophilum* in a young dog: a case report. Vet Med (Praha), 52, 207-212.
- Passamonti F, Veronesi F, Cappelli K, Capomaccio S, Coppola G, Marenzoni ML, Piergili FD, Verini SA, Coletti M (2010) . *Anaplasma phagocytophilum* in horses and ticks: a preliminary survey of Central Italy. Comp Immunol Microbiol Infect Dis, 33, 73–83.
- Pasa S, Kıral F, Karagenc T, Atasoy A, Seyrek K (2009). Description of dogs naturally infected with *Hepatozoon canis* in the Aegean region of Turkey. Turk J Vet Anim Sci, 33, 289-295.
- Petrovec M, Lotric Furlan S, Zupanc TA, Strle F, Brouqui P, Roux V, et al. (1997). Human disease in Europe caused by a granulocytic Ehrlichia species. J Clin Microbiol 35, 1556-1559.
- Potkonjak A, Vracar V, Savić S, Lako B, Radosavljevic V, Cincovic M, Suvajdzic L, Jurisic A, Petrovic A (2015). The seroprevalence of *Anaplasma phagocytophilum* infection in dogs in the canine babesiosis, Serbia. Veterinarski Arhiv, 85, 385-394.
- Sainz A, Roura X, Miró G, Estrada-Peña A, Kohn B, Harrus S, Solano-Gallego L (2015). Guideline for veterinary practitioners on canine ehrlichiosis and anaplasmosis in Europe. Parasitol Vectors, 8, 75.
- Santos HA, Pires MS, Vilela JA, Sanotos TM,Facinni JL, Baldani CD,Thome SM, Sanavria A, Massard CL (2011). Detection of *Anaplasma phagocytophilum* in Brazilian dogs by real-time polymerase chain reaction. J Vet Diagn Invest, 23, 770-774
- Sen E, Uchishima Y, Okamoto Y, Fukui T, Kadosaka T, Ohashi N, Masuzawa T (2011). Molecular detection of Anaplasma phagocytophilum and Borrelia burgdorferi in Ixodes ricinus ticks from Istanbul metropolitan area and rural Trakya (Thrace) region of northwestern Turkey. Ticks Tick-borne Dis, 2(2), 94-98.
- Ulutas B, Bayramli G, Karagenc T (2007). First case of Anaplasma (Ehrlichia) platys infection in a dog in Turkey. Turk J Vet Anim Sci, 31(4), 279-282.
- Vargas-Hernandez MR, André DM, Cendales KCM, de Sousa LR, Gonçalves MCH, Rondelli, Machado RZ, Tinucci-Costa M (2016). Molecular detection of Anaplasma species in dogs in Colombia, Rev. Bras. Parasitol Veterinária, 25, 459-464.
- Woldehiwet Z (2010). The natural history of *Anaplasma phagocytophilum*. Vet Parasitol, 167(2-4), 108-22.