# PAPER DETAILS

TITLE: Nonlinear Integrodifferential Equations with Time Varying Delay

AUTHORS: Kamalendra KUMAR, Manoj KARNATAK, Rakesh KUMAR

PAGES: 433-444

ORIGINAL PDF URL: https://dergipark.org.tr/tr/download/article-file/1356397

Advances in the Theory of Nonlinear Analysis and its Applications 5 (2021) No. 3, 433–444. https://doi.org/10.31197/atnaa.814109 Available online at www.atnaa.org Research Article



# Nonlinear Integrodifferential Equations with Time Varying Delay

Kamalendra Kumar<sup>a</sup>, Manoj Karnatak<sup>b</sup>, Rakesh Kumar<sup>b</sup>

<sup>a</sup> Department of Mathematics, SRMS College of Engineering and Technology, Bareilly, 243001, U.P., India. <sup>b</sup> Department of Mathematics, Hindu College, Moradabad, U.P., India.

# Abstract

By practicing the manner of semigroup theory and Banach contraction theorem, the existence and uniqueness of mild and classical solutions of nonlinear integrodifferential equations with time varying delay in Banach spaces is showed. Certainly, an example is revealed to justify the abstract idea.

*Keywords:* Nonlinear integrodifferential equations; Time varying delay; Nonlocal condition; Mild and classical solution; Banach contraction theorem. 2010 MSC: Subject Classification 34G20,34K30, 47D03, 47H10.

#### 1. Introduction

In this work, we examine the class of nonlinear integrodifferential equations with time varying delay of form:

$$x'(t) + Ax(t) = F_1\Big(t, x(\gamma_1(t)), \dots, x(\gamma_n(t)), \int_{t_0}^t h_1(t, s, x(\gamma_{n+1}(s)))ds\Big) + F_2\Big(t, x(\eta_1(t)), \dots, x(\eta_m(t)), \int_{t_0}^t h_2(t, s, x(\eta_{m+1}(s)))ds\Big), t \in (t_0, t_0 + b]$$

$$(1)$$

 $\operatorname{and}$ 

$$x(t_0) + g(x) = x_0,$$
(2)

*Email addresses:* kamlendra.14kumar@gmail.com (Kamalendra Kumar), karnatak.manoj@gmail.com (Manoj Karnatak), rakeshnaini1@gmail.com (Rakesh Kumar)

Received October 22, 2020; Accepted: June 14, 2021; Online: June 16, 2021.

in which  $t_0 \ge 0, b > 0$ . The infinitesimal generator that is expressed by -A, of a  $C_0$  semigroup of operators on a Banach spaces.  $F_1, h_1, F_2, h_2$  are functions which is stated in (1) and these functions gratifying some assumptions and  $x_0 \in E$ . Assigning the tool of semigroup, the existence of solutions for semilinear evolution equations is analyzed by Pazy [11]. The same classes of evolution equations as present in [11] with nonlocal condition are explored by Byszewskii [6]. During previous years, differential and integrodifferential system with time varying delay is considered by various investigators like [1] - [5], [7] - [10], [13], [14]. They have used different tools and techniques for discussing the outcomes.

# 2. Preliminaries

In this section, we give some definitions, notations and basic facts which are applied in the next sections.

**Definition 2.1.** [11] A one parameter family  $T(t), 0 \le t < \infty$ , of bounded linear operators from  $E \to E$ , where E is a Banach space, is said to be the semigroup of bounded linear operators on E if

(i) T(0) = I, the identity operator on E;

(ii)  $T(t)T(s) = T(t+s); \forall t, s \ge 0$  (the semigroup property).

A semigroup of bounded linear operators T(t) is uniformly continuous

$$\lim_{t \to 0} \| T(t) - I \| = 0.$$

If the linear operator A explained by

$$D(A) = \left[ y \in E : \lim_{t \to 0} \frac{T(t)y - y}{t} exists \right]$$

and

$$Ay = \lim_{t \to 0} \frac{T(t)y - y}{t}, y \in D(A)$$

is the infinitesimal generator of the semigroup T(t). Here D(A) is the domain of A.

**Definition 2.2.** [11] A semigroup  $T(t), 0 \le t < \infty$ , of bounded linear operators on E is a strongly continuous semigroup of bounded linear operators if

$$\lim_{t \to 0} T(t)y = y, \quad \forall y \in E$$

A strongly continuous semigroup of bounded linear operators on E will be termed as  $C_0$ -semigroup.

**Theorem 2.3.** Suppose T(t) be a  $C_0$ -semigroup. The constants  $\Omega \ge 0$  and  $M \ge 1$  exist such that

$$|| T(t) || \le M e^{\Omega t}, 0 \le t < \infty.$$

If  $\Omega = 0$  then T(t) is called uniformly bounded and if M = 1 it is said to be  $C_0$ -semigroup of contraction.

Now, if E is assumed as a Banach space with norm  $\| . \|$ . Also,  $C_0$ -semigroup of operators on E is written by  $\{T(t)\}_{t\geq 0}$ . Throughout paper, infinitesimal generator is represented by -A and the same is  $C_0$ -semigroup of operators on E. Here the domain of A is given by D(A) and also  $t_0 \geq 0, b > 0$ ,

$$J := [t_0, t_0 + b],$$

$$\Delta := \{ (t, s) : t_0 \le s \le t \le t_0 + b \}$$
$$M := \sup_{t \in [0, b]} \| T(t) \|_{BL(E, E)},$$
$$X := C(J, E)$$

and  $F_1: J \times E^{n+1} \to E, h_1: \Delta \times E \to E, F_2: J \times E^{m+1} \to E, h_2: \Delta \times E \to E, g: X \to E, \gamma_i: J \to J$   $(i = 1, 2, ..., n + 1), \eta_j: J \to J$  (j = 1, 2, ..., m + 1) are stated functions and these functions meet some assumptions.

For the suitability, the operator norm  $\| . \|_{BL(E,E)}$  will be indicated by  $\| T(t) \|$ .

The following two definitions will be mandatory for the mild and classical solutions of the system (1) - (2).

**Definition 2.4.** The following integral equation is fulfilled by the function  $x \in X$ ,

$$\begin{aligned} x(t) &= T(t-t_0)x_0 - T(t-t_0)g(x) \\ &+ \int_{t_0}^t T(t-s)F_1\Big(s, x(\gamma_1(s)), ..., x(\gamma_n(s)), \int_{t_0}^s h_1(s, \tau, x(\gamma_{n+1}(\tau)))d\tau\Big)ds \\ &+ \int_{t_0}^t T(t-s)F_2\Big(s, x(\eta_1(s)), ..., x(\eta_m(s)), \int_{t_0}^s h_2(s, \tau, x(\eta_{m+1}(\tau)))d\tau\Big)ds, \\ &\quad t \in (t_0, t_0 + b] \end{aligned}$$
(3)

is remarked to be mild solution of the system (1) - (2) on J.

**Definition 2.5.** A function  $x: J \to E$  is termed as classical solution of the system (1) - (2) on J if:

(i) x is a continuous on J and is continuously differentiable on  $J/{t_0}$ .

$$(ii) \ x'(t) + Ax(t) = F_1\Big(t, x(\gamma_1(t)), \dots, x(\gamma_n(t)), \int_{t_0}^t h_1(t, s, x(\gamma_{n+1}(s)))ds\Big) + F_2\Big(t, x(\eta_1(t)), \dots, x(\eta_m(t)), \int_{t_0}^t h_2(t, s, x(\eta_{m+1}(s)))ds\Big), t \in J/\{t_0\}$$

(*iii*)  $x(t_0) + g(x) = x_0$ 

# 3. Main Results

3.1. Existence of Mild Solution

The existence of mild solution is discussed by means of following theorem.

#### **Theorem 3.1.** Presume that

- (i) -A is the infinitesimal generator of a  $C_0$ -semigroup  $T(t), t \ge 0$  in E such that  $||T(t)|| \le M$ , for some  $M \ge 1$ .
- (ii) Here the function  $F_1: J \times E^{n+1} \to E$  and  $F_2: J \times E^{m+1} \to E$  are continuous. We take constants  $M_1 > 0, M_2 > 0$  in such a manner that  $\forall x_i, y_i \in E, i = 1, 2, ..., n+1$  and  $\forall x_j, y_j \in E, j = 1, 2, ..., m+1$ , we get

$$\|F_1(t, x_1, x_2, ..., x_{n+1}) - F_1(t, y_1, y_2, ..., y_{n+1})\| \le M_1 \left(\sum_{i=1}^{n+1} \|x_i - y_i\|\right)$$
(4)

and

$$\|F_2(t, x_1, x_2, ..., x_{m+1}) - F_2(t, y_1, y_2, ..., y_{m+1})\| \le M_2 \left(\sum_{j=1}^{m+1} \|x_j - y_j\|\right)$$
(5)

(iii) Next,  $h_1, h_2: \Delta \times E \to E$  are continuous functions and we consider constants  $H_1 > 0, H_2 > 0$  in such a way that  $\forall x, y \in E$ ,

$$|| h_1(t,s,x) - h_1(t,s,y)|| \le H_1 || x - y||$$

and

$$|| h_2(t,s,x) - h_2(t,s,y)|| \le H_2 || x - y||$$

(iv) The function  $g: X \to E$  and there is a constant G > 0 such that

$$|| g(u) - g(v)|| \le G || u - v ||_X, \forall u, v \in E$$

(v) The functions  $\gamma_i \in C(J,J), i = 1, 2, ..., n + 1$  and the function  $\eta_j \in C(J,J), j = 1, 2, ..., m + 1$ .

(vi) finally

$$M[G + M_1b(n + H_1b) + M_2b(m + H_2b)] < 1$$
(6)

If all the above conditions are satisfied then the equations (1) - (2) has a unique mild solution on J. Proof. By explaining a mapping  $\phi$  on X by the formula

$$\begin{aligned} (\phi u)(t) &= T(t-t_0)x_0 - T(t-t_0)g(u) \\ &+ \int_{t_0}^t T(t-s)F_1\Big(s, u(\gamma_1(s)), ..., u(\gamma_n(s)), \int_{t_0}^s h_1(s, \tau, u(\gamma_{n+1}(\tau)))d\tau\Big)ds \\ &+ \int_{t_0}^t T(t-s)F_2\Big(s, u(\eta_1(s)), ..., u(\eta_m(s)), \int_{t_0}^s h_2(s, \tau, u(\eta_{m+1}(\tau)))d\tau\Big)ds, \end{aligned}$$

for  $u \in X$  and  $t \in J$ .

It is simple to understand that  $\phi: X \to X$ .

Just now, we shall try to demonstrate that  $\phi$  is a contraction on x. For this plan, make the difference

$$\begin{aligned} (\phi u)(t) &- (\phi v)(t) = -T(t-t_0)[g(u) - g(v)] \\ &+ \int_{t_0}^t T(t-s) \left[ F_1 \Big( s, u(\gamma_1(s)), ..., u(\gamma_n(s)), \int_{t_0}^s h_1(s, \tau, u(\gamma_{n+1}(\tau))) d\tau \Big) \right. \\ &- F_1 \Big( s, v(\gamma_1(s)), ..., v(\gamma_n(s)), \int_{t_0}^s h_1(s, \tau, v(\gamma_{n+1}(\tau))) d\tau \Big) \right] ds \\ &+ \int_{t_0}^t T(t-s) \left[ F_2 \Big( s, u(\eta_1(s)), ..., u(\eta_m(s)), \int_{t_0}^s h_2(s, \tau, u(\eta_{m+1}(\tau))) d\tau \Big) \right. \\ &- F_2 \Big( s, v(\eta_1(s)), ..., v(\eta_m(s)), \int_{t_0}^s h_2(s, \tau, v(\eta_{m+1}(\tau))) d\tau \Big) \right] ds \end{aligned}$$

Now, taking norm both sides, we obtain

$$\| (\phi u)(t) - (\phi v)(t) \| \leq \| T(t - t_0) \| \| g(u) - g(v) \|$$
  
 
$$+ \int_{t_0}^t \| T(t - s) \| \left\| F_1(s, u(\gamma_1(s)), ..., u(\gamma_n(s)), \int_{t_0}^s h_1(s, \tau, u(\gamma_{n+1}(\tau))) d\tau \right)$$
  
 
$$- F_1(s, v(\gamma_1(s)), ..., v(\gamma_n(s)), \int_{t_0}^s h_1(s, \tau, v(\gamma_{n+1}(\tau))) d\tau ) \left\| ds \right\|$$

$$+ \int_{t_0}^{t} || T(t-s) || \left\| F_2\left(s, u(\eta_1(s)), ..., u(\eta_m(s)), \int_{t_0}^{s} h_2(s, \tau, u(\eta_{m+1}(\tau))) d\tau\right) \right\|$$

$$- F_2\left(s, v(\eta_1(s)), ..., v(\eta_m(s)), \int_{t_0}^{s} h_2(s, \tau, v(\eta_{m+1}(\tau))) d\tau\right) \left\| ds$$

$$\leq MG || u - v ||_X + MM_1 \int_{t_0}^{t} \left( \sum_{i=1}^{n} \left\| u(\gamma_i(s)) - v(\gamma_i(s)) \right\| \right.$$

$$+ \int_{t_0}^{s} \left\| h_1(s, \tau, u(\gamma_{n+1}(\tau))) - h_1(s, \tau, v(\gamma_{n+1}(\tau))) \right\| d\tau \right) ds$$

$$+ MM_2 \int_{t_0}^{t} \left( \sum_{j=1}^{m} \left\| u(\eta_j(s)) - v(\eta_j(s)) \right\| \right.$$

$$+ \int_{t_0}^{s} \left\| h_2(s, \tau, u(\eta_{m+1}(\tau))) - h_2(s, \tau, v(\eta_{m+1}(\tau))) \right\| d\tau \right) ds$$

$$\leq MG || u - v ||_X + MM_1 b \Big[ n || u - v ||_X + H_1 b || u - v ||_X \Big]$$

$$+ MM_2 b \Big[ m || u - v ||_X + H_2 b || u - v ||_X \Big], \forall u, v \in J, t \in I.$$

Let  $\lambda = M [G + M_1 b(n + H_1 b) + M_2 b(m + H_2 b)]$ . Then, by (7) and by assumption (6), we have  $\| (\phi u)(t) - (\phi v)(t) \| \leq \lambda \| u - v \|_X$ , for  $u, v \in X$ 

with  $0 < \lambda < 1$ . This shows that the operator  $\phi$  is a contraction on X.

## 3.2. Existence of Classical Solution

In this section, we shall study the existence of classical solution through the following theorem.

#### **Theorem 3.2.** Suppose that

- (i) The assumptions (i) and (iv) of Theorem 3.1 holds.
- (ii) E is reflexive Banach space,  $x_0 \in D(A)$  and  $g(x) \in D(A)$ , where x reveals the unique mild solution of system (1) (2).
- (iii) There are constants  $M_1 > 0, M_2 > 0$  in such way that

$$\| F_1(t, x_1, x_2, ..., x_{n+1}) - F_1(s, \tilde{x}_1, \tilde{x}_2, ..., \tilde{x}_{n+1}) \| \le M_1 \left[ |t-s| + \sum_{i=1}^{n+1} \| x_i - \tilde{x}_i \| \right]$$

for  $t, s \in J, x_i, \tilde{x}_i \in E, i = 1, 2, ..., n + 1$ and

$$\| F_2(t, x_1, x_2, ..., x_{m+1}) - F_2(s, \tilde{x}_1, \tilde{x}_2, ..., \tilde{x}_{m+1}) \| \le M_2 \left[ |t - s| + \sum_{j=1}^{m+1} \| x_j - \tilde{x}_j \| \right] \le J, x_j, \tilde{x}_j \in E, j = 1, 2, ..., m + 1;$$

(iv) There exist constants  $H_1, H_2 > 0$  such that

$$|| h_1(t_1, s, x) - h_1(t_2, s, \tilde{x})|| \le H_1[|t_1 - t_2| + || x - \tilde{x}||]$$

and

for t, s

$$|| h_2(t_1, s, x) - h_2(t_2, s, \tilde{x})|| \le H_2[|t_1 - t_2| + || x - \tilde{x}||]$$

(v) There are constants  $C_3 > 0, C_4 > 0$  such that

 $|| x(\gamma_i(t)) - x(\gamma_i(s))|| \le C_3 || x(t) - x(s)||, i = 1, 2, ..., n + 1$ 

and

$$|| x(\eta_j(t)) - x(\eta_j(s))|| \le C_4 || x(t) - x(s)||, j = 1, 2, ..., m + 1$$

for  $t, s \in J$ .

If all the above assumptions are fulfil, then x is the unique classical solution of the system (1) - (2) on J.

*Proof.* The equation (1) - (2) possess a unique mild solution if all the conditions of Theorem 3.1 are satisfied, which is represented by x.

Next, we want to manifest that x is the unique classical solution of (1) - (2) on J. At this stage, we initiate

$$C_{5} = \max_{s \in J} \left\| F_{1}\left(s, x(\gamma_{1}(s)), ..., x(\gamma_{n}(s)), \int_{t_{0}}^{s} h_{1}(s, \tau, x(\gamma_{n+1}(\tau))) d\tau\right) \right\|,$$
  

$$C_{6} = \max_{s \in J} \left\| F_{2}\left(s, x(\eta_{1}(s)), ..., x(\eta_{m}(s)), \int_{t_{0}}^{s} h_{2}(s, \tau, x(\eta_{m+1}(\tau))) d\tau\right) \right\|,$$
  

$$C_{7} = \max_{t,s \in \Delta} \left\| h_{1}(t, s, x(\gamma_{n+1}(s))) \right\|, \text{and } C_{8} = \max_{t,s \in \Delta} \left\| h_{2}(t, s, x(\eta_{n+1}(s))) \right\|.$$

For this purpose, consider the difference

$$\begin{split} & x(t+h) - x(t) \\ &= \left[ T(t+h-t_0)x_0 - T(t-t_0)x_0 \right] - \left[ T(t+h-t_0)g(x) - T(t-t_0)g(x) \right] \\ &+ \int_{t_0}^{t_0+h} T(t+h-s)F_1\left(s,x(\gamma_1(s)),...,x(\gamma_n(s)),\int_{t_0}^s h_1(s,\tau,x(\gamma_{n+1}(\tau)))d\tau \right) ds \\ &+ \int_{t_0+h}^{t+h} T(t+h-s)F_1\left(s,x(\gamma_1(s)),...,x(\gamma_n(s)),\int_{t_0}^s h_1(s,\tau,x(\gamma_{n+1}(\tau)))d\tau \right) ds \\ &- \int_{t_0}^t T(t-s)F_1\left(s,x(\gamma_1(s)),...,x(\gamma_n(s)),\int_{t_0}^s h_1(s,\tau,x(\gamma_{n+1}(\tau)))d\tau \right) ds \\ &+ \int_{t_0}^{t_0+h} T(t+h-s)F_2\left(s,x(\eta_1(s)),...,x(\eta_m(s)),\int_{t_0}^s h_2(s,\tau,x(\eta_{m+1}(\tau)))d\tau \right) ds \\ &+ \int_{t_0+h}^{t+h} T(t+h-s)F_2\left(s,x(\eta_1(s)),...,x(\eta_m(s)),\int_{t_0}^s h_2(s,\tau,x(\eta_{m+1}(\tau)))d\tau \right) ds \\ &- \int_{t_0}^t T(t-s)F_2\left(s,x(\eta_1(s)),...,x(\eta_m(s)),\int_{t_0}^s h_2(s,\tau,x(\eta_{m+1}(\tau)))d\tau \right) ds \\ &= T(t-t_0)[T(h)-T]x_0 - T(t-t_0)[T(h)-T]g(x) \\ &+ \int_{t_0}^{t_0+h} T(t+h-s)F_1\left(s,x(\gamma_1(s)),...,x(\gamma_n(s)),\int_{t_0}^s h_1(s,\tau,x(\gamma_{n+1}(\tau)))d\tau \right) ds \\ &+ \int_{t_0}^t T(t-s) \times \\ &\left[ F_1\left(s+h,x(\gamma_1(s+h)),...,x(\gamma_n(s+h)),\int_{t_0}^{s+h} h_1(s+h,\tau,x(\gamma_{n+1}(\tau)))d\tau \right) ds \\ &+ \int_{t_0}^{t_0+h} T(t+h-s)F_2\left(s,x(\eta_1(s)),...,x(\eta_m(s)),\int_{t_0}^s h_2(s,\tau,x(\eta_{m+1}(\tau)))d\tau \right) ds \\ &+ \int_{t_0}^{t_0+h} T(t+h-s)F_2\left(s,x(\eta_1(s)),...,x(\eta_m(s)),\int_{t_0}^{s+h} h_1(s+h,\tau,x(\gamma_{n+1}(\tau)))d\tau \right) ds \\ &+ \int_{t_0}^{t_0+h} T(t+h-s)F_2\left(s,x(\eta_1(s)),...,x(\eta_m(s)),\int_{t_0}^{s+h} h_2(s,\tau,x(\eta_{m+1}(\tau)))d\tau \right) ds \\ &+ \int_{t_0}^{t_0+h} T(t+h-s)F_2\left(s,x(\eta_1(s)),...,x(\eta_m(s)),\int_{t_0}^{s} h_2(s,\tau,x(\eta_{m+1}(\tau)))d\tau \right) ds \\ &+ \int_{t_0}^{t_0+h} T(t+h-s)F_2\left(s,x(\eta_1(s)),....,x(\eta_m$$

$$\begin{split} &+ \int_{t_0}^t T(t-s) \times \\ &\left[ F_2 \Big( s+h, x(\eta_1(s+h)), ..., x(\eta_m(s+h)), \int_{t_0}^{s+h} h_2(s+h, \tau, x(\eta_{m+1}(\tau))) d\tau \Big) \right] \\ &- F_2 \Big( s, x(\eta_1(s)), ..., x(\eta_m(s)), \int_{t_0}^s h_2(s, \tau, x(\eta_{m+1}(\tau))) d\tau \Big) \right] ds \\ &\leq Mh \| Ax_0 \| + Mh \| Ag(x) \| + hMC_5 \\ &+ \int_{t_0}^t MM_1 \left[ h + \sum_{i=1}^n \left\| x(\gamma_i(s+h)) - x(\gamma_i(s)) \right\| + \int_{t_0}^s H_1 |s+h-s| d\tau \\ &+ \int_s^{s+h} C_7 d\tau \right] ds + hMC_6 + \int_{t_0}^t MM_2 \left[ h + \sum_{j=1}^m \left\| x(\eta_j(s+h)) - x(\eta_j(s)) \right\| \\ &+ \int_s^s H_2 |s+h-s| d\tau + \int_s^{s+h} C_8 d\tau \right] ds \\ &\leq Mh \| Ax_0 \| + Mh \| Ag(x) \| + hMC_5 + MM_1 bh \\ &+ MM_1 \int_{t_0}^t \sum_{i=1}^n \| x(\gamma_i(s+h)) - x(\gamma_i(s)) \| ds + MM_1 hbH_1 + MM_1 C_7 hb \end{split}$$

for  $t \in [t_0, t_0 + h), h > 0$  and  $t + h \in (t_0, t_0 + b]$ , where

$$Q := M \left[ \| Ax_0 \| + \| Ag(x) \| + C_5 + M_1 b(1 + H_1) + M_1 C_7 b + C_6 + M_2 b(1 + H_2) + M_2 C_8 b \right]$$

With the use of Gronwall's inequality and use of (8), we have

$$||x(t+h) - x(t)|| \le Qh \exp^{bM[M_1nC_3 + M_2mC_4]}$$

for  $t \in [t_0, t_0 + h), h > 0$  and  $t + h \in (t_0, t_0 + b]$ . Hence, x is Lipschitz continuous on J.

The Lipschitz continuity of x on J and inequalities (4), (5) imply that the function

$$\begin{aligned} t \in J \to Z(t) &:= F_1\Big(t, x(\gamma_1(t)), ..., x(\gamma_n(t)), \int_{t_0}^t h_1(t, s, x(\gamma_{n+1}(t)))ds\Big) \\ &+ F_2\Big(t, x(\eta_1(t)), ..., x(\eta_m(t)), \int_{t_0}^t h_2(t, s, x(\eta_{m+1}(t)))ds\Big) \in E \end{aligned}$$

is Lipschitz continuous on J. This property of  $t \to z(t)$  along with assumptions of Theorem 3.2 suggested by Theorem 1 given in of [12] and by Theorem 3.1 together with equation (3), we conclude that the linear Cauchy problem

$$v'(t) + Av(t) = z(t), t \in J/\{t_0\}$$
  
 $v(t_0) = x_0 - g(x)$ 

has a unique classical solution v in such a manner

$$v(t) = T(t - t_0)x_0 - T(t - t_0)g(x) + \int_{t_0}^t T(t - s)z(s)ds$$

$$= T(t - t_0)x_0 - T(t - t_0)g(x) + \int_{t_0}^t T(t - s)F_1\Big(s, x(\gamma_1(s)), \dots, x(\gamma_n(s)), \int_{t_0}^s h_1(s, \tau, x(\gamma_{n+1}(\tau)))d\tau\Big)ds + \int_{t_0}^t T(t - s)F_2\Big(s, x(\eta_1(s)), \dots, x(\eta_m(s)), \int_{t_0}^s h_2(s, \tau, x(\eta_{m+1}(\tau)))d\tau\Big)ds = x(t), t \in J$$

As a result, x(t) is the unique classical solution of the initial value problem (1) - (2) on J. This completes the proof of Theorem 3.2.

### 3.3. Applications

Now, we discuss two examples in favour of our results.

(1) We assume the following partial integrodifferential equation of the form:

$$\frac{\partial z(t,x)}{\partial t} - \frac{\partial^2 z(t,x)}{\partial x^2} = f_1 \Big( t, z(\gamma_1(t), x), \dots, z(\gamma_n(t), x), \int_{t_0}^t H_1(t, s, z(\gamma_{n+1}(s), x)) ds \Big) \\
+ f_2 \Big( t, z(\eta_1(t), x), \dots, z(\eta_m(t), x), \int_{t_0}^t H_2(t, s, z(\eta_{m+1}(s), x)) ds \Big), \\
0 < x < \pi, t \ge 0,$$
(9)

with initial and boundary conditions

$$z(0,t) = z(\pi,t) = 0, t \ge 0$$
(10)

$$z_0(x) = z(t_0, x) + \sum_{p=1}^k c_p z(t_p, x), x \in [0, \pi].$$
(11)

In continuation  $E = L^2[0,\pi]$  and  $A: D(A) \subset E \to E$  is the operator Az = z'' with domain  $D(A) = \{z \in E: z, z' \text{ are absolutely continuous, } z'' \in E, z(0) = z(\pi) = 0\}.$ 

It is well known that A is the infinitesimal generator of  $C_0$ -semigroup  $\{T(t)\}_{t\geq 0}$  on E. It is assumed that for certain constants  $N_i > 0, i = 1, 2, 3, 4, 5$ , the following conditions are satisfied:

$$\| f_1(t, y_1, y_2, ..., y_{n+1}) - f_1(t, z_1, z_2, ..., z_{n+1}) \| \le N_1 \sum_{i=1}^{n+1} \| y_i - z_i \|$$
  
$$\| f_2(t, y_1, y_2, ..., y_{m+1}) - f_2(t, z_1, z_2, ..., z_{m+1}) \| \le N_2 \sum_{j=1}^{m+1} \| y_j - z_j \|$$
  
$$\| H_1(t, s, y) - H_1(t, s, z) \| \le N_3(\| y - z \|)$$
  
$$\| H_2(t, s, y) - H_2(t, s, z) \| \le N_4(\| y - z \|)$$
  
$$\| G(s_1) - G(s_2) \| \le N_5(\| s_1 - s_2 \|)$$

where  $(Gz)(x) = \sum_{p=1}^{k} c_p z(t_p, x).$ 

Define the function  $F_1: J \times E^{n+1} \to E; \quad F_2: J \times E^{m+1} \to E; \quad h_1, h_2: J \times J \times E \to E \text{ and } G: X \to E$  as follows

$$F_1(t, x_1(t), \dots, x_{n+1}(t))(x) = f_1(t, x_1(x, t), \dots, x_{n+1}(x, t))$$

$$F_2(t, x_1(t), \dots, x_{m+1}(t))(x) = f_2(t, x_1(x, t), \dots, x_{m+1}(x, t))$$

$$h_1(t, s, x_1(t))(x) = H_1(t, s, x_1(x, t))$$

$$h_2(t, s, x_1(t))(x) = H_2(t, s, x_1(x, t))$$

for  $t \in J$  and  $0 < x < \pi$ . Then the above problem (9) -(11) can be formulated in (1) -(2). Since all the hypothesis of Theorem 3.1 are satisfied. Consequently, Theorem 3.1 can be applied for the equations (9) -(11).

(2) Consider the another partial integrodifferential equation of the form:

$$\frac{\partial w(t,y)}{\partial t} - \frac{\partial^2 w(t,y)}{\partial y^2} = c_1(t)w(\sin t,y) + c_2(t)\sin w(t,y) + \frac{1}{t^2+1}\int_{t_0}^t c_3(s)w(\sin s,y)ds + \tilde{c}_1(t)w(\sin t,y) + \tilde{c}_2(t)\sin w(t,y) + \frac{1}{t^2+1}\int_{t_0}^t \tilde{c}_3(s)w(\sin s,y)ds,$$
(12)

$$w(t,0) = w(t,\pi) = 0;$$
(13)

$$w(0,y) + \sum_{p=1}^{k} C_p w(t_p, y) = w_0(y)$$
(14)

where we state the conditions as follows:

(i) The function  $c_j(.)$  and  $\tilde{c}_j(.), j = 1, 2, 3$  are continuous on [0, 1] with condition

$$l_j = \sup_{0 \le s \le 1} ||c_j(s)| < 1, j = 1, 2, 3$$

and

$$\tilde{l}_j = \sup_{0 \le s \le 1} \|\tilde{c}_j(s)\| < 1, j = 1, 2, 3$$

(b) The function  $C_p \in R, p = 1, 2, ..., k$ .

Let us consider that  $E = L^2[0, \pi]$ . Explain  $A = D(A) \subset E \to E$  is linear operator which is described by Aw = w'' with domain  $D(A) = \{w \in E : w, w' \text{ are absolutely continuous, } w'' \in E, w(0) = w(\pi) = 0\}$ Then operator A can be expressed

$$Aw = \sum_{n=1}^{\infty} n^2(w, w_n)w_n, w \in D(A)$$

where  $w_n(y) = \left(\frac{\sqrt{2}}{\pi}\right) \sin ny$ , n = 1, 2, ... is the orthogonal set of eigenvalues of A. Further, for  $w \in E$ , we have

$$T(t)w = \sum_{n=1}^{\infty} exp\left(\frac{-n^2t}{1+n^2}\right)(w, w_n)w_n.$$

It common that A is the infinitesimal generator of  $C_0$ -semigroup  $\{T(t)\}_{\geq 0}$  on E.

To solve this system, we will define the operators  $F_1, F_2: J \times E \times E \to E; h_1, h_2: J \times J \times E \to E; g: X \to E$  by

$$\begin{split} F_1\bigg(t, w(\alpha(t)), \int_0^t h_1(t, s, w(\alpha(t)))ds\bigg)(y) \\ &= c_1(t)w(\sin t, y) + c_2(t)\sin w(t, y) + \frac{1}{t^2 + 1} \int_{t_0}^t c_3(s)w(\sin s, y)ds; \\ F_2\bigg(t, w(\alpha(t)), \int_0^t h_2(t, s, w(\alpha(t)))ds\bigg)(y) \\ &= \tilde{c}_1(t)w(\sin t, y) + \tilde{c}_2(t)\sin w(t, y) + \frac{1}{t^2 + 1} \int_{t_0}^t \tilde{c}_3(s)w(\sin s, y)ds; \\ \int_0^t h_1(t, s, w(\alpha(t)))ds &= \frac{1}{t^2 + 1} \int_{t_0}^t c_3(s)w(\sin s, y)ds; \\ \int_0^t h_2(t, s, w(\alpha(t)))ds &= \frac{1}{t^2 + 1} \int_{t_0}^t \tilde{c}_3(s)w(\sin s, y)ds \\ g(w)(y) &= \sum_{p=1}^k t_p w(t_p, y) \end{split}$$

Then system (12) - (14) yields the abstract form (1) - (2). With the choice of the above functions it is clear that all the conditions of the Theorem 3.1 are fulfilled. Thus with the help of Theorem 3.1, we assume that the system (12) - (14) has a mild solution on J.

#### 4. Acknowledgement

We are thankful to the referee for his valuable suggestions and comments on the manuscript for its improvement.

#### References

- Hamdy M. Ahmed, Boundary controllability of impulsive nonlinear fractional delay integro-differential system, Cogent Engineering 3:1, DOI: 10.1080/23311916.2016.1215766.
- [2] H. Akca, V. Covachev and Z. Covacheva, Existence theorem for a second-order impulsive functional-differential equation with a nonlocal condition, J. Nonlinear Convex Anal. 17 (2016), no. 6, 1129–1136.
- [3] K. Balachandran and E.R. Anandhi, Boundary controllability of delay integrodifferential systems in Banach spaces, J. Korean Soc. Ind. Appl. Math. 4 (2000), no. 2, 67–75.
- [4] K. Balachandran and M. Chandrasekaran, Existence of solutions of delay differential equation with nonlocal condition, Indian J. Pure Appl. Math. 27 (1996), no. 5, 443–449.
- [5] K. Balachandran and R.R. Kumar, Existence of solutions of integrodifferential evolution equations with time varying delays, Appl. Math. E-Notes 7 (2007), 1-8.
- [6] L. Byszewski, Theorems about the existence and uniqueness of solutions of a semilinear evolution nonlocal Cauchy problem, J. Math. Anal. Appl. 162 (1991), no. 1, 496-505.
- [7] X. Fu and X. Liu, Existence of solutions for neutral non-autonomous evolution equations with nonlocal conditions, Indian J. Pure Appl. Math. 37 (2006), no. 3, 179–192.
- [8] H. Gabsia, A. Ardjounib and A. Djoudic, Existence of periodic solutions for two types of second-order nonlinear neutral integro-differential equations with infinite distributed mixed-delays, Advances in the Theory of Nonlinear Analysis and its Applications 2 (2018) No. 4, 184–194
- K. Kumar and R. Kumar, Controllability results for general integrodifferential evolution equations in Banach space, Differ. Uravn. Protsessy Upr. 2015 (2015), no. 3, 1-15.
- [10] D.G. Park, K. Balachandran and F.P. Samuel, Regularity of solutions of abstract quasilinear delay integrodifferential equations, J. Korean Math. Soc. 48 (2011), no. 3, 585–597.
- [11] A. Pazy, Semigroup of Linear Operators and Applications to Partial Differential Equations, Springer-Verlag, New York, 1983.
- [12] T. Winirska, Nonlinear evolution equation with parameter, Bull. Pol. Acad. Sci. Math. 37 (1989), 157–162.
- [13] S. Xie, Existence of solutions for nonlinear mixed type integro-differential functional evolution equations with nonlocal conditions, Bound. Value Probl. 2012 (2012), 100. https://doi.org/10.1186/1687-2770-2012-100.
- [14] Z. Yan, Existence for a nonlinear impulsive functional integrodifferential equation with nonlocal conditions in Banach spaces, J. Appl. Math. Inform. 29 (2011), no. 3-4, 681-696.