

PAPER DETAILS

TITLE: SOME NEW SIMPSON TYPE INEQUALITIES FOR THE p -CONVEX AND p -CONCAVE FUNCTIONS

AUTHORS: İmdat İSCAN, Nihan Kalyoncu KONUK, Mahir KADAKAL

PAGES: 252-263

ORIGINAL PDF URL: <https://dergipark.org.tr/tr/download/article-file/594484>



SOME NEW SIMPSON TYPE INEQUALITIES FOR THE p -CONVEX AND p -CONCAVE FUNCTIONS

İMDAT İŞCAN, NİHAN KALYONCU KONUK, AND MAHİR KADAKAL

ABSTRACT. In this paper, we establish some new Simpson type inequalities for the class of functions whose derivatives in absolute values at certain powers are p -convex and p -concave.

1. INTRODUCTION

A function $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ is said to be convex if the inequality

$$f(tx + (1-t)y) \leq tf(x) + (1-t)f(y)$$

is valid for all $x, y \in I$ and $t \in [0, 1]$. If this inequality reverses, then f is said to be concave on interval $I \neq \emptyset$. This definition is well known in the literature.

It is well known that theory of convex sets and convex functions play an important role in mathematics and the other pure and applied sciences. In recent years, the concept of convexity has been extended and generalized in various directions using novel and innovative techniques. For some inequalities, generalizations and applications concerning convexity see [1, 2, 4, 5, 6, 16, 20].

In [9], the author gave definition harmonically convex and concave functions as follow.

Definition 1. Let $I \subset \mathbb{R} \setminus \{0\}$ be a real interval. A function $f : I \rightarrow \mathbb{R}$ is said to be harmonically convex, if

$$f\left(\frac{xy}{tx + (1-t)y}\right) \leq tf(y) + (1-t)f(x)$$

for all $x, y \in I$ and $t \in [0, 1]$. If this inequality is reversed, then f is said to be harmonically concave.

Received by the editors: June 21, 2017, Accepted: July 21, 2017.

2010 *Mathematics Subject Classification.* 26A51, 26D10, 26D15.

Key words and phrases. Convex function, p -Convex function, Simpson type inequalities.

©2018 Ankara University.

Communications Faculty of Sciences University of Ankara-Series A1 Mathematics and Statistics.
Communications de la Faculté des Sciences de l'Université d'Ankara-Séries A1 Mathématiques et Statistiques.

Definition 2. Let $I \subset (0, \infty)$ be a real interval and $p \in \mathbb{R} \setminus \{0\}$. A function $f : I \rightarrow \mathbb{R}$ is said to be a p -convex function, if

$$f\left([tx^p + (1-t)y^p]^{1/p}\right) \leq tf(x) + (1-t)f(y)$$

for all $x, y \in I$ and $t \in [0, 1]$. If this inequality is reversed, then f is said to be p -concave.

According to Definition 2, It can be easily seen that for $p = 1$ and $p = -1$, p -convexity reduces to ordinary convexity and harmonically convexity of functions defined on $I \subset (0, \infty)$, respectively.

Hermite-Hadamard inequality for the p -convex function is following:

Theorem 1. Let $f : I \subset (0, \infty) \rightarrow \mathbb{R}$ be a p -convex function, $p \in \mathbb{R} \setminus \{0\}$, and $a, b \in I$ with $a < b$. If $f \in L[a, b]$ then we have

$$f\left(\left[\frac{a^p + b^p}{2}\right]^{\frac{1}{p}}\right) \leq \frac{p}{b^p - a^p} \int_a^b \frac{f(x)}{x^{1-p}} dx \leq \frac{f(a) + f(b)}{2}.$$

These inequalities are sharp [5, 8]. If these inequalities are reversed, then f is said to be p -concave.

Many papers have been written by a number of mathematicians concerning inequalities for different classes of harmonically convex and p -convex functions see for instance the recent papers [3, 7, 8, 9, 10, 11, 12, 17, 18, 19, 21, 22, 24] and the references within these papers.

The following integral inequality, named Simpson's integral inequality, is one of the best known results in the literature.

Theorem 2. (Simpson's Integral Inequality). Let $f : I = [a, b] \subset \mathbb{R} \rightarrow \mathbb{R}$ be a four time continuously differentiable on I° , where I° is the interior of I and $\|f^{(4)}\|_\infty < \infty$. Then

$$\left| \frac{1}{3} \left[\frac{f(a) + f(b)}{2} + 2f\left(\frac{a+b}{2}\right) \right] - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{1}{2880} \|f^{(4)}\|_\infty (b-a)^4.$$

There are substantial literature on Simpson type integral inequalities. Here we mention the result of [13, 14, 15] and the corresponding references cited therein.

Throughout this paper we will use the following notations. Let $0 < a < b$ and $p \in \mathbb{R} \setminus \{0\}$.

$$\begin{aligned} A_p &= A_p(a, b) = \frac{a^p + b^p}{2}, \quad A_1 = A = A(a, b) = \frac{a + b}{2} \\ M_p &= M_p(a, b) = A_p^{\frac{1}{p}} = \left[\frac{a^p + b^p}{2} \right]^{\frac{1}{p}}, \quad H = H(a, b) = \frac{2ab}{a + b} \\ I_t(x, A_p; u, v) &= \frac{|t - \frac{1}{3}|^u}{[(1-t)x^p + tA_p]^{v - \frac{v}{p}}}, \\ J_t(x, A_p; u, v) &= \frac{|t - \frac{1}{3}|^u (1-t)}{[(1-t)x^p + tA_p]^{v - \frac{v}{p}}}, \\ K_t(x, A_p; u, v) &= \frac{|t - \frac{1}{3}|^u t}{[(1-t)x^p + tA_p]^{v - \frac{v}{p}}}. \end{aligned}$$

where $t \in [0, 1]$ and $u, v \geq 0$.

2. MAIN RESULTS

In this section, we will use the following Lemma for we obtain the main results:

Lemma 1. *Let $f : I \subset (0, \infty) \rightarrow \mathbb{R}$ be a differentiable mapping on I° (interior of I) and $a, b \in I^\circ$ with $a < b$ and $p \in \mathbb{R} \setminus \{0\}$. If $f' \in L[a, b]$, then the following equality holds:*

$$\begin{aligned} & \frac{1}{6} [f(a) + 4f(M_p) + f(b)] - \frac{p}{b^p - a^p} \int_a^b \frac{f(x)}{x^{1-p}} dx \\ &= \frac{b^p - a^p}{4p} \left[\int_0^1 \frac{t - \frac{1}{3}}{[(1-t)a^p + tA_p]^{1 - \frac{1}{p}}} f' \left([(1-t)a^p + tA_p]^{\frac{1}{p}} \right) dt \right. \\ & \quad \left. + \int_0^1 \frac{t - \frac{2}{3}}{[(1-t)A_p + tb^p]^{1 - \frac{1}{p}}} f' \left([(1-t)A_p + tb^p]^{\frac{1}{p}} \right) dt \right]. \end{aligned}$$

Proof. Firstly, let's calculate the following integral:

$$\begin{aligned} & \frac{b^p - a^p}{4p} \left[\int_0^1 \frac{t - \frac{1}{3}}{[(1-t)a^p + tA_p]^{1 - \frac{1}{p}}} f' \left([(1-t)a^p + tA_p]^{\frac{1}{p}} \right) dt \right. \\ & \quad \left. + \int_0^1 \frac{t - \frac{2}{3}}{[(1-t)A_p + tb^p]^{1 - \frac{1}{p}}} f' \left([(1-t)A_p + tb^p]^{\frac{1}{p}} \right) dt \right]. \end{aligned}$$

For shortness, we will use the notations

$$\begin{aligned} I_1 &= \int_0^1 \left(t - \frac{1}{3}\right) df \left([(1-t) a^p + t A_p]^{\frac{1}{p}}\right), \\ I_2 &= \int_0^1 \left(t - \frac{2}{3}\right) df \left([(1-t) A_p + t b^p]^{\frac{1}{p}}\right). \end{aligned}$$

Using the partial integration method and the method of changing variables respectively for the integrals I_1 and I_2 as following, we get

$$\begin{aligned} I_1 &= \int_0^1 \left(t - \frac{1}{3}\right) df \left([(1-t) a^p + t A_p]^{\frac{1}{p}}\right) \\ &= \left(t - \frac{1}{3}\right) f \left([(1-t) a^p + t A_p]^{\frac{1}{p}}\right) \Big|_0^1 - \int_0^1 f \left([(1-t) a^p + t A_p]^{\frac{1}{p}}\right) dt \\ &= \frac{2}{3} f(M_p) + \frac{1}{3} f(a) - \frac{2p}{b^p - a^p} \int_a^{A_p} \frac{f(x)}{x^{1-p}} dx \end{aligned} \quad (2.1)$$

$$\begin{aligned} I_2 &= \int_0^1 \left(t - \frac{2}{3}\right) df \left([(1-t) A_p + t b^p]^{\frac{1}{p}}\right) \\ &= \left(t - \frac{2}{3}\right) f \left([(1-t) A_p + t b^p]^{\frac{1}{p}}\right) \Big|_0^1 - \int_0^1 f \left([(1-t) A_p + t b^p]^{\frac{1}{p}}\right) dt \\ &= \frac{1}{3} f(b) + \frac{2}{3} f(M_p) - \frac{2p}{b^p - a^p} \int_{A_p}^b \frac{f(x)}{x^{1-p}} dx. \end{aligned} \quad (2.2)$$

Summing up side by side (2.1) and (2.2), we have

$$\begin{aligned} I_1 + I_2 &= \frac{1}{3} [f(a) + f(b)] + \frac{4}{3} f(M_p) - \frac{2p}{b^p - a^p} \int_a^b \frac{f(x)}{x^{1-p}} dx, \\ \frac{I_1 + I_2}{2} &= \frac{1}{6} [f(a) + 4f(M_p) + f(b)] - \frac{p}{b^p - a^p} \int_a^b \frac{f(x)}{x^{1-p}} dx. \end{aligned}$$

□

Theorem 3. Let $f : I \subset (0, \infty) \rightarrow \mathbb{R}$ be a differentiable mapping on I° (the interior of I) and $a, b \in I^\circ$ with $a < b$ and $p \in \mathbb{R} \setminus \{0\}$. If $f' \in L[a, b]$ and $|f'|^q$ is p -convex on I for $q \geq 1$, then the following inequality holds:

$$\begin{aligned} &\left| \frac{1}{6} [f(a) + 4f(M_p) + f(b)] - \frac{p}{b^p - a^p} \int_a^b \frac{f(x)}{x^{1-p}} dx \right| \\ &\leq \frac{b^p - a^p}{4p} [C_p(a, b)]^{1-\frac{1}{q}} [|f'(a)|^q D_p(a, b) + |f'(M_p)|^q E_p(a, b)]^{\frac{1}{q}} \\ &\quad + \frac{b^p - a^p}{4p} [F_p(a, b)]^{1-\frac{1}{q}} [|f'(M_p)|^q G_p(a, b) + |f'(b)|^q H_p(a, b)]^{\frac{1}{q}} \end{aligned}$$

where

$$\begin{aligned} C_p(a, b) &= \int_0^1 I_t(a, A_p; 1, 1) dt, & D_p(a, b) &= \int_0^1 J_t(a, A_p; 1, 1) dt, \\ E_p(a, b) &= \int_0^1 K_t(a, A_p; 1, 1) dt, & F_p(a, b) &= \int_0^1 I_{1-t}(b, A_p; 1, 1) dt, \\ G_p(a, b) &= \int_0^1 K_{1-t}(b, A_p; 1, 1) dt, & H_p(a, b) &= \int_0^1 J_{1-t}(b, A_p; 1, 1) dt. \end{aligned}$$

Proof. Using Lemma 1 and the power mean inequality, we have

$$\begin{aligned} & \left| \frac{1}{6} [f(a) + 4f(M_p) + f(b)] - \frac{p}{b^p - a^p} \int_a^b \frac{f(x)}{x^{1-p}} dx \right| \\ & \leq \frac{b^p - a^p}{4p} \left[\int_0^1 \frac{|t - \frac{1}{3}|}{[(1-t)a^p + tA_p]^{1-\frac{1}{p}}} \left| f' \left([(1-t)a^p + tA_p]^{\frac{1}{p}} \right) \right| dt \right] \\ & \quad + \frac{b^p - a^p}{4p} \left[\int_0^1 \frac{|t - \frac{2}{3}|}{[(1-t)A_p + tb^p]^{1-\frac{1}{p}}} \left| f' \left([(1-t)A_p + tb^p]^{\frac{1}{p}} \right) \right| dt \right] \\ & \leq \frac{b^p - a^p}{4p} \left(\int_0^1 \frac{|t - \frac{1}{3}|}{[(1-t)a^p + tA_p]^{1-\frac{1}{p}}} dt \right)^{1-\frac{1}{q}} \\ & \quad \times \left(\int_0^1 \frac{|t - \frac{1}{3}|}{[(1-t)a^p + tA_p]^{1-\frac{1}{p}}} \left| f' \left([(1-t)a^p + tA_p]^{\frac{1}{p}} \right) \right|^q dt \right)^{\frac{1}{q}} \\ & \quad + \frac{b^p - a^p}{4p} \left(\int_0^1 \frac{|t - \frac{2}{3}|}{[(1-t)A_p + tb^p]^{1-\frac{1}{p}}} dt \right)^{1-\frac{1}{q}} \\ & \quad \times \left(\int_0^1 \frac{|t - \frac{2}{3}|}{[(1-t)A_p + tb^p]^{1-\frac{1}{p}}} \left| f' \left([(1-t)A_p + tb^p]^{\frac{1}{p}} \right) \right|^q dt \right)^{\frac{1}{q}} \\ & \leq \frac{b^p - a^p}{4p} \left(\int_0^1 I_t(a, A_p; 1, 1) dt \right)^{1-\frac{1}{q}} \\ & \quad \times \left(\int_0^1 \frac{|t - \frac{1}{3}| [(1-t)|f'(a)|^q + t|f'(M_p)|^q]}{[(1-t)a^p + tA_p]^{1-\frac{1}{p}}} dt \right)^{\frac{1}{q}} \\ & \quad + \frac{b^p - a^p}{4p} \left(\int_0^1 I_{1-t}(b, A_p; 1, 1) dt \right)^{1-\frac{1}{q}} \\ & \quad \times \left(\int_0^1 \frac{|t - \frac{2}{3}| [(1-t)|f'(M_p)|^q + t|f'(b)|^q]}{[(1-t)A_p + tb^p]^{1-\frac{1}{p}}} dt \right)^{\frac{1}{q}} \end{aligned}$$

$$\begin{aligned}
 &= \frac{b^p - a^p}{4p} \left[\int_0^1 I_t(a, A_p; 1, 1) dt \right]^{1-\frac{1}{q}} \\
 &\quad \times \left[|f'(a)|^q \int_0^1 J_t(a, A_p; 1, 1) dt + |f'(M_p)|^q \int_0^1 K_t(a, A_p; 1, 1) dt \right]^{\frac{1}{q}} \\
 &\quad + \frac{b^p - a^p}{4p} \left[\int_0^1 I_{1-t}(b, A_p; 1, 1) dt \right]^{1-\frac{1}{q}} \\
 &\quad \times \left[|f'(M_p)|^q \int_0^1 K_{1-t}(b, A_p; 1, 1) dt + |f'(b)|^q \int_0^1 J_{1-t}(b, A_p; 1, 1) dt \right]^{\frac{1}{q}} \\
 &\leq \frac{b^p - a^p}{4p} [C_p(a, b)]^{1-\frac{1}{q}} [|f'(a)|^q D_p(a, b) + |f'(M_p)|^q E_p(a, b)]^{\frac{1}{q}} \\
 &\quad + \frac{b^p - a^p}{4p} [F_p(a, b)]^{1-\frac{1}{q}} [|f'(M_p)|^q G_p(a, b) + |f'(b)|^q H_p(a, b)]^{\frac{1}{q}}.
 \end{aligned}$$

This completes the proof of theorem. $\square$

Corollary 1. Under conditions of Theorem 3

i. If we take $p = 1$, then we obtain the following inequality for convex function:

$$\begin{aligned}
 &\left| \frac{1}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] - \frac{1}{b-a} \int_a^b f(x) dx \right| \\
 &\leq \frac{b-a}{4} \left(\frac{5}{18}\right)^{1-\frac{1}{q}} \left[\frac{8}{81} |f'(a)|^q + \frac{29}{162} \left| f'\left(\frac{a+b}{2}\right) \right|^q \right]^{\frac{1}{q}} \\
 &\quad + \frac{b-a}{4} \left(\frac{5}{18}\right)^{1-\frac{1}{q}} \left[\frac{29}{162} \left| f'\left(\frac{a+b}{2}\right) \right|^q + \frac{8}{81} |f'(b)|^q \right]^{\frac{1}{q}}
 \end{aligned}$$

which is the same of the inequality [6, Corollary 10] for $s = 1$.

ii. If we take $p = -1$, then we obtain the following inequality for harmonically convex function:

$$\begin{aligned}
 &\left| \frac{1}{6} \left[f(a) + 4f\left(\frac{2ab}{a+b}\right) + f(b) \right] - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \right| \\
 &\leq \frac{b-a}{4ab} [C_{-1}(a, b)]^{1-\frac{1}{q}} \left[|f'(a)|^q D_{-1}(a, b) + \left| f'\left(\frac{2ab}{a+b}\right) \right|^q E_{-1}(a, b) \right]^{\frac{1}{q}} \\
 &\quad + \frac{b-a}{4ab} [F_{-1}(a, b)]^{1-\frac{1}{q}} \left[\left| f'\left(\frac{2ab}{a+b}\right) \right|^q G_{-1}(a, b) + |f'(b)|^q H_{-1}(a, b) \right]^{\frac{1}{q}}.
 \end{aligned}$$

Theorem 4. Let $f : I \subset (0, \infty) \rightarrow \mathbb{R}$ be a differentiable mapping on I° (the interior of I) and $a, b \in I^\circ$ with $a < b$ and $p \in \mathbb{R} \setminus \{0\}$. If $f' \in L[a, b]$ and $|f'|^q$ is p -convex

on I for $q > 1$, $\frac{1}{r} + \frac{1}{q} = 1$, then

$$\begin{aligned} & \left| \frac{1}{6} [f(a) + 4f(M_p) + f(b)] - \frac{p}{b^p - a^p} \int_a^b \frac{f(x)}{x^{1-p}} dx \right| \\ & \leq \frac{b^p - a^p}{4p} \left[N_{p,r}^{\frac{1}{r}}(a, b) A^{\frac{1}{q}}(|f'(a)|^q, |f'(M_p)|^q) + O_{p,r}^{\frac{1}{r}}(a, b) A^{\frac{1}{q}}(|f'(M_p)|^q, |f'(b)|^q) \right], \end{aligned}$$

where

$$\begin{aligned} N_{p,r}(a, b) &= \int_0^1 I_t(a, A_p; r, r) dt, \\ O_{p,r}(a, b) &= \int_0^1 I_{1-t}(b, A_p; r, r) dt. \end{aligned}$$

Proof. From Lemma 1, Hölder's integral inequality and the p -convexity of $|f'|^q$ on $[a, b]$, we have,

$$\begin{aligned} & \left| \frac{1}{6} [f(a) + 4f(M_p) + f(b)] - \frac{p}{b^p - a^p} \int_a^b \frac{f(x)}{x^{1-p}} dx \right| \\ & \leq \frac{b^p - a^p}{4p} N_{p,r}^{\frac{1}{r}}(a, b) \left(\int_0^1 \left| f' \left(((1-t)a^p + tA_p)^{\frac{1}{p}} \right) \right|^q dt \right)^{\frac{1}{q}} \\ & \quad + \frac{b^p - a^p}{4p} O_{p,r}^{\frac{1}{r}}(a, b) \left(\int_0^1 \left| f' \left(((1-t)A_p + tb^p)^{\frac{1}{p}} \right) \right|^q dt \right)^{\frac{1}{q}} \\ & \leq \frac{b^p - a^p}{4p} \left[N_{p,r}^{\frac{1}{r}}(a, b) \left(\int_0^1 ((1-t)|f'(a)|^q + t|f'(M_p)|^q) dt \right)^{\frac{1}{q}} \right. \\ & \quad \left. + O_{p,r}^{\frac{1}{r}}(a, b) \left(\int_0^1 ((1-t)|f'(M_p)|^q + t|f'(b)|^q) dt \right)^{\frac{1}{q}} \right] \\ & = \frac{b^p - a^p}{4p} \left[N_{p,r}^{\frac{1}{r}}(a, b) \left(\frac{|f'(a)|^q + |f'(M_p)|^q}{2} \right) + O_{p,r}^{\frac{1}{r}}(a, b) \left(\frac{|f'(M_p)|^q + |f'(b)|^q}{2} \right) \right] \\ & = \frac{b^p - a^p}{4p} \left[N_{p,r}^{\frac{1}{r}}(a, b) M_q(|f'(a)|, |f'(M_p)|) + O_{p,r}^{\frac{1}{r}}(a, b) M_q(|f'(M_p)|, |f'(b)|) \right]. \end{aligned}$$

This completes the proof of theorem. $\square$

Corollary 2. Under conditions of Theorem 4,

i. If we take $p = 1$, then we obtain the following inequality for convex function:

$$\begin{aligned} & \left| \frac{1}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ & \leq \frac{b-a}{12} \left[\frac{1+2^{r+1}}{3(r+1)} \right]^{\frac{1}{r}} \left[M_q \left(|f'(a)|, \left| f' \left(\frac{a+b}{2} \right) \right| \right) + M_q \left(\left| f' \left(\frac{a+b}{2} \right) \right|, |f'(b)| \right) \right] \end{aligned}$$

which is the same as the inequality in [22, Theorem 8] for $s = 1$.

ii. If we take $p = -1$, then we obtain the following inequality for harmonically convex function:

$$\begin{aligned} & \left| \frac{1}{6} \left[f(a) + 4f\left(\frac{2ab}{a+b}\right) + f(b) \right] - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \right| \\ & \leq \frac{b-a}{4ab} \left[N_{-1,r}^{\frac{1}{r}}(a,b) M_q(|f'(a)|, |f'(H)|) + O_{-1,r}^{\frac{1}{r}}(a,b) M_q(|f'(H)|, |f'(b)|) \right]. \end{aligned}$$

Theorem 5. Let $f : I \subset (0, \infty) \rightarrow \mathbb{R}$ be a differentiable mapping on I° (the interior of I) and $a, b \in I^\circ$ with $a < b$ and $p \in \mathbb{R} \setminus \{0\}$. If $f' \in L[a, b]$ and $|f'|^q$ is p -convex on I for $q > 1$, $\frac{1}{r} + \frac{1}{q} = 1$, then

$$\begin{aligned} & \left| \frac{1}{6} [f(a) + 4f(M_p) + f(b)] - \frac{p}{b^p - a^p} \int_a^b \frac{f(x)}{x^{1-p}} dx \right| \\ & \leq \frac{b^p - a^p}{12p} \left[\frac{1 + 2^{r+1}}{3(r+1)} \right]^{\frac{1}{r}} \left\{ [Q_{p,q}(a,b) |f'(a)|^q + R_{p,q}(a,b) |f'(M_p)|^q]^{\frac{1}{q}} \right. \\ & \quad \left. + [S_{p,q}(a,b) |f'(b)|^q + T_{p,q}(a,b) |f'(M_p)|^q]^{\frac{1}{q}} \right\}, \end{aligned}$$

where

$$\begin{aligned} Q_{p,q}(a,b) &= \int_0^1 J_t(a, A_p; 0, q) dt, & S_{p,q}(a,b) &= \int_0^1 K_{1-t}(b, A_p; 0, q) dt, \\ R_{p,q}(a,b) &= \int_0^1 K_t(a, A_p; 0, q) dt, & T_{p,q}(a,b) &= \int_0^1 J_{1-t}(b, A_p; 0, q) dt. \end{aligned}$$

Proof. From Lemma 1, Hölder's integral inequality and the p -convexity of $|f'|^q$ on $[a, b]$, we obtain,

$$\begin{aligned}
& \left| \frac{1}{6} [f(a) + 4f(M_p) + f(b)] - \frac{p}{b^p - a^p} \int_a^b \frac{f(x)}{x^{1-p}} dx \right| \\
& \leq \frac{b^p - a^p}{4p} \int_0^1 \left| t - \frac{1}{3} \right| \left| \frac{1}{[(1-t)a^p + tA_p]^{1-\frac{1}{p}}} f' \left([(1-t)a^p + tA_p]^{\frac{1}{p}} \right) \right| dt \\
& \quad + \frac{b^p - a^p}{4p} \int_0^1 \left| t - \frac{2}{3} \right| \left| \frac{1}{[(1-t)A_p + tb^p]^{1-\frac{1}{p}}} f' \left([(1-t)A_p + tb^p]^{\frac{1}{p}} \right) \right| dt \\
& \leq \frac{b^p - a^p}{12p} \left[\frac{1 + 2^{r+1}}{3(r+1)} \right]^{\frac{1}{r}} \left(\int_0^1 \frac{1}{[(1-t)a^p + tA_p]^{q-\frac{q}{p}}} \left| f' \left([(1-t)a^p + tM_p^p]^{\frac{1}{p}} \right) \right|^q dt \right)^{\frac{1}{q}} \\
& \quad + \frac{b^p - a^p}{12p} \left[\frac{1 + 2^{r+1}}{3(r+1)} \right]^{\frac{1}{r}} \left(\int_0^1 \frac{1}{[(1-t)A_p + tb^p]^{q-\frac{q}{p}}} \left| f' \left([(1-t)M_p^p + tb^p]^{\frac{1}{p}} \right) \right|^q dt \right)^{\frac{1}{q}} \\
& \leq \frac{b^p - a^p}{12p} \left[\frac{1 + 2^{r+1}}{3(r+1)} \right]^{\frac{1}{r}} \left\{ [Q_{p,q}(a, b) |f'(a)|^q + R_{p,q}(a, b) |f'(M_p)|^q]^{\frac{1}{q}} \right. \\
& \quad \left. + [S_{p,q}(a, b) |f'(b)|^q + T_{p,q}(a, b) |f'(M_p)|^q]^{\frac{1}{q}} \right\}.
\end{aligned}$$

This completes the proof of theorem. $\square$

Corollary 3. Under conditions of Theorem 5,

i. If we take $p = 1$, then we obtain the following inequality for convex function:

$$\begin{aligned}
& \left| \frac{1}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] - \frac{1}{b-a} \int_a^b f(x) dx \right| \\
& \leq \frac{b-a}{12} \left[\frac{1 + 2^{r+1}}{3(r+1)} \right]^{\frac{1}{r}} \left[M_q \left(|f'(a)|, \left| f' \left(\frac{a+b}{2} \right) \right| \right) + M_q \left(\left| f' \left(\frac{a+b}{2} \right) \right|, |f'(b)| \right) \right]
\end{aligned}$$

which reduce the inequality in Corollary 2 (i).

ii. If we take $p = -1$, then we obtain the following inequality for harmonically convex function:

$$\begin{aligned}
& \left| \frac{1}{6} \left[f(a) + 4f\left(\frac{2ab}{a+b}\right) + f(b) \right] - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \right| \\
& \leq \frac{b-a}{12ab} \left[\frac{1 + 2^{r+1}}{3(r+1)} \right]^{\frac{1}{r}} \left\{ [Q_{-1,q}(a, b) |f'(a)|^q + R_{-1,q}(a, b) |f'(H)|^q]^{\frac{1}{q}} \right. \\
& \quad \left. + [S_{-1,q}(a, b) |f'(b)|^q + T_{-1,q}(a, b) |f'(H)|^q]^{\frac{1}{q}} \right\}.
\end{aligned}$$

Theorem 6. Let $f : I \subset (0, \infty) \rightarrow \mathbb{R}$ be a differentiable mapping on I° (the interior of I) and $a, b \in I^\circ$ with $a < b$ and $p \in \mathbb{R} \setminus \{0\}$. If $f' \in L[a, b]$ and $|f'|^q$ is p -concave on I for $q > 1$, $\frac{1}{r} + \frac{1}{q} = 1$, then

$$\begin{aligned} & \left| \frac{1}{6} [f(a) + 4f(M_p) + f(b)] - \frac{p}{b^p - a^p} \int_a^b \frac{f(x)}{x^{1-p}} dx \right| \\ & \leq \frac{b^p - a^p}{4p} \left[N_{p,r}^{\frac{1}{r}}(a, b) \left| f' \left(\left[\frac{3a^p + b^p}{4} \right]^{\frac{1}{p}} \right) \right| + O_{p,r}^{\frac{1}{r}}(a, b) \left| f' \left(\left[\frac{a^p + 3b^p}{4} \right]^{\frac{1}{p}} \right) \right| \right]. \end{aligned}$$

Proof. From Lemma 1, Hölder's integral inequality and the p -concavity of $|f'|^q$ on $[a, b]$, we have,

$$\begin{aligned} & \left| \frac{1}{6} [f(a) + 4f(M_p) + f(b)] - \frac{p}{b^p - a^p} \int_a^b \frac{f(x)}{x^{1-p}} dx \right| \\ & \leq \frac{b^p - a^p}{4p} N_{p,r}^{\frac{1}{r}}(a, b) \left(\int_0^1 \left| f' \left(((1-t)a^p + tA_p)^{\frac{1}{p}} \right) \right|^q dt \right)^{\frac{1}{q}} \\ & \quad + \frac{b^p - a^p}{4p} O_{p,r}^{\frac{1}{r}}(a, b) \left(\int_0^1 \left| f' \left(((1-t)A_p + tb^p)^{\frac{1}{p}} \right) \right|^q dt \right)^{\frac{1}{q}} \\ & \leq \frac{b^p - a^p}{4p} \left[N_{p,r}^{\frac{1}{r}}(a, b) \left| f' \left(\left[\frac{3a^p + b^p}{4} \right]^{\frac{1}{p}} \right) \right| + O_{p,r}^{\frac{1}{r}}(a, b) \left| f' \left(\left[\frac{a^p + 3b^p}{4} \right]^{\frac{1}{p}} \right) \right| \right]. \end{aligned}$$

This completes the proof of theorem. $\square$

Corollary 4. Under conditions of Theorem 6,

i. If we take $p = 1$, then we obtain the following inequality for concave function:

$$\begin{aligned} & \left| \frac{1}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] - \frac{1}{b-a} \int_a^b f(x) dx \right| \\ & \leq \frac{b-a}{12} \left[\frac{1+2^{r+1}}{3(r+1)} \right]^{\frac{1}{r}} \left[\left| f' \left(\frac{3a+b}{4} \right) \right| + \left| f' \left(\frac{a+3b}{4} \right) \right| \right] \end{aligned}$$

This is the same of the inequality in [6, Corollary 28] for $s = 1$.

ii. If we take $p = -1$, then we obtain the following inequality for harmonically concave function:

$$\begin{aligned} & \left| \frac{1}{6} \left[f(a) + 4f\left(\frac{2ab}{a+b}\right) + f(b) \right] - \frac{ab}{b-a} \int_a^b \frac{f(x)}{x^2} dx \right| \\ & \leq \frac{b-a}{4ab} \left[N_{-1,r}^{\frac{1}{r}}(a, b) \left| f' \left(\frac{4ab}{a+3b} \right) \right| + O_{-1,r}^{\frac{1}{r}}(a, b) \left| f' \left(\frac{4ab}{3a+b} \right) \right| \right]. \end{aligned}$$

3. CONCLUSION

The paper deals with Simpson type inequalities for p -convex and p -concave functions. Firstly, we give a new identity for differentiable functions and get some new integral inequalities for the p -convex and p -concave functions by using this identity.

REFERENCES

- [1] Anderson, G.D., Vamanamurthy M.K. and Vuorinen, M., Generalized convexity and inequalities, *Journal of Mathematical Analysis and Applications*, 335(2) (2007), 1294-1308.
- [2] Avcı, M., Kavurmacı H. and Özdemir, M. E., New inequalities of Hermite-Hadamard type via s -convex functions in the second sense with applications, *Appl. Math. Comput.*, 217 (2011), 5171-5176
- [3] I. A. Baloch and İ. İşcan, Some Ostrowski Type Inequalities for Harmonically (s, m) -convex functions in Second Sense, *International Journal of Analysis*, vol. 2015 (2015), Article ID 672675, 9 pages, <http://dx.doi.org/10.1155/2015/672675>.
- [4] Dragomir S.S. and Agarwal, R.P., Two Inequalities for Differentiable Mappings and Applications to Special Means of Real Numbers and to Trapezoidal Formula, *Appl. Math. Lett.* 11(5) (1998), 91-95.
- [5] Fang Z. B. and Shi, R., On the $(p; h)$ -convex function and some integral inequalities, *J. Inequal. Appl.*, 2014(45) (2014), 16 pages.
- [6] İşcan, İ., New estimates on generalization of some integral inequalities for s -convex functions and their applications, *International Journal of Pure and Applied Mathematics*, 86(4) (2013), 727-746.
- [7] İşcan, İ., Hermite-Hadamard Type Inequalities for p -Convex Functions, *International Journal of Analysis and Applications*, Volume 11, Number 2 (2016), 137-145.
- [8] İşcan, İ., Ostrowski type inequalities for p -convex functions, *New Trends in Mathematical Sciences*, NTMSCI 4, No. 3, 140-150 (2016).
- [9] İşcan, İ., Hermite-Hadamard type inequalities for harmonically convex functions, *Hacettepe Journal of Mathematics and Statistics*, 43(6) (2014), 935-942.
- [10] İşcan, İ., M. Kunt, Hermite-Hadamard-Fejer Type Inequalities for Harmonically s -convex Functions via Fractional Integrals, *Australian Journal of Mathematical Analysis and Applications*, 12(1) (2015), Article 10, 1-16.
- [11] İşcan, İ., M. Aydın, S. Dikmenoglu, New integral inequalities via harmonically convex functions, *Mathematics and Statistics*, 3(5): 134-140, 2015. DOI: 10.13189/ms.2015.030504.
- [12] İşcan, İ., On generalization of different type inequalities for harmonically quasi-convex functions via fractional integrals, *Applied Mathematics and Computation*, 275 (2016) 287-298.
- [13] İşcan, İ., Hermite-Hadamard and Simpson-like type inequalities for differentiable harmonically convex functions, *Journal of Mathematics*, Volume 2014 (2014), Article ID 346305, 10 pages.
- [14] İşcan, İ., K. Bekar and S. Numan, Hermite-Hadamard and Simpson type inequalities for differentiable quasi-geometrically convex functions, *Turkish Journal of Analysis and Number Theory*, 2014, vol. 2, no. 2, 42-46.
- [15] İşcan, İ., S. Numan and K. Bekar, Hermite-Hadamard and Simpson type inequalities for differentiable harmonically p -functions, *British Journal of Mathematics & Computer Science*, 4(14) (2014), 1908-1920.
- [16] Kırmacı, U.S., Inequalities for differentiable mappings and applications to special means of real numbers and to midpoint formula, *Appl. Math. Comput.* 147 (2004), 137-146.
- [17] Kunt, M. İşcan, İ., Hermite-Hadamard-Fejer type inequalities for p -convex functions, *Arab Journal of Mathematical Sciences*, vol.23, pp.215-230, 2017.

- [18] Kunt, M. İscan, İ., Hermite-Hadamard-Fejer type inequalities for p -convex functions via fractional integrals, *Communication in Mathematical Modeling and Applications*, vol.2, no.1, pp.1-15, 2017.
- [19] Kunt, M. İscan, İ., Hermite-Hadamard type inequalities for p -convex functions via fractional integrals, *Moroccan Journal of Pure and Applied Analysis*, vol.3, no.1, pp.22-35, 2017.
- [20] Niculescu, C. P., Convexity according to the geometric mean, *Math. Inequal. Appl.*, 3(2) (2000), 155-167.
- [21] Noor, M. A., Noor, K. I., Iftikhar, S., Nonconvex Functions and Integral Inequalities, *Punjab University Journal of Mathematics*, Vol. 47(2)(2015), 19-27.
- [22] Noor, M. A., Noor, K. I., Mihai, M. V. and Awan, M. U., Hermite-Hadamard inequalities for differentiable p -convex functions using hyper geometric functions, *Publications De L'Institut Mathématique, Nouvelle série*, tome 100(114) (2016), 251-257.
- [23] Sarikaya, M.Z., Set, E. and Özdemir, M.E., On new inequalities of Simpson's type for s -convex functions, *Computers and Mathematics with Applications* 60 (2010), 2191-2199.
- [24] Zhang K.S. and Wan, J.P., p -convex functions and their properties, *Pure Appl. Math.* 23(1), (2007), 130-133.

Current address: İMDAT İŞCAN: Department of Mathematics, Faculty of Arts and Sciences, Giresun University, 28100, Giresun, Turkey.

E-mail address: imdat.iscan@giresun.edu.tr

ORCID Address: <http://orcid.org/0000-0001-6749-0591>

Current address: NİHAN KALYONCU KONUK: Institute of Sciences and Arts, Giresun University-Giresun-Turkey

E-mail address: nihankalyoncu@hotmail.com

ORCID Address: <http://orcid.org/0000-0003-1287-3370>

Current address: MAHİR KADAKAL: Department of Mathematics, Faculty of Arts and Sciences, Giresun University, 28100, Giresun, Turkey.

E-mail address: mahirkadakal@gmail.com

ORCID Address: <http://orcid.org/0000-0002-0240-918X>