

## PAPER DETAILS

TITLE: The mean remaining strength of parallel systems in a stress-strength model based on exponential distribution

AUTHORS: Fatih KIZILASLAN

PAGES: 1435-1451

ORIGINAL PDF URL: <https://dergipark.org.tr/tr/download/article-file/669562>



## THE MEAN REMAINING STRENGTH OF PARALLEL SYSTEMS IN A STRESS-STRENGTH MODEL BASED ON EXPONENTIAL DISTRIBUTION

FATİH KIZILASLAN

**ABSTRACT.** The mean remaining strength of any coherent system is one of the important characteristics in stress-strength reliability. It shows that the system on the average how long can be safe under the stress. In this paper, we consider the mean remaining strength of the parallel systems in the stress-strength model. We assume that the strength and stress components constitute parallel systems separately. The mean remaining strength and its estimations are obtained when the all components follow the exponential distribution. The likelihood ratio order between the remaining strength of the parallel systems is presented for two-component case. The simulation study is performed to compare the derived estimates and their results are presented.

### 1. INTRODUCTION

In the reliability theory, the stress-strength model describes the reliability of a component or system in terms of random variables. The reliability is defined as  $R = P(X > Y)$  where  $Y$  is the random stress experienced by the system and  $X$  is the random strength of the system available to overcome the stress. The system fails if the stress exceeds the strength. This main idea was introduced by Birnbaum [1] and developed by Birnbaum and McCarty [2]. The last few decades, the problem of estimating  $R$  has been considerable investigated by many authors for the different data types and the distributional assumptions on  $X$  and  $Y$ . Examples of such results and references can be found in Kotz et al. [3], Kundu and Gupta [4], Basirat et al. [5, 6], Asgharzadeh et al. [7]. However, some results in the multicomponent stress-strength models can be found in Bhattacharyya and Johnson [8, 9], Eryilmaz [10, 11], Pakdaman and Ahmadi [12, 13], Hassan [14], Kızılaslan [15].

Let  $X$  and  $Y$  be two independent random variables. It is assumed that  $X$  is the strength to failure of a component subject to a random stress  $Y$  and the component

---

Received by the editors: February 01, 2018; Accepted: October 03, 2018.

2010 *Mathematics Subject Classification.* Primary 62N05, 60K10; Secondary 60E15, 62N02.

*Key words and phrases.* Mean remaining strength, exponential and generalized exponential distributions, stress-strength model.

works if its strength is greater than the applied stress, that is  $X > Y$ . Then, we may estimate the component's survival function under the stress  $Y$ . We may also wish to learn for how long, on average, the component can still be safe under the stress. The mean remaining strength (MRS) of the component can be defined as the expected remaining strength under the stress  $Y$ , i.e.  $\Phi = E(X - Y | X > Y)$ .

The MRS of the systems has been presented by Gurler [16] for the simple stress-strength model,  $k$ -out-of- $n$  :  $F$  system, series and parallel systems under the common stress. When the component is alive at the strength level  $t$  under the applied stress  $Y$ , the MRS of the component was defined as  $\Phi(t) = E(X - Y - t | X - Y > t)$  for  $t > 0$  by Bairamov et al. [17]. They obtained that the MRS of the  $k$ -out-of- $n$  :  $F$  system, series and parallel systems for the exchangeable strength components under the common stress. The MRS of the two-component parallel and series systems were considered by Gurler et al. [18] for the dependent strength components which are subject to a common stress.

In this study, the parallel stress and strength systems are considered. It is assumed that  $X_1, \dots, X_{n_1}$  and  $Y_1, \dots, Y_{n_2}$  are independent and identical strength and stress random variables follow the exponential distribution with parameters  $\lambda_1$  and  $\lambda_2$ , respectively. Stochastic comparison of the remaining strength of two-component parallel strength and stress systems are studied. Maximum likelihood (ML) and Bayesian estimations of the MRS of this system are obtained. Bayesian estimates are derived by using Lindley's approximation and Markov Chain Monte Carlo (MCMC) method due to the lack of explicit forms. In Section 2, we introduce preliminaries for our system and obtain some distributional properties and stochastic ordering results. In Section 3, we derive ML and Bayesian estimations of the MRS of our system. Moreover, the asymptotic confidence and the highest probability density (HPD) credible intervals of the MRS are constructed. In Section 4, we present a simulation study to compare the proposed estimates of the MRS.

## 2. MODEL DESCRIPTION

Let  $X$  be a random variable with exponential distribution with parameter  $\lambda$  and mean  $1/\lambda$ . Then, it is known that the cdf and pdf of  $X$  are given by

$$F_X(x) = 1 - e^{-\lambda x}, \quad f_X(x) = \lambda e^{-\lambda x}, \quad x > 0, \quad \lambda > 0,$$

respectively and denoted by  $X \sim \text{Exp}(\lambda)$ .

For our case, it is assumed that  $X_1, \dots, X_{n_1}$  strength and  $Y_1, \dots, Y_{n_2}$  stress variables follow the exponential distribution with parameters  $\lambda_1$  and  $\lambda_2$ . It is known that the distribution of the parallel system (or its maximum) is generalized exponential (GE) or exponentiated exponential distribution when the components are independent and identical exponential distribution. The GE distribution was introduced by Gupta and Kundu [19]. This distribution has been studied extensively in the literature since then.

If we assume that  $X_{1:n_1} \leq X_{2:n_1} \leq \dots \leq X_{n_1:n_1}$  are the ordered strength of the components, then  $X_{1:n_1}$  and  $X_{n_1:n_1}$  are the weakest and strongest components. It is clear that the strength and stress of the parallel systems are  $\max_{1 \leq i \leq n_1} (X_i) = X_{n_1:n_1}$  and  $\max_{1 \leq i \leq n_2} (Y_i) = Y_{n_2:n_2}$ . The cdfs and pdfs of  $X_{n_1:n_1}$  and  $Y_{n_2:n_2}$  are

$$F_{X_{n_1:n_1}}(x) = (1 - e^{-\lambda_1 x})^{n_1}, \quad f_{X_{n_1:n_1}}(x) = n_1 \lambda_1 e^{-\lambda_1 x} (1 - e^{-\lambda_1 x})^{n_1-1},$$

and

$$F_{Y_{n_2:n_2}}(y) = (1 - e^{-\lambda_2 y})^{n_2}, \quad f_{Y_{n_2:n_2}}(y) = n_2 \lambda_2 e^{-\lambda_2 y} (1 - e^{-\lambda_2 y})^{n_2-1},$$

that is  $X_{n_1:n_1} \sim GE(n_1, \lambda_1)$  and  $Y_{n_2:n_2} \sim GE(n_2, \lambda_2)$  where  $n_i$  and  $\lambda_i$   $i = 1, 2$  are the shape and scale parameters.

In this case, the reliability for the strength and stress of the parallel systems is given by

$$\begin{aligned} R_{n_1, n_2} &= P(X_{n_1:n_1} > Y_{n_2:n_2}) = \int_0^\infty F_{Y_{n_2:n_2}}(x) f_{X_{n_1:n_1}}(x) dx \\ &= n_1 \sum_{i=0}^{n_1-1} \sum_{j=0}^{n_2} \binom{n_2}{j} \binom{n_1-1}{i} (-1)^{i+j} \frac{\lambda_1}{\lambda_1(i+1) + \lambda_2 j}. \end{aligned} \quad (1)$$

It is also obtained by Pakdaman and Ahmadi [12, 13] (see Equations 2.8 and 9, respectively).

Our system works if the strength is greater than the applied stress, that is  $X_{n_1:n_1} > Y_{n_2:n_2}$ . It is important to learn this system on the average how long can be safe under the stress. Hence, we want to estimate the mean remaining strength (MRS) of this system when the stress  $Y_{n_2:n_2}$  is applied. The MRS of our parallel systems are the expected remaining strength under the stress  $Y_{n_2:n_2}$  and given by

$$\Phi_{n_1, n_2} = E(X_{n_1:n_1} - Y_{n_2:n_2} | X_{n_1:n_1} > Y_{n_2:n_2}). \quad (2)$$

The cdf of the conditional random variable  $\psi \equiv (X_{n_1:n_1} - Y_{n_2:n_2} | X_{n_1:n_1} > Y_{n_2:n_2})$  is

$$\begin{aligned} F_\psi(x) &= P(X_{n_1:n_1} - Y_{n_2:n_2} \leq x | X_{n_1:n_1} > Y_{n_2:n_2}) \\ &= \frac{P(X_{n_1:n_1} \leq Y_{n_2:n_2} + x, X_{n_1:n_1} > Y_{n_2:n_2})}{P(X_{n_1:n_1} > Y_{n_2:n_2})} \\ &= \frac{P(X_{n_1:n_1} \leq Y_{n_2:n_2} + x, X_{n_1:n_1} > Y_{n_2:n_2})}{R_{n_1, n_2}}. \end{aligned}$$

Then, conditioning on  $Y_{n_2:n_2} = y$ ,

$$\begin{aligned} P(X_{n_1:n_1} \leq Y_{n_2:n_2} + x, X_{n_1:n_1} > Y_{n_2:n_2}) &= \int_0^\infty P(y < X_{n_1:n_1} \leq y + x) dF_{Y_{n_2:n_2}}(y) \\ &= \int_0^\infty (F_{X_{n_1:n_1}}(y + x) - F_{X_{n_1:n_1}}(y)) dF_{Y_{n_2:n_2}}(y) \end{aligned}$$

$$\begin{aligned}
&= \int_0^\infty F_{X_{n_1:n_1}}(y+x) dF_{Y_{n_2:n_2}}(y) - \int_0^\infty F_{X_{n_1:n_1}}(y) dF_{Y_{n_2:n_2}}(y) \\
&\equiv I_1 - I_2
\end{aligned}$$

and

$$\begin{aligned}
I_1 &= \int_0^\infty F_{X_{n_1:n_1}}(y+x) dF_{Y_{n_2:n_2}}(y) \\
&= \int_0^\infty (1 - e^{-\lambda_1(y+x)})^{n_1} n_2 \lambda_2 e^{-\lambda_2 y} (1 - e^{-\lambda_2 y})^{n_2-1} dy \\
&= n_2 \lambda_2 \sum_{i=0}^{n_1} \sum_{j=0}^{n_2-1} \binom{n_2-1}{j} \binom{n_1}{i} (-1)^{i+j} \frac{e^{-\lambda_1 i x}}{\lambda_1 i + \lambda_2 (j+1)}, \\
I_2 &= \int_0^\infty F_{X_{n_1:n_1}}(y) dF_{Y_{n_2:n_2}}(y) \\
&= \int_0^\infty (1 - e^{-\lambda_1 y})^{n_1} n_2 \lambda_2 e^{-\lambda_2 y} (1 - e^{-\lambda_2 y})^{n_2-1} dy \\
&= n_2 \lambda_2 \sum_{i=0}^{n_1} \sum_{j=0}^{n_2-1} \binom{n_2-1}{j} \binom{n_1}{i} (-1)^{i+j} \frac{1}{\lambda_1 i + \lambda_2 (j+1)}.
\end{aligned}$$

Hence,

$$F_\psi(x) = \frac{n_2 \lambda_2}{R_{n_1, n_2}} \sum_{i=1}^{n_1} \sum_{j=0}^{n_2-1} \binom{n_2-1}{j} \binom{n_1}{i} (-1)^{i+j} \frac{(e^{-\lambda_1 i x} - 1)}{\lambda_1 i + \lambda_2 (j+1)} \quad (3)$$

and

$$f_\psi(x) = \frac{dF_\psi(x)}{dx} = \frac{n_2 \lambda_2}{R_{n_1, n_2}} \sum_{i=1}^{n_1} \sum_{j=0}^{n_2-1} \binom{n_2-1}{j} \binom{n_1}{i} (-1)^{i+j+1} \frac{\lambda_1 i e^{-\lambda_1 i x}}{\lambda_1 i + \lambda_2 (j+1)}. \quad (4)$$

Then,

$$\begin{aligned}
\Phi_{n_1, n_2} &= E(X_{n_1:n_1} - Y_{n_2:n_2} | X_{n_1:n_1} > Y_{n_2:n_2}) \\
&= E_\psi(x) = \int_0^\infty x f_\psi(x) dx \\
&= \frac{n_2}{R_{n_1, n_2}} \sum_{i=1}^{n_1} \sum_{j=0}^{n_2-1} \binom{n_2-1}{j} \binom{n_1}{i} (-1)^{i+j+1} \frac{\lambda_2}{\lambda_1 i (\lambda_1 i + \lambda_2 (j+1))}. \quad (5)
\end{aligned}$$

It can be also rewritten as

$$\Phi_{n_1, n_2} = \frac{R_{n_1, n_2}^*}{R_{n_1, n_2}}, \quad (6)$$

where

$$R_{n_1, n_2}^* = n_2 \sum_{i=1}^{n_1} \sum_{j=0}^{n_2-1} \binom{n_2-1}{j} \binom{n_1}{i} (-1)^{i+j+1} \frac{\lambda_2}{\lambda_1 i (\lambda_1 i + \lambda_2 (j+1))}. \quad (7)$$

In Figure 1, some plots of  $\Phi_{n_1, n_2}$  with respect to the parameters  $\lambda_1$  and  $\lambda_2$  are presented. It is observed that  $\Phi_{n_1, n_2}$  is a decreasing function of  $\lambda_1$  for fixed value of  $\lambda_2$  and increasing function of  $\lambda_2$  for fixed value of  $\lambda_1$ .




FIGURE 1. Plots of  $\Phi_{n_1, n_2}$  with respect to the parameters  $\lambda_1$  and  $\lambda_2$ .

**2.1. Stochastic ordering results.** In this section, we present the likelihood ratio ordering result associated with the remaining strength of parallel systems i.e. the conditional random variable  $\psi \equiv X_{n_1:n_1} - Y_{n_2:n_2} | X_{n_1:n_1} > Y_{n_2:n_2}$ . This random variable is a special case of the residual life of a random variable  $X$  at random time  $\Theta$  which is defined as  $X_\Theta = X - \Theta | X > \Theta$  (see Dewan and Khaledi [20] and Misra and Naqvi [21, 22]).

Let  $X$  and  $Y$  be two lifetime random variables with pdfs  $f(x)$  and  $g(x)$ , respectively.  $X$  is said to be smaller than  $Y$  in the likelihood ratio order (denoted by  $X \leq_{lr} Y$ ) if  $g(x)/f(x)$  is increasing in  $x$  for all  $x$  for which this ratio is well defined. It is known that the likelihood ratio order implies other stochastic orders. Hence the likelihood ratio order is the most interesting order in stochastic comparison. For more details on stochastic comparisons, see Shaked and Shanthikumar [23].

The coefficients of the cdf and pdf of  $\psi \equiv X_{n_1:n_1} - Y_{n_2:n_2} | X_{n_1:n_1} > Y_{n_2:n_2}$  in equations (3) and (4) can be negative or positive. That is why general stochastic comparisons is not possible for  $\psi$  random variable. As a special case we consider two-component parallel systems (i.e.  $n_1 = n_2 = 2$ ). In this case, we have

$$f_\psi(x) = \frac{4\lambda_1\lambda_2^2}{R_{2,2}(\lambda_1 + \lambda_2)} \left[ \frac{e^{-\lambda_1 x}}{\lambda_1 + 2\lambda_2} - \frac{e^{-2\lambda_1 x}}{2(2\lambda_1 + \lambda_2)} \right],$$

where

$$R_{2,2} = 1 - \frac{5\lambda_1}{\lambda_1 + \lambda_2} + \frac{2\lambda_1}{\lambda_1 + 2\lambda_2} + \frac{4\lambda_1}{2\lambda_1 + \lambda_2},$$

from equations (4) and (1).

**Theorem 1.** Suppose  $X_i, X_i^*$   $i = 1, 2$  are the strength and  $Y_i, Y_i^*$   $i = 1, 2$  are the stress variables with  $X_i \sim \text{Exp}(\lambda_1)$ ,  $X_i^* \sim \text{Exp}(\lambda_1^*)$ ,  $Y_i \sim \text{Exp}(\lambda_2)$  and  $Y_i^* \sim \text{Exp}(\lambda_2^*)$ ,  $i = 1, 2$ . If  $\lambda_1^* < \lambda_1$  and  $\lambda_2 < \lambda_2^*$ , then we have  $\psi \leq_{lr} \psi^*$  where  $\psi = X_{2:2} - Y_{2:2} | X_{2:2} > Y_{2:2}$  and  $\psi^* = X_{2:2}^* - Y_{2:2}^* | X_{2:2}^* > Y_{2:2}^*$ .

*Proof.* If we show that  $f_{\psi^*}(x)/f_{\psi}(x)$  is increasing function in  $x$ , it completes the proof. Then, we have

$$\Lambda(x) = \frac{f_{\psi}(x)}{f_{\psi^*}(x)} = \frac{\lambda_1 \lambda_2^2}{\lambda_1^* \lambda_2^{*2}} \frac{R_{2,2}^* (\lambda_1^* + \lambda_2^*) D_1^* D_2^*}{R_{2,2} (\lambda_1 + \lambda_2) D_1 D_2} \frac{2D_1 e^{-\lambda_1 x} - D_2 e^{-2\lambda_1 x}}{2D_1^* e^{-\lambda_1^* x} - D_2^* e^{-2\lambda_1^* x}},$$

where  $D_1 = 2\lambda_1 + \lambda_2$ ,  $D_1^* = 2\lambda_1^* + \lambda_2^*$ ,  $D_2 = \lambda_1 + 2\lambda_2$  and  $D_2^* = \lambda_1^* + 2\lambda_2^*$ . After some computations

$$\begin{aligned} \Lambda'(x) &\stackrel{\text{sign}}{=} 4D_1 D_1^* e^{-\lambda_1 x} e^{-\lambda_1^* x} (\lambda_1^* - \lambda_1) + 2D_2 D_2^* e^{-2\lambda_1 x} e^{-2\lambda_1^* x} (\lambda_1^* - \lambda_1) \\ &\quad + 2D_1 D_2^* e^{-\lambda_1 x} e^{-2\lambda_1^* x} (\lambda_1 - 2\lambda_1^*) + 2D_1^* D_2 e^{-2\lambda_1 x} e^{-\lambda_1^* x} (2\lambda_1 - \lambda_1^*) \\ &< 2D_1 [2 D_1^* (\lambda_1^* - \lambda_1) + D_2^* (\lambda_1 - 2\lambda_1^*)] + 2 D_2 [D_2^* (\lambda_1^* - \lambda_1) + D_1^* (2\lambda_1 - \lambda_1^*)] \\ &= 6\lambda_1 \lambda_1^* [(\lambda_1^* - \lambda_1) + (\lambda_2 - \lambda_2^*)], \end{aligned}$$

where  $a \stackrel{\text{sign}}{=} b$  means that  $a$  and  $b$  have the same sign. The last inequality implies that  $\Lambda(x)$  is a decreasing function in  $x$  for  $\lambda_1^* < \lambda_1$  and  $\lambda_2 < \lambda_2^*$ . Hence, it completes the proof.  $\square$

**Example 2.** Theorem 1 results are observed in Figure 2. When the theorem conditions are not satisfied in Figure 3 (A) and (B), the graphic of  $f_{\psi^*}(x)/f_{\psi}(x)$  can be concave or convex. However, it is observed that all these results are also valid for  $n_1, n_2 > 2$ .

### 3. ESTIMATION OF $\Phi_{n_1, n_2}$

In this section, we consider the estimation problem of MRS. Although the estimation of the stress-strength reliability of different systems has been considered extensively, the similar problem for MRS has not been studied in the literature except for Gurler et al. [18]. In our case, ML and Bayes estimations of the MRS are studied.




FIGURE 2. Plot of  $f_{\psi^*}(x)/f_{\psi}(x)$  for  $(\lambda_1, \lambda_2) = (2, 3)$  and  $(\lambda_1^*, \lambda_2^*) = (1.5, 3.5)$ .




FIGURE 3. Plots of  $f_{\psi^*}(x)/f_{\psi}(x)$  for different parameters.

**3.1. MLE case.** The random strength and stress of the parallel systems are denoted by  $V = \max_{1 \leq i \leq n_1} (X_i)$  and  $W = \max_{1 \leq j \leq n_2} (Y_j)$ . It is known that  $V \sim GE(n_1, \lambda_1)$  and  $W \sim GE(n_2, \lambda_2)$  when  $X_i$ ,  $i = 1, \dots, n_1$  and  $Y_j$ ,  $j = 1, \dots, n_2$  are exponential distributions with parameters  $\lambda_1$  and  $\lambda_2$ . Let  $V_1, \dots, V_n$  be a random sample of size  $n$  from  $GE(n_1, \lambda_1)$  and  $W_1, \dots, W_m$  be a random sample of size  $m$  from  $GE(n_2, \lambda_2)$ . Then, the likelihood function based on the observed sample  $\{\mathbf{v} = (v_1, \dots, v_n), \mathbf{w} = (w_1, \dots, w_m)\}$  is given by

$$L(\lambda_1, \lambda_2 | \mathbf{v}, \mathbf{w}) = \prod_{i=1}^n \prod_{j=1}^m f_{V_i}(v_i) f_{W_j}(w_j)$$



$$= n_1^n \lambda_1^n n_2^m \lambda_2^m \exp \left( -\lambda_1 \sum_{i=1}^n v_i + (n_1 - 1) \sum_{i=1}^n \ln(1 - e^{-\lambda_1 v_i}) \right) \\ \exp \left( -\lambda_2 \sum_{j=1}^m w_j + (n_2 - 1) \sum_{j=1}^m \ln(1 - e^{-\lambda_2 w_j}) \right).$$

Hence, the MLEs of  $\lambda_1$  and  $\lambda_2$ , say  $\hat{\lambda}_1$  and  $\hat{\lambda}_2$ , are the solution of the following nonlinear equations:

$$\frac{n}{\lambda_1} - \sum_{i=1}^n \frac{v_i}{1 - e^{-\lambda_1 v_i}} + n_1 \sum_{i=1}^n \frac{v_i e^{-\lambda_1 v_i}}{1 - e^{-\lambda_1 v_i}} = 0, \\ \frac{m}{\lambda_2} - \sum_{j=1}^m \frac{w_j}{1 - e^{-\lambda_2 w_j}} + n_2 \sum_{j=1}^m \frac{w_j e^{-\lambda_2 w_j}}{1 - e^{-\lambda_2 w_j}} = 0.$$

$\hat{\lambda}_1$  and  $\hat{\lambda}_2$  can be obtained by using the fixed point method or Newton-Raphson method or other numerical methods. Ghitany et al. [24] proved that if at least one observation is different minimum of the all observations, then this type nonlinear equations have unique solution. When we obtain  $\hat{\lambda}_1$  and  $\hat{\lambda}_2$ , the MLE of  $\Phi_{n_1, n_2}$ , say  $\hat{\Phi}_{n_1, n_2}^{MLE}$ , is obtained from (5) by using the invariance property of MLEs.

Moreover, an asymptotic confidence interval of  $\Phi_{n_1, n_2}$  can be constructed based on the MLEs. The Fisher information matrix of  $\boldsymbol{\lambda} = (\lambda_1, \lambda_2)$  is

$$I(\boldsymbol{\lambda}) = - \begin{pmatrix} E \left( \frac{\partial^2 l}{\partial \lambda_1^2} \right) & E \left( \frac{\partial^2 l}{\partial \lambda_1 \partial \lambda_2} \right) \\ E \left( \frac{\partial^2 l}{\partial \lambda_2 \partial \lambda_1} \right) & E \left( \frac{\partial^2 l}{\partial \lambda_2^2} \right) \end{pmatrix} = \begin{pmatrix} I_{11} & I_{12} \\ I_{21} & I_{22} \end{pmatrix}.$$

where  $l = \ln(L(\lambda_1, \lambda_2 | v, \mathbf{w}))$ . The elements of the matrix are obtained as  $I_{12} = I_{21} = 0$ ,

$$I_{11} = \frac{n}{\lambda_1^2} + \frac{n_1 n}{\lambda_1^2 (n_1 - 2)} \left\{ (\psi(2) - \psi(n_1))^2 + \psi'(2) - \psi'(n_1) \right\}$$

and

$$I_{22} = \frac{m}{\lambda_2^2} + \frac{n_2 m}{\lambda_2^2 (n_2 - 2)} \left\{ (\psi(2) - \psi(n_2))^2 + \psi'(2) - \psi'(n_2) \right\},$$

for  $n_1 > 2$  and  $n_2 > 2$  by using the formula 4.261(17) in Gradshteyn and Ryzhik [25] where  $\psi(x) = d \ln \Gamma(x) / dx$  is a Psi function.  $\hat{\Phi}_{n_1, n_2}^{MLE}$  is asymptotically normal with mean  $\Phi_{n_1, n_2}$  and asymptotic variance

$$\sigma_{\Phi_{n_1, n_2}}^2 = \sum_{j=1}^2 \sum_{i=1}^2 \frac{\partial \Phi_{n_1, n_2}}{\partial \lambda_i} \frac{\partial \Phi_{n_1, n_2}}{\partial \lambda_j} I_{ij}^{-1},$$

where  $I_{ij}^{-1}$  is the  $(i, j)$ th element of the inverse of the  $I(\boldsymbol{\lambda})$ ; see Rao [26]. Then,

$$\sigma_{\Phi_{n_1, n_2}}^2 = \left( \frac{\partial \Phi_{n_1, n_2}}{\partial \lambda_1} \right)^2 \frac{1}{I_{11}} + \left( \frac{\partial \Phi_{n_1, n_2}}{\partial \lambda_2} \right)^2 \frac{1}{I_{22}}.$$

The partial derivatives of  $R_{n_1, n_2}$  and  $R_{n_1, n_2}^*$  with respect to  $\lambda_1$  and  $\lambda_2$  are given by

$$\begin{aligned}\frac{\partial R_{n_1, n_2}}{\partial \lambda_1} &= n_1 \sum_{i=0}^{n_1-1} \sum_{j=0}^{n_2} \binom{n_2}{j} \binom{n_1-1}{i} (-1)^{i+j} \frac{\lambda_2 j}{(\lambda_1(i+1) + \lambda_2 j)^2}, \\ \frac{\partial R_{n_1, n_2}}{\partial \lambda_2} &= n_1 \sum_{i=0}^{n_1-1} \sum_{j=0}^{n_2} \binom{n_2}{j} \binom{n_1-1}{i} (-1)^{i+j+1} \frac{\lambda_1 j}{(\lambda_1(i+1) + \lambda_2 j)^2}, \\ \frac{\partial R_{n_1, n_2}^*}{\partial \lambda_1} &= n_2 \sum_{i=1}^{n_1} \sum_{j=0}^{n_2-1} \binom{n_2-1}{j} \binom{n_1}{i} (-1)^{i+j} \frac{\lambda_2(2\lambda_1 i + \lambda_2(j+1))}{\lambda_1^2 i (\lambda_1 i + \lambda_2(j+1))^2}, \\ \frac{\partial R_{n_1, n_2}^*}{\partial \lambda_2} &= n_2 \sum_{i=1}^{n_1} \sum_{j=0}^{n_2-1} \binom{n_2-1}{j} \binom{n_1}{i} (-1)^{i+j+1} \frac{1}{(\lambda_1 i + \lambda_2(j+1))^2}.\end{aligned}$$

Then,  $\partial \Phi_{n_1, n_2} / \partial \lambda_1$  and  $\partial \Phi_{n_1, n_2} / \partial \lambda_2$  are evaluated by using these partial derivatives. Therefore, an asymptotic  $100(1 - \gamma)\%$  confidence interval of  $\Phi_{n_1, n_2}$  is given by

$$\Phi_{n_1, n_2} \in \left( \hat{\Phi}_{n_1, n_2}^{MLE} \pm z_{\gamma/2} \hat{\sigma}_{\Phi_{n_1, n_2}} \right),$$

where  $z_{\gamma/2}$  is the upper  $\gamma/2$ th quantile of the standard normal distribution and  $\hat{\sigma}_{\Phi_{n_1, n_2}}$  is the value of  $\sigma_{\Phi_{n_1, n_2}}$  at the MLE of the parameters.

**3.2. Bayesian case.** In this section, we assume that the parameters  $\lambda_1$  and  $\lambda_2$  are random variables and have statistically independent gamma prior distributions with parameters  $(a_i, b_i)$ ,  $i = 1, 2$ , respectively. The pdf of a gamma random variable  $X$  with parameters  $(a_i, b_i)$  is

$$f(x) = \frac{b_i^{a_i}}{\Gamma(a_i)} x^{a_i-1} e^{-x b_i}, \quad x > 0, \quad a_i, b_i > 0$$

where  $a_i, b_i > 0$ ,  $i = 1, 2$ . Then, the joint posterior density function of  $\lambda_1$  and  $\lambda_2$  is

$$\begin{aligned}\pi(\lambda_1, \lambda_2 | \mathbf{v}, \mathbf{w}) &\propto \lambda_1^{n+a_1-1} \exp \left( -\lambda_1 \left( b_1 + \sum_{i=1}^n v_i \right) + (n_1 - 1) \sum_{i=1}^n \ln(1 - e^{-\lambda_1 v_i}) \right) \\ &\quad \lambda_2^{m+a_2-1} \exp \left( -\lambda_2 \left( b_2 + \sum_{j=1}^m w_j \right) + (n_2 - 1) \sum_{j=1}^m \ln(1 - e^{-\lambda_2 w_j}) \right).\end{aligned}\tag{8}$$

The Bayes estimator of  $\Phi_{n_1, n_2}$  under the SE loss function is given by

$$\hat{\Phi}_{n_1, n_2}^{Bayes} = \int_0^\infty \int_0^\infty \Phi_{n_1, n_2} \pi(\lambda_1, \lambda_2 | \mathbf{v}, \mathbf{w}) d\lambda_1 d\lambda_2.\tag{9}$$

Since the integrals given in (9) is not computed analytically, Lindley's approximation and MCMC methods can be applied to approximate (9).

3.2.1. *Lindley's approximation.* Lindley [27] introduced an approximate procedure for the computation of the ratio of two integrals. This procedure, applied to the posterior expectation of the function  $U(\lambda)$  for a given  $\mathbf{x}$ , is

$$E(u(\theta) | \mathbf{x}) = \frac{\int u(\theta) e^{Q(\theta)} d\theta}{\int e^{Q(\theta)} d\theta},$$

where  $Q(\theta) = l(\theta) + \rho(\theta)$ ,  $l(\theta)$  is the logarithm of the likelihood function and  $\rho(\theta)$  is the logarithm of the prior density of  $\theta$ . Using Lindley's approximation,  $E(u(\theta) | \mathbf{x})$  is approximately estimated by

$$E(u(\theta) | \mathbf{x}) = \left[ u + \frac{1}{2} \sum_i \sum_j (u_{ij} + 2u_i \rho_j) \sigma_{ij} + \frac{1}{2} \sum_i \sum_j \sum_k \sum_l L_{ijk} \sigma_{ij} \sigma_{kl} u_l \right]_{\hat{\lambda}} \\ + \text{terms of order } n^{-2} \text{ or smaller,}$$

where  $\theta = (\theta_1, \theta_2, \dots, \theta_m)$ ,  $i, j, k, l = 1, \dots, m$ ,  $\hat{\theta}$  is the MLE of  $\theta$ ,  $u = u(\theta)$ ,  $u_i = \partial u / \partial \theta_i$ ,  $u_{ij} = \partial^2 u / \partial \theta_i \partial \theta_j$ ,  $L_{ijk} = \partial^3 l / \partial \theta_i \partial \theta_j \partial \theta_k$ ,  $\rho_j = \partial \rho / \partial \theta_j$  and  $\sigma_{ij} = (i, j)$ th element in the inverse of the matrix  $\{-L_{ij}\}$  all evaluated at the MLE of the parameters.

For the two parameter case  $\boldsymbol{\lambda} = (\lambda_1, \lambda_2)$ , Lindley's approximation leads to

$$\hat{u}_{Lin} = u(\boldsymbol{\lambda}) + \frac{1}{2} [B + Q_{30}B_{12} + Q_{21}C_{12} + Q_{12}C_{21} + Q_{03}B_{21}],$$

where  $B = \sum_{i=1}^2 \sum_{j=1}^2 u_{ij} \tau_{ij}$ ,  $Q_{ij} = \partial Q^{i+j} / \partial^i \theta \lambda_1 \partial^j \lambda_2$  for  $i, j = 0, 1, 2, 3$ ,  $i+j=3$ ,  $u_i = \partial u / \partial \lambda_i$ ,  $u_{ij} = \partial^2 u / \partial \lambda_i \partial \lambda_j$  for  $i, j = 1, 2$  and  $B_{ij} = (u_i \tau_{ii} + u_j \tau_{ij}) \tau_{ii}$ ,  $C_{ij} = 3u_i \tau_{ii} \tau_{ij} + u_j (\tau_{ii} \tau_{ij} + 2\tau_{ij}^2) \tau_{ij}$  for  $i \neq j$ .  $\tau_{ij}$  is the  $(i, j)$ th element in the inverse of matrix  $Q^* = (-Q_{ij}^*)$ ,  $i, j = 1, 2$  such that  $Q_{ij}^* = \partial Q^2 / \partial \lambda_i \partial \lambda_j$ . The approximate Bayes estimate  $\hat{u}_{Lin}$  is evaluated at  $(\tilde{\lambda}_1, \tilde{\lambda}_2)$  which is the mode of the posterior density.

In our case,  $u(\boldsymbol{\lambda}) = \Phi_{n_1, n_2}$ ,

$$Q = \ln \pi(\lambda_1, \lambda_2 | \mathbf{v}, \mathbf{w}) \propto (n + a_1 - 1) \ln \lambda_1 + (m + a_2 - 1) \ln \lambda_2 - \lambda_1 \left( b_1 + \sum_{i=1}^n v_i \right) \\ - \lambda_2 \left( b_2 + \sum_{j=1}^m w_j \right) + (n_1 - 1) \sum_{i=1}^n \ln(1 - e^{-\lambda_1 v_i}) + (n_2 - 1) \sum_{j=1}^m \ln(1 - e^{-\lambda_2 w_j}).$$

The posterior mode of  $\lambda_1$  and  $\lambda_2$ , say  $\tilde{\lambda}_1$  and  $\tilde{\lambda}_2$ , are the solution of the following nonlinear equations from  $Q$

$$\frac{n + a_1 - 1}{\lambda_1} - \left( b_1 + \sum_{i=1}^n v_i \right) + (n_1 - 1) \sum_{i=1}^n \frac{v_i e^{-\lambda_1 v_i}}{1 - e^{-\lambda_1 v_i}} = 0,$$

$$\frac{m + a_2 - 1}{\lambda_2} - \left( b_2 + \sum_{j=1}^m w_j \right) + (n_2 - 1) \sum_{j=1}^m \frac{w_j e^{-\lambda_2 w_j}}{1 - e^{-\lambda_2 w_j}} = 0.$$

Moreover, it is obtained that

$$\tau_{11} = \left[ \frac{n + a_1 - 1}{\lambda_1^2} + (n_1 - 1) \sum_{i=1}^n \frac{v_i^2 e^{-\lambda_1 v_i}}{(1 - e^{-\lambda_1 v_i})^2} \right]^{-1},$$

$$\tau_{22} = \left[ \frac{m + a_2 - 1}{\lambda_2^2} + (n_2 - 1) \sum_{j=1}^m \frac{w_j^2 e^{-\lambda_2 w_j}}{(1 - e^{-\lambda_2 w_j})^2} \right]^{-1},$$

$$\tau_{12} = \tau_{21} = 0, Q_{12} = Q_{21} = 0, Q_{03} = 2(m + a_2 - 1)/\lambda_2^3, Q_{30} = 2(n + a_1 - 1)/\lambda_1^3, \\ B_{12} = u_1 \tau_{11}^2, B_{21} = u_2 \tau_{22}^2, B = u_{11} \tau_{11} + u_{22} \tau_{22},$$

$$u_{11} = \frac{\partial^2 \Phi_{n_1, n_2}}{\partial \lambda_1^2} = \frac{1}{(R_{n_1, n_2})^2} \left( R_{n_1, n_2} \frac{\partial^2 R_{n_1, n_2}^*}{\partial \lambda_1^2} - R_{n_1, n_2}^* \frac{\partial^2 R_{n_1, n_2}}{\partial \lambda_1^2} \right) \\ - \frac{2}{(R_{n_1, n_2})^3} \frac{\partial R_{n_1, n_2}}{\partial \lambda_1} \left( R_{n_1, n_2} \frac{\partial R_{n_1, n_2}^*}{\partial \lambda_1} - R_{n_1, n_2}^* \frac{\partial R_{n_1, n_2}}{\partial \lambda_1} \right)$$

and

$$u_{22} = \frac{\partial^2 \Phi_{n_1, n_2}}{\partial \lambda_2^2} = \frac{1}{(R_{n_1, n_2})^2} \left( R_{n_1, n_2} \frac{\partial^2 R_{n_1, n_2}^*}{\partial \lambda_2^2} - R_{n_1, n_2}^* \frac{\partial^2 R_{n_1, n_2}}{\partial \lambda_2^2} \right) \\ - \frac{2}{(R_{n_1, n_2})^3} \frac{\partial R_{n_1, n_2}}{\partial \lambda_2} \left( R_{n_1, n_2} \frac{\partial R_{n_1, n_2}^*}{\partial \lambda_2} - R_{n_1, n_2}^* \frac{\partial R_{n_1, n_2}}{\partial \lambda_2} \right).$$

$u_{11}$  and  $u_{22}$  are evaluated by using the second partial derivatives of  $R_{n_1, n_2}$  and  $R_{n_1, n_2}^*$  with respect to  $\lambda_1$  and  $\lambda_2$ . Therefore, the approximate Bayes estimate of  $\Phi_{n_1, n_2}$  is

$$\hat{\Phi}_{n_1, n_2}^{Lin} = \Phi_{n_1, n_2} + \frac{1}{2} [B + Q_{30} B_{12} + Q_{03} B_{21}]_{(\lambda_1, \lambda_2) = (\tilde{\lambda}_1, \tilde{\lambda}_2)}. \quad (10)$$

**3.2.2. MCMC method.** The joint posterior density function of  $\lambda_1$  and  $\lambda_2$  is given in (8). The marginal posterior density functions of  $\lambda_1$  and  $\lambda_2$  are given respectively as

$$\pi_1(\lambda_1 | \lambda_2, \mathbf{v}, \mathbf{w}) \propto \lambda_1^{n+a_1-1} \exp \left( -\lambda_1 \left( b_1 + \sum_{i=1}^n v_i \right) + (n_1 - 1) \sum_{i=1}^n \ln(1 - e^{-\lambda_1 v_i}) \right),$$

and

$$\pi_2(\lambda_2 | \lambda_1, \mathbf{v}, \mathbf{w}) \propto \lambda_2^{m+a_2-1} \exp \left( -\lambda_2 \left( b_2 + \sum_{j=1}^m w_j \right) + (n_2 - 1) \sum_{j=1}^m \ln(1 - e^{-\lambda_2 w_j}) \right).$$

Since these density functions are not well-known distribution, it is not possible to sample directly by standard methods. If the posterior density function is unimodal and roughly symmetric, then it is often convenient to approximate it by a normal distribution (see Gelman et al., [28]). To see the marginal posterior densities are unimodal and roughly symmetric, we check whether the posterior densities have the log-concavity property. It is easily seen that the marginal posterior densities of  $\lambda_1$  and  $\lambda_2$  are log-concave. Therefore, we use the Metropolis-Hasting algorithm with the normal proposal distribution to generate a random sample from the posterior densities of  $\lambda_1$  and  $\lambda_2$  in our implementation. The following algorithm is used.

Step 1: Start with initial guess  $\lambda_1^{(0)}$  and  $\lambda_2^{(0)}$ .

Step 2: Set  $i = 1$ .

Step 3: Generate  $\lambda_1^{(i)}$  from  $\pi_1(\lambda_1 | \lambda_2, \mathbf{v}, \mathbf{w})$  using the Metropolis-Hastings algorithm with the proposal distribution  $q_1(\lambda_1) \equiv N(\lambda_1^{(i-1)}, V_{\lambda_1})$  as follows.

a) Let  $v = \lambda_1^{(i-1)}$ .

b) Generate  $w$  from the proposal distribution  $q_1$ .

c) Let  $p(v, w) = \min \left\{ 1, \frac{\pi_1(w | \lambda_2^{(i)}, \mathbf{v}, \mathbf{w}) q_1(v)}{\pi_1(v | \lambda_1^{(i)}, \mathbf{v}, \mathbf{w}) q_1(w)} \right\}$ .

d) Generate  $u$  from  $Uniform(0, 1)$ . If  $u \leq p(v, w)$ , then accept the proposal and set  $\lambda_1^{(i)} = w$ ; otherwise, set  $\lambda_1^{(i)} = v$ .

Step 4: Similarly,  $\lambda_2^{(i)}$  is generated from  $\pi_2(\lambda_2 | \lambda_1, \mathbf{v}, \mathbf{w})$  using the Metropolis-Hastings algorithm with the proposal distribution  $q_2(\lambda_2) \equiv N(\lambda_2^{(i-1)}, V_{\lambda_2})$ .

Step 5: Compute the  $\Phi_{n_1, n_2}^{(i)}$  at  $(\lambda_1^{(i)}, \lambda_2^{(i)})$ .

Step 6: Set  $i = i + 1$ .

Step 7: Repeat Steps 2 through -7,  $N$  times and obtain the posterior sample  $\Phi_{n_1, n_2}^{(i)}$ ,  $i = 1, \dots, N$ .

This sample is used to compute the Bayes estimate and to construct the HPD credible interval for  $\Phi_{n_1, n_2}$ . The Bayes estimate of  $R_{s, k}$  under a SE loss function is given by

$$\hat{\Phi}_{n_1, n_2}^{MCMC} = \frac{1}{N - M} \sum_{i=M+1}^{N-M} \Phi_{n_1, n_2}^{(i)}, \quad (11)$$

where  $M$  is the burn-in period.

The HPD  $100(1 - \gamma)\%$  credible interval of  $R_{s, k}$  is obtained by the method of Chen and Shao [29].

#### 4. SIMULATION STUDY

In this section, some numerical results are presented to compare the performance of the ML and Bayes estimates of  $\Phi_{n_1, n_2}$  for different parameters and sample sizes. The performances of the point estimators are compared by using mean squared

error (MSE) and estimated risks (ERs). The performances of the asymptotic confidence and credible intervals are compared by using average confidence lengths and coverage probabilities (cps). The coverage probability of a confidence interval is the proportion of the time that the interval contains the true value of interest. The ER of  $\theta$ , when  $\theta$  is estimated by  $\hat{\theta}$ , is given by

$$ER(\theta) = \frac{1}{N} \sum_{i=1}^N \left( \hat{\theta}_i - \theta_i \right)^2,$$

under the SE loss function. All of the computations are performed by using MATLAB. All the results are based on 2500 replications.

In Tables 1-4, strength and stress samples are generated for  $(n_1, n_2) = (5, 5), (10, 10), (15, 10)$  and  $(\lambda_1, \lambda_2) = (0.5, 10), (1, 10), (1.5, 10), (2, 10)$  and different sample sizes  $n$  and  $m = 10(10)50$ . The hyperparameters are chosen that prior means are exactly equal to the true values of the parameters. For this reason  $(a_1, b_1) = (5, 10), (10, 10), (15, 10), (20, 10)$  and  $(a_2, b_2) = (5, 1/2)$  are used for  $(\lambda_1, \lambda_2) = (0.5, 10), (1, 10), (1.5, 10), (2, 10)$ , respectively. For these samples, estimations of  $\Phi_{n_1, n_2}$  are listed based on the MLE and Bayesian estimates which are obtained by using Lindley's approximation and MCMC method. Moreover, 95% asymptotic confidence interval and HPD credible interval of  $\Phi_{n_1, n_2}$  with its coverage probabilities (cps) are presented.

In the MCMC case, we run three MCMC chains with fairly different initial values and generate 5000 iterations for each chain. To diminish the effect of the starting distribution, a certain number of the first 2500 draws is discarded. This is known as the burn-in. In our case, we discard the first 2500 iterations of each sequence and focus on the other 2500 iterations. In order to break the dependence between draws in the Markov chain, it is suggested only to keep every  $d$ th draw of the chain. This is known as thinning. In our case, we calculate the Bayesian MCMC estimates by the means of every 5th sampled values after discarding the first 2500 iterations of the chains. To monitor convergence of MCMC simulations the scale reduction factor estimate is used. The estimate is given by  $\sqrt{Var(\psi)/W}$ , where  $\psi$  is the estimand of interest,  $Var(\psi) = (n-1)W/n + B/n$  with the iteration number  $n$  for each chain, the between-sequence variance  $B$  and the within-sequence variance  $W$ , see Gelman et al. [28]. In our case, the scale factor values of the MCMC estimators are found to be below 1.1, which is an acceptable value for their convergence.

From Tables 1-4, it is observed that the average MSEs of ML estimates and ERs of the Bayes estimates of  $\Phi_{n_1, n_2}$  decrease as the sample size increases in all cases, as expected. The Bayes estimates of  $\Phi_{n_1, n_2}$  have smaller errors than that of MLEs. Moreover, the ERs of the Bayes estimates which are obtained from the MCMC method are smaller than those obtained from Lindley's approximation. The average lengths of the intervals decrease as the sample size increases. The average lengths of the Bayesian credible intervals are smaller than those of the asymptotic

TABLE 1. Estimates and confidence interval of  $\Phi_{n_1, n_2}$ .

$\lambda_1 = 0.5$ and $\lambda_2 = 10$									
$n_1$	$n_2$	$n$	$m$	$\Phi_{n_1, n_2}$	$\hat{\Phi}_{n_1, n_2}^{MLE}$	$\hat{\Phi}_{n_1, n_2}^{Lindley}$	$\hat{\Phi}_{n_1, n_2}^{MCMC}$	$ACI$ of $\Phi_{n_1, n_2}$	$HPDCI$ of $\Phi_{n_1, n_2}$
5	5	10	10	4.3387	4.3587	4.5344	4.4223	(2.8714, 5.8460)	(3.0964, 5.8736)
					0.5969	0.5405	0.4843	2.9746/0.9332	2.7773/0.9524
		20	20		4.3463	4.4411	4.3779	(3.2975, 5.3951)	(3.4074, 5.4112)
					0.2838	0.2704	0.2543	2.0976/0.9428	2.0037/0.9496
		30	30		4.3452	4.4100	4.3660	(3.4891, 5.2013)	(3.5637, 5.2104)
					0.1908	0.1849	0.1772	1.7122/0.9488	1.6467/0.9484
		40	40		4.3493	4.3984	4.3651	(3.6073, 5.0913)	(3.6655, 5.0973)
					0.1434	0.1407	0.1363	1.4840/0.9564	1.4318/0.9504
		50	50		4.3476	4.3871	4.3599	(3.6841, 5.0110)	(3.7324, 5.0131)
					0.1138	0.1121	0.1091	1.3270/0.9460	1.2807/0.9484
10	10	10	10	5.6550	5.6204	5.7751	5.6556	(4.1033, 7.1374)	(4.2848, 7.1228)
					0.6173	0.5974	0.5391	3.0341/0.9444	2.8380/0.9488
		20	20		5.5861	5.6682	5.6031	(4.5199, 6.6522)	(4.6182, 6.6403)
					0.2945	0.2895	0.2743	2.1323/0.9460	2.0221/0.9464
		30	30		5.5887	5.6443	5.5994	(4.7178, 6.4596)	(4.7880, 6.4430)
					0.2111	0.2095	0.2025	1.7418/0.9512	1.6550/0.9424
		40	40		5.5788	5.6210	5.5865	(4.8258, 6.3317)	(4.8867, 6.3141)
					0.1480	0.1470	0.1436	1.5059/0.9480	1.4274/0.9344
		50	50		5.5667	5.6008	5.5738	(4.8947, 6.2387)	(4.9488, 6.2203)
					0.1218	0.1204	0.1185	1.3440/0.9460	1.2715/0.9320
15	10	10	10	6.3436	6.3792	6.5217	6.4019	(4.8683, 7.8901)	(5.0233, 7.8625)
					0.5917	0.5764	0.5284	3.0218/0.9464	2.8392/0.9468
		20	20		6.3653	6.4397	6.3746	(5.2994, 7.4313)	(5.3864, 7.4065)
					0.2979	0.2948	0.2818	2.1319/0.9528	2.0201/0.9408
		30	30		6.3483	6.3988	6.3551	(5.4802, 7.2165)	(5.5465, 7.1931)
					0.1912	0.1890	0.1845	1.7363/0.9536	1.6466/0.9472
		40	40		6.3619	6.3999	6.3670	(5.6086, 7.1153)	(5.6660, 7.0888)
					0.1465	0.1463	0.1429	1.5067/0.9568	1.4227/0.9392
		50	50		6.3434	6.3741	6.3479	(5.6715, 7.0154)	(5.7204, 6.9906)
					0.1186	0.1176	0.1160	1.3439/0.9472	1.2702/0.9312

Notes: 1st row represents the average estimates, 95% confidence interval and 2nd row represents corresponding MSE or ERs, interval lengths and cps for the point and interval estimates, respectively.

TABLE 2. Estimates and confidence interval of  $\Phi_{n_1, n_2}$ .

$\lambda_1 = 1$ and $\lambda_2 = 10$									
$n_1$	$n_2$	$n$	$m$	$\Phi_{n_1, n_2}$	$\hat{\Phi}_{n_1, n_2}^{MLE}$	$\hat{\Phi}_{n_1, n_2}^{Lindley}$	$\hat{\Phi}_{n_1, n_2}^{MCMC}$	$ACI$ of $\Phi_{n_1, n_2}$	$HPDCI$ of $\Phi_{n_1, n_2}$
5	5	10	10	2.0583	2.0612	2.1311	2.0919	(1.3207, 2.8018)	(1.4553, 2.7841)
					0.1466	0.1027	0.0938	1.4811/0.9392	1.3288/0.9676
		20	20		2.0723	2.1120	2.0858	(1.5459, 2.5986)	(1.6075, 2.5936)
					0.0715	0.0604	0.0566	1.0527/0.9492	0.9862/0.9632
		30	30		2.0638	2.0924	2.0731	(1.6357, 2.4919)	(1.6754, 2.4909)
					0.0481	0.0428	0.0409	0.8561/0.9460	0.8156/0.9564
		40	40		2.0628	2.0850	2.0697	(1.6922, 2.4334)	(1.7215, 2.4332)
					0.0359	0.0329	0.0318	0.7412/0.9492	0.7117/0.9548
		50	50		2.0607	2.0788	2.0664	(1.7295, 2.3919)	(1.7533, 2.3919)
					0.0270	0.0252	0.0245	0.6624/0.9552	0.6385/0.9596
10	10	10	10	2.6362	2.6484	2.7128	2.2660	(1.8914, 3.4055)	(1.9922, 3.3828)
					0.1532	0.1238	0.1131	1.5140/0.9428	1.3906/0.9596
		20	20		2.6503	2.6860	2.6575	(2.1145, 3.1860)	(2.1624, 3.1755)
					0.0749	0.0678	0.0641	1.0715/0.9504	1.0131/0.9584
		30	30		2.6433	2.6683	2.6481	(2.2069, 3.0796)	(2.2397, 3.0724)
					0.0520	0.0485	0.0468	0.8728/0.9412	0.8326/0.9468
		40	40		2.6395	2.6588	2.6429	(2.2622, 3.0168)	(2.2880, 3.0095)
					0.0381	0.0361	0.0351	0.7546/0.9504	0.7215/0.9476
		50	50		2.6392	2.6548	2.6420	(2.3017, 2.9766)	(2.3231, 2.9700)
					0.0292	0.0280	0.0274	0.6749/0.9440	0.6468/0.9432
15	10	10	10	3.0253	3.0237	3.0849	3.0377	(2.2690, 3.7783)	(2.3546, 3.7561)
					0.1396	0.1175	0.1095	1.5093/0.9568	1.4015/0.9688
		20	20		3.0367	3.0695	3.0412	(2.5015, 3.5718)	(2.5431, 3.5591)
					0.0766	0.0709	0.0678	1.0704/0.9456	1.0160/0.9464
		30	30		3.0331	3.0558	3.0358	(2.5966, 3.4696)	(2.6249, 3.4597)
					0.0509	0.0483	0.0469	0.8729/0.9488	0.8348/0.9496
		40	40		3.0283	3.0457	3.0305	(2.6509, 3.4057)	(2.6734, 3.3968)
					0.0366	0.0351	0.0345	0.7548/0.9524	0.7235/0.9520
		50	50		3.0280	3.0421	3.0295	(2.6905, 3.3655)	(2.7097, 3.3565)
					0.0293	0.0284	0.0280	0.6749/0.9584	0.6468/0.9520

Notes: 1st row represents the average estimates, 95% confidence interval and 2nd row represents corresponding MSE or ERs, interval lengths and cps for the point and interval estimates, respectively.

TABLE 3. Estimates and confidence interval of  $\Phi_{n_1, n_2}$ .

$\lambda_1 = 1.5$ and $\lambda_2 = 10$									
$n_1$	$n_2$	$n$	$m$	$\Phi_{n_1, n_2}$	$\hat{\Phi}_{n_1, n_2}^{MLE}$	$\hat{\Phi}_{n_1, n_2}^{Lindley}$	$\hat{\Phi}_{n_1, n_2}^{MCMC}$	$ACI$ of $\Phi_{n_1, n_2}$	$HPDCI$ of $\Phi_{n_1, n_2}$
5	5	10	10	1.3032	1.3086	1.3452	1.3306	(0.8180, 1.7991)	(0.9194, 1.7751)
			20		0.0614	0.0348	0.0323	0.9811/0.9444	0.8558/0.9832
			30		1.3063	1.3299	1.3175	(0.9596, 1.6531)	(1.0064, 1.6477)
			40		0.0324	0.0240	0.0228	0.6934/0.9444	0.6413/0.9704
		30	30		1.3058	1.3230	1.3131	(1.0227, 1.5890)	(1.0521, 1.5873)
			40		0.0214	0.0174	0.0168	0.5663/0.9452	0.5352/0.9628
			50		1.3057	1.3192	1.3112	(1.0605, 1.5509)	(1.0822, 1.5503)
			40		0.0151	0.0129	0.0126	0.4905/0.9476	0.4681/0.9608
		50	50		1.3029	1.3142	1.3073	(1.0839, 1.5218)	(1.1010, 1.5214)
			40		0.0125	0.0110	0.0107	0.4379/0.9492	0.4204/0.9576
			30		1.6686	1.7037	1.6837	(1.1628, 2.1744)	(1.2394, 2.1553)
			20		0.0680	0.0472	0.0440	1.0117/0.9452	0.9159/0.9676
10	10	10	10	1.6609	1.6700	1.6908	1.6760	(1.3118, 2.0282)	(1.3469, 2.0202)
			20		0.0336	0.0281	0.0267	0.7164/0.9516	0.6733/0.9640
			30		1.6661	1.6811	1.6701	(1.3744, 1.9579)	(1.3975, 1.9529)
			40		0.0218	0.0193	0.0187	0.5836/0.9496	0.5554/0.9528
		30	30		1.6690	1.6805	1.6719	(1.4160, 1.9220)	(1.4336, 1.9178)
			40		0.0163	0.0150	0.0146	0.5060/0.9544	0.4842/0.9532
			50		1.6622	1.6719	1.6648	(1.4366, 1.8878)	(1.4509, 1.8847)
			40		0.0131	0.0121	0.0119	0.4512/0.9512	0.4338/0.9528
		50	50		1.9289	1.9617	1.9401	(1.4209, 2.4370)	(1.4829, 2.4192)
			20		0.0690	0.0520	0.0486	1.0161/0.9400	0.9363/0.9692
			30		1.9290	1.9479	1.9330	(1.5698, 2.8882)	(1.5982, 2.2801)
			40		0.0339	0.0295	0.0283	0.7184/0.9532	0.6819/0.9588
15	15	10	10	1.9194	1.9234	1.9370	1.9260	(1.6310, 2.2157)	(1.6503, 2.2098)
			20		0.0222	0.0202	0.0196	0.5847/0.9472	0.5595/0.9520
			30		1.9178	1.9285	1.9197	(1.6652, 2.1704)	(1.6800, 2.1656)
			40		0.0163	0.0150	0.0148	0.5052/0.9532	0.4856/0.9524
		30	30		1.9242	1.9327	1.9255	(1.6977, 2.1508)	(1.7100, 2.1464)
			40		0.0139	0.0132	0.0129	0.4531/0.9444	0.4364/0.9428
		50	50		1.9242	1.9327	1.9255	(1.6977, 2.1508)	(1.7100, 2.1464)
			40		0.0139	0.0132	0.0129	0.4531/0.9444	0.4364/0.9428
			30		1.9242	1.9327	1.9255	(1.6977, 2.1508)	(1.7100, 2.1464)
			20		0.0139	0.0132	0.0129	0.4531/0.9444	0.4364/0.9428
		50	50		1.9242	1.9327	1.9255	(1.6977, 2.1508)	(1.7100, 2.1464)
			40		0.0139	0.0132	0.0129	0.4531/0.9444	0.4364/0.9428

Notes: 1st row represents the average estimates, 95% confidence interval and 2nd row represents corresponding MSE or ERs, interval lengths and cps for the point and interval estimates, respectively.

TABLE 4. Estimates and confidence interval of  $\Phi_{n_1, n_2}$ .

$\lambda_1 = 2$ and $\lambda_2 = 10$									
$n_1$	$n_2$	$n$	$m$	$\Phi_{n_1, n_2}$	$\hat{\Phi}_{n_1, n_2}^{MLE}$	$\hat{\Phi}_{n_1, n_2}^{Lindley}$	$\hat{\Phi}_{n_1, n_2}^{MCMC}$	$ACI$ of $\Phi_{n_1, n_2}$	$HPDCI$ of $\Phi_{n_1, n_2}$
5	5	10	10	0.9311	0.9382	0.9592	0.9553	(0.5748, 1.3015)	(0.6567, 1.2788)
			20		0.0356	0.0167	0.0160	0.7267/0.9396	0.6221/0.9888
			30		0.9357	0.9507	0.9448	(0.6789, 1.1925)	(0.7167, 1.1871)
			40		0.0165	0.0110	0.0106	0.5136/0.9484	0.4704/0.9776
		30	30		0.9345	0.9460	0.9406	(0.7250, 1.1441)	(0.7489, 1.1425)
			40		0.0112	0.0085	0.0082	0.4191/0.9456	0.3936/0.9696
			50		0.9325	0.9419	0.9373	(0.7513, 1.1137)	(0.7688, 1.1138)
			40		0.0083	0.0067	0.0065	0.3625/0.9580	0.3449/0.9732
		50	50		0.9303	0.9383	0.9343	(0.7686, 1.0920)	(0.7821, 1.0925)
			40		0.0069	0.0058	0.0057	0.3234/0.9424	0.3103/0.9624
			30		1.1804	1.2022	1.1943	(0.8023, 1.5585)	(0.8643, 1.5436)
			20		0.0370	0.0224	0.0212	0.7561/0.9452	0.6793/0.9800
10	10	10	10	1.1756	1.1808	1.1947	1.1867	(0.9131, 1.4484)	(0.9413, 1.4433)
			20		0.0186	0.0144	0.0138	0.5353/0.9428	0.5020/0.9644
			30		1.1759	1.1864	1.1801	(0.9579, 1.3938)	(0.9760, 1.3913)
			40		0.0122	0.0102	0.0099	0.4359/0.9516	0.4154/0.9660
		30	30		1.1780	1.1861	1.1809	(0.9891, 1.3670)	(1.0027, 1.3649)
			40		0.0097	0.0085	0.0083	0.3780/0.9440	0.3622/0.9544
			50		1.1780	1.1846	1.1801	(1.0888, 1.3471)	(1.0196, 1.3451)
			40		0.0075	0.0068	0.0066	0.3383/0.9480	0.3254/0.9508
		50	50		1.3729	1.3933	1.3840	(0.9907, 1.7552)	(1.0413, 1.7431)
			20		0.0385	0.0258	0.0245	0.7645/0.9404	0.7018/0.9708
			30		1.3710	1.3836	1.3755	(1.1007, 1.6413)	(1.1228, 1.6366)
			40		0.0193	0.0157	0.0152	0.5407/0.9484	0.5137/0.9648
15	15	10	10	1.3671	1.3740	1.3828	1.3762	(1.1530, 1.5950)	(1.1668, 1.5914)
			20		0.0121	0.0106	0.0103	0.4420/0.9576	0.4245/0.9644
			30		1.3669	1.3742	1.3689	(1.1764, 1.5575)	(1.1875, 1.5549)
			40		0.0096	0.0086	0.0085	0.3811/0.9488	0.3674/0.9520
		30	30		1.3707	1.3765	1.3721	(1.1998, 1.5415)	(1.2086, 1.5390)
			40		0.0078	0.0072	0.0070	0.3417/0.9480	0.3304/0.9492
			50		1.3707	1.3765	1.3721	(1.1998, 1.5415)	(1.2086, 1.5390)
			40		0.0078	0.0072	0.0070	0.3417/0.9480	0.3304/0.9492
		50	50		1.3707	1.3765	1.3721	(1.1998, 1.5415)	(1.2086, 1.5390)
			40		0.0078	0.0072	0.0070	0.3417/0.9480	0.3304/0.9492
			30		1.3707	1.3765	1.3721	(1.1998, 1.5415)	(1.2086, 1.5390)
			20		0.0078	0.0072	0.0070	0.3417/0.9480	0.3304/0.9492

Notes: 1st row represents the average estimates, 95% confidence interval and 2nd row represents corresponding MSE or ERs, interval lengths and cps for the point and interval estimates, respectively.



confidence intervals. Their coverage probabilities are close to the nominal level 95%.

## 5. CONCLUSIONS

In this paper, we have studied the mean remaining strength of the parallel systems in the stress-strength model. We obtain the conditional random variable for the remaining strength of the parallel system under the applied parallel stress system. The likelihood ratio ordering between two systems is established for two-component case. Currently, we do not prove it is true in number of components are greater than two. The proof of this general case can be considered as a future work. Moreover, the maximum likelihood and Bayes estimates of the mean remaining strength of the system is derived and compared.

**Acknowledgement.** The author would like to thank the Editor and two anonymous referees for their valuable comments which led to this improved version.

## REFERENCES

- [1] Birnbaum, Z. W., On a use of Mann-Whitney statistics, in *Proc. 3rd Berkeley Symposium on Mathematical Statistics and Probability* 1, (1956), 13-17.
- [2] Birnbaum, Z. W. and McCarty, B. C., A distribution-free upper confidence bounds for  $Pr(Y < X)$  based on independent samples of  $X$  and  $Y$ , *The Annals of Mathematical Statistic* 29(2), (1958), 558-562.
- [3] Kotz, S., Lumelskii, Y., Pensky, M., *The Stress-Strength Model and its Generalizations: Theory and Applications*, World Scientific, Singapore, 2003.
- [4] Kundu, D., Gupta, R. D., Estimation of  $P(Y < X)$  for generalized exponential distribution, *Metrika* 61, (2005), 291-308.
- [5] Basirat, M., Baratpour, S. and Ahmadi, J., Statistical inferences for stress-strength in the proportional hazard models based on progressive Type-II censored samples, *Journal of Statistical Computation and Simulation* 85(3), (2015), 431-449.
- [6] Basirat, M., Baratpour, S. and Ahmadi, J., On estimation of stress-strength parameter using record values from proportional hazard rate models, *Communications in Statistics - Theory and Methods* 45(19), (2016), 5787-5801.
- [7] Asgharzadeh, A., Kazemi, M. and Kundu, D., Estimation of  $P(X > Y)$  for Weibull distribution based on hybrid censored samples, *International Journal of System Assurance Engineering and Management* 8(1), (2017), 489-498.
- [8] Bhattacharyya, G. K. and Johnson, R. A. Estimation of reliability in multicomponent stress-strength model, *Journal of American Statistical Association* 69, (1974), 966-970.
- [9] Bhattacharyya, G. K. and Johnson, R. A. Stress-strength models for system reliability, *Proceedings of the Symposium on Reliability and Fault Tree Analysis SIAM*, (1975), 509-532.
- [10] Eryilmaz, S., Consecutive k-Out-of-n : G System in Stress-Strength Setup, *Communications in Statistics - Simulation and Computation*, 37(3), (2008), 579-589.
- [11] Eryilmaz, S., On system reliability in stress-strength setup, *Statistics & Probability Letters* 80, (2010), 834-839.
- [12] Pakdaman, Z. and Ahmadi, J., Stress-strength reliability for  $P(X_{r:n_1} < Y_{k:n_2})$  in the exponential case, *Istatistik: Journal of The Turkish Statistical Association* 6(3), (2013), 92-102.

- [13] Pakdaman, Z. and Ahmadi, J., Point estimation of the stress-strength reliability parameter for parallel system with independent and non-identical components, *Communications in Statistics-Simulation and Computation*, 47(4), (2018), 1193-1203.
- [14] Hassan, M. K. H., Estimation a stress-strength model for  $P(Y_{r:n_1} < X_{k:n_2})$  using the Lindley distribution, *Revista Colombiana de Estadística* 40(1), (2017), 105-121.
- [15] Kızılaslan, F., Classical and Bayesian estimation of reliability in a multicomponent stress-strength model based on the proportional reversed hazard rate mode, *Mathematics and Computers in Simulation* 136, (2017), 36-62.
- [16] Gürler, S., The mean remaining strength of systems in a stress-strength model, *Hacettepe Journal of Mathematics and Statistics* 42(2), (2013), 181-187.
- [17] Bairamov (Bayramoglu), I., Gurler, S. and Ucer, B., On the mean remaining strength of the  $k$ -out-of- $n$  :  $F$  system with exchangeable components, *Communications in Statistics-Simulation and Computation* 44(1), (2015), 1-13.
- [18] Gurler, S., Ucer, B. H. and Bairamov, I., On the mean remaining strength at the system level for some bivariate survival models based on exponential distribution, *Journal of Computational and Applied Mathematics* 290, (2015), 535-542.
- [19] Gupta, R. D. and Kundu, D., Generalized exponential distributions, *Australian & New Zealand Journal of Statistics* 41, (1999), 173-188.
- [20] Dewan, I. and Khaledi, B., On stochastic comparisons of residual life time at random time, *Statistics and Probability Letters* 88, (2014), 73-79.
- [21] Misra, N. and Naqvi, S., Stochastic comparison of residual lifetime mixture models, *Operations Research Letters* 46, (2018), 122-127.
- [22] Misra, N. and Naqvi, S., Some unified results on stochastic properties of residual lifetime at random times, *Brazilian Journal of Probability and Statistics* 32(2), (2018), 422-436.
- [23] Shakedand, M. and Shanthikumar, J. G., Stochastic Orders, Springer Series in Statistics, 2007.
- [24] Ghitany, M. E., Al-Jarallah, R. A. and Balakrishnan, N., On the existence and uniqueness of the MLEs of the parameters of a general class of exponentiated distributions, *Statistics* 47(3), (2013), 605-612.
- [25] Gradshteyn, I. S and Ryzhik, I. M., Table of Integrals, Series and Products, 5th ed. Boston, USA: Academic Press, 2007.
- [26] Rao, C. R., Linear Statistical Inference and Its Applications. New York: Wiley; 1965.
- [27] Lindley, D. V., Approximate Bayes method, *Trabajos de Estadística* 3, (1980), 281-288.
- [28] Gelman, A., Carlin, J. B., Stern, H. S. and Rubin, D. B., Bayesian Data Analysis. Chapman Hall, London, 2003.
- [29] Chen, M. H. and Shao, Q. M., Monte Carlo estimation of Bayesian credible and HPD intervals, *Journal of Computational and Graphical Statistics* 8(1), (1999), 69-92.

*Current address:* Fatih Kızılaslan: Department of Statistics, Marmara University, Istanbul, Turkey.

*E-mail address:* fatih.kizilaslan@marmara.edu.tr

*ORCID Address:* <http://orcid.org/0000-0001-6457-0967>