# PAPER DETAILS

TITLE: Report on the determination of the coordinates of the Astronomical Observatory of the

Ankara University

AUTHORS: Arif ÇÖKLÜ

PAGES: 0-0

ORIGINAL PDF URL: https://dergipark.org.tr/tr/download/article-file/1626882

## Report on the determination of the coordinates of the Astronomical Observatory of the . Ankara University

### by Arif ÇÖKLÜ and Faruk UZEL

#### (Submitted by the Department of Astronomy)

Özet: Ankara Fen Fakültesi Astronomi rasathanesinin coğrafya koordinatları üzerine kısa bir rapor verilmektedir.

Sonuçlar :

Cografya boylam	$\lambda = z^h  11^m  07^s,  131 \pm 0^s.006$
Coğrafya enlem	$\varphi = 39^{\circ} 50' 37'', 002 \pm 0'', 14$
Denizden olan yükseklik	h = 1256,69 metre

#### \*\*

Summary: An abbreviated report is submitted concerning the determination of the geographic coordinates of the Astronomical Observatory of the Fen Fakültesi. The results are:

Geographic	longitude	
Geographic latitude		
Height abov	ve sea level	

 $\lambda = 2^{h} 11^{m} 07^{s}, 181 \pm 0^{s}.006$   $\varphi = 89^{\circ} 50' 87'', 002 \pm 0'', 14$ h = 1256.69 meter

The new Observatory of Ankara is situated on a flat ridge at a distance, along the road, of 18 km from the city. The observations needed to find the exact longitude and latitude started September 27 - 1960.

\*\*

#### 1. Determination of latitude.

Observer : Arif Çöklü.

Instrument: Wild T<sub>4</sub>

Method : Horrebor - Talcott

#### THE COORDINATES OF THE OBSERVATORY OF ANKARA 21

As is known with this method in the meridian pairs of stars are measured while with each pair the positions of the two stars relative to the zenith is symmetric. In this way during two nights  $12 \times 2$  stars were measured. From these observations we find:

$$\varphi = \frac{\delta_n + \delta_s}{2}$$

Where  $\delta_n$  and  $\delta_n$  are the declinations of the stars south and north of the zenith respectively. In this way from the twelve pairs for the geographic latitude we find

$$39^{\circ} 50' 37''.213 \pm 0''.14$$

For the reduction to sea level (Hayford - Ellipsoide) we apply the coorection  $D = 0,000171 h_m \sin 2\varphi = -0^{\circ},211$ . Consequently we have:

latitude observed	39° 50′ 37″, 213 ± 0″,14
correction	0",211
corrected latitude	39° 50′ 37″, 002 ± 0″,14

#### 2. Determination of longitude:

Observer: Faruk Uzel. Instrument: Wild T<sub>4</sub> choronometer Ulysse Nardin

choronograph Wetzer receiver RG 37 Marconi

#### Method: Registration.

During four nights 3 groups of six stars each were observed. Each group consisted out of 5 stars with declination  $18^{\circ} < \delta < 55^{\circ}$ and one circumpolar star for which  $\delta > 70^{\circ}$ . Altogether during these four nights 72 stars were observed. After each night the mean value  $\Delta u$  for the group of 18 stars was calculated and next the longitude was determined with the help of the signals TQG5, RWM and GBR.

## $\lambda = 2^{h} \, 11^{m} \, 07^{s}, \, 131 \pm 0^{s}, 006$

With both set of measurements the following catalogues were used: «FK 3 Apparent places of fundamental stars» and Boss «General Catalogue».

## 3. Heights :

Through comparison with other points the altitude was found to be h = 1256,69 meter.

(Manuscript received 21 th June 1961)