

PAPER DETAILS

TITLE: On 6- Figures In Moufang Projective Planes

AUTHORS: Süleyman ÇİFTÇİ

PAGES: 0-0

ORIGINAL PDF URL: <https://dergipark.org.tr/tr/download/article-file/1640074>

ON 6- FIGURES IN MOUFANG PROJECTIVE PLANES

SÜLEYMAN ÇİFTÇİ*

* Department of Mathematics, Uludağ University, Bursa /TURKEY.

ABSTRACT

In this paper, the concept of cross-ratio is extended to the whole Moufang plane and some properties of 6-figures in the π Moufang plane is examined. Essentially the geometric properties of the π which is equivalent to the existence of a square root of an element of an alternative division ring R is determined.

INTRODUCTION

Let Π be a Moufang projective plane. It is well known that π determines a unique alternative ring R . One of the main problems in projective geometry is to find geometric properties of π which are equivalent to certain algebraic properties of the alternative ring R . For instance, Π is Desarguesian if and only if R is associative. In this paper, we extend some of the properties of the 6-figures, which have been given by Cater [1] for Desarguesian planes, to the Moufang planes. Namely, we determined the geometric properties of Π equivalent to the existence of a square root of an element of R . And if $(A,B:C,D)$ denotes the cross-ratio of distinct collinear points A,B,C,D in Π , we construct points J and N such that $(A,B:C,J) = (A,B:C,D)^2$ and $(A,B:C,N) = (A,B:C,D)^3$. In fact, these are investigated in [2]. But some mappings which is obtained by using the identity

$$(1) \quad x^{-1}(y(xz)) = (x^{-1}yx)z$$

asserted in [5] for Cayley-Dickson algebras are used. However [3] demonstrated here that (1) does not valid for Cayley-Dickson algebras. But also the existence of 4-point transitivity is shown by using of some new mappings. Essentially, in this paper, [2] is rearranged and modified under the light of [3].

Throughout the paper we use the terminology in [1] and [3]: A 6-figure is a sequence of 6 distinct points $(ABC, A'B'C')$ such that ABC is a triangle, and $A' \in BC$, $B' \in CA$, $C' \in AB$. The points A, B, C, A', B', C' are called vertices of this 6-figure. A 6-figure $(ABC, A'B'C')$ is said to be equivalent to any 6-figure $(DEF, D'E'F')$ if there exist a projective collineation of Π which maps A, B, C, A', B', C' to D, E, F, D', E', F' respectively; in symbols $ABCA'B'C' \sim DEFD'E'F'$.

$(ABC, A'B'C')$ is called a menelaus 6-figure if A', B', C' are collinear; and $(ABC, A'B'C')$ is called a ceva 6-figure if the lines AA', BB', CC' are concurrent.

Throughout the paper, we assume that R is an alternative division ring with center K of arbitrary characteristic and that Π is the Moufang plane coordinatized by R , [4].

It is easy to see that the mappings

$$\begin{aligned} I_1: (x, y) &\rightarrow (x^{-1}, yx^{-1}), x \neq 0, (0, y) \leftrightarrow (y), (\infty) \rightarrow (\infty) \\ [m, k] &\rightarrow [k, m], [k] \rightarrow [k^{-1}], k \neq 0, [0] \leftrightarrow [\infty] \end{aligned}$$

and

$$\begin{aligned} I_2: (x, y) &\rightarrow (xy^{-1}, y^{-1}), y \neq 0, (x, 0) \leftrightarrow (x^{-1}), x \neq 0, (0, 0) \leftrightarrow (\infty), (0) \rightarrow (0) \\ [m, k] &\rightarrow [-k^{-1}m, k^{-1}], k \neq 0, [m, 0] \leftrightarrow [m^{-1}], m \neq 0, [0, 0] \leftrightarrow [\infty], [0] \rightarrow [0] \end{aligned}$$

are collineations of Π .

Let $A=(0)$, $B=(\infty)$, $C=(0,0)$, $A'=(0,1)$, $B'=(-1,0)$, $C'=(-m)$ for some $m \in R$. Here, $(-1,0) (\infty) (0,1) (0) = (-1,1)$ and

$$I_2 I_1: (0), (\infty), (0,0), (-1,1) \rightarrow (\infty), (0,0), (0), (1,-1).$$

According to the Theorem 1 in [3], the mapping g which maps $(0), (\infty), (0,0), (1,-1)$, to $(0), (\infty), (0,0), (-1,m)$ is a composition of the collineations F_c and $S_{\alpha, \beta}$ where $c \in R$, $\alpha, \beta \in K$. We already known that such a collineation, maps (x,y) to (x', yd) . Now, since g maps $(1,-1)$ to $(-1,m)$ than $(-1) d = m$ and so $-m^{-1} = d^{-1}$.

Therefore $(0, -m^{-1})$ maps to $(0, -m^{-1} d) = (0,1)$. Thus

$$f = g I_2 I_1: (0), (\infty), (0,0), (0,1), (-1,0), (-m) \rightarrow (\infty), (0,0), (0), (-1,0), (-m), (0,1).$$

Consequently

$$ABCA'B'C' \sim BCAB'C'A' \sim CABC'A'B'$$

is shown by using f and f^2 .

Furthermore, any 6-figure $(DEF, D'E'F')$ is equivalent to $((0)(\infty)(0,0), (0,1)(-1,0)(-m))$ for some $m \in R$; since there exist by Theorem 1 in [3], a collineation mapping D, E, D', E' 4-point on $(0), (\infty), (0,1), (-1,0)$. Thus $(ABC, A'B'C'), (BCA, B'C'A'), (CAB, C'A'B')$ will be regarded as the same 6-figure μ , and likewise $(ACB, A'C'B'), (CBA, C'B'A'), (BAC, B'A'C')$ as the same 6-figure λ . μ and λ are called opposite 6-figures of each other; in symbols, $\lambda = \mu^{-1}$ and $\mu = \lambda^{-1}$.

Let Π be a Moufang plane satisfying the Fano's Axiom. It follows, (see [7]) that there exist unique points $A'' \in BC, B'' \in CA, C'' \in AB$ such that $H(AB, C'C''), H(BC, A'A''), H(CA, B'B'')$. The 6-figure $(ABC, A''B''C'')$ is called the conjugate of μ , in symbol $-\mu$. Likewise μ is the conjugate of $-\mu$.

Let $C^d \in AB$ be the point such that C, C^d and AA', BB' are collinear. Let $A^d \in BC$ and $B^d \in CA$ be the points such that A, A^d and BB', CC' are collinear and B, B^d and AA', CC' are collinear. The 6-figure $(ACB, A^d C^d B^d)$ is called the first descendant of μ , written μ^d . μ is called a first ancestor of μ^d .

Let $A^c = BC \cdot B'C', B^c = CA \cdot C'A', C^c = AB \cdot A'B'$. The figure $(ACB, A^c C^c B^c)$ is called the first codescendant of μ , written μ^c . μ is called a first coancestor of μ^c .

Figure 1 represents the 6-figure $\mu_1 = ((0)(\infty)(0,0), (0,1)(-1,0)(-m))$ and μ_1^c and construction of B^{cc} . In figure 2 μ_1, μ_1^d are drawn and B^{dc} is constructed. In figure 3 the points $A''^d, B''^d, C''^d, B''^c$ are drawn.




Figure 1




Figure 3




Figure 2

SOME PROPERTIES OF 6-FIGURES IN MOUFANG PLANES

The classical definition of the cross-ratio for Desarguesian planes is not available for the Moufang planes. Ferrar [5] gives the following algebraic definition of the cross-ratio for the points on the line $[0,0]$.

$$(A,B;C,D) = (a,b;c,d) = \{((a-d)^{-1}(b-d))((b-c)^{-1}(a-c))\}$$

where $A = (a,0)$, $B = (b,0)$, $C = (c,0)$, $D = (d,0)$ and $\{x\}$ denotes the conjugacy class of x in the alternative ring R . We extend this definition to whole plane as follows:

(i) If $A = (a, a_1)$, $B = (b, b_1)$, $C = (c, c_1)$, $D = (d, d_1)$ are on a line of type $[m, k]$ let $(A, B; C, D) = (a, b; c, d)$. For this case one of A, B, C, D is the ideal point (m) , use $\infty \notin R$ instead of the corresponding component in $(a, b; c, d)$. (Notice the perspectivity from $[m, k]$ to $[0, 0]$ with center (∞)).

(ii) If A, B, C, D are on a line of type $[k]$ that is, $A = (k, a)$, $B = (k, b)$, $C = (k, c)$, $D = (k, d)$ let $(A, B; C, D) = (a, b; c, d)$. In the case one of A, B, C, D is (∞) use again ∞ in $(a, b; c, d)$. (Notice that the perspectivity from $[k]$ to $[0, 0]$ with center (-1) maps $(k, x) \rightarrow (k+x, 0)$; and taking $k+x$ instead of x does not change the cross ratio.).

(iii) If the points are on $[\infty]$, that is $A = (a)$, $B = (b)$, $C = (c)$, $D = (d)$, let $(A, B; C, D) = (a, b; c, d)$. (Notice that the perspectivity from $[\infty]$ to $[0, 0]$ with center $(0, -1)$ maps $(m) \rightarrow (m^{-1}, 0)$, $(0) \rightarrow (0)$, $(\infty) \rightarrow (0, 0)$, and $(a, b; c, d) = (a^{-1}, b^{-1}; c^{-1}, d^{-1})$).

LEMMA 1. In a Moufang plane a perspectivity preserve the cross ratio.

Proof: Let $A, B, C, D \in l$, φ_M and φ_N be perspectivities from any line l to $[0, 0]$ which have center M and N , respectively. Where φ_M is the perspectivity in definition of generalized cross - ratio, and $N \neq M$, $N \in l$, $N \notin [0, 0]$. In addition let $\varphi_M: A, B, C, D \rightarrow A_1, B_1, C_1, D_1$ and $\varphi_N: A, B, C, D \rightarrow A_0, B_0, C_0, D_0$.

To prove, it is sufficient to show that φ_N preserves the cross-ratio. According to the definition $(A, B; C, D) = (A_1 B_1; C_1 D_1)$. Thus

$$\varphi = \varphi_M \varphi_N^{-1}: A_0, B_0, C_0, D_0 \rightarrow A_1, B_1, C_1, D_1$$

Therefore φ is a projectivity of $[0, 0]$.

Since φ is a projectivity of $[0, 0]$ which preserves the cross-ratio (See [5], Theorem 3,7) $(A_1, B_1; C_1, D_1)$ is equal to $(A_0, B_0; C_0, D_0)$. Consequently $(A, B; C, D) = (A_0, B_0; C_0, D_0)$, that is, φ_N preserves the cross-ratio.

In what follows we use the fact that distinct collinear point A, B, C, D in Π are in harmonic position if and only if $(A, B; C, D) = -1$.

LEMMA 2. If $\mu_1 = (ABC, A'B'C') = ((0) (\infty) (0, 0), (0, 1) (-1, 0) (-m))$ then

$$(A, B; C', C^e) = (B, C; A', A^e) = (C, A; B', B^e) = \{-m\}.$$

Proof is trivial.

Conjugacy class of $-(A,B:C',C^c) = -(C,A:B',B^c)$ is called the ratio of the 6-figure $\mu = (ABC,A'B'C')$, denoted by $r(\mu)$.

THEOREM 1:

(i) μ is a menelaus 6-figure if and only if $r(\mu) = -1$

(ii) μ is a ceva 6-figure if and only if $r(\mu) = 1$

Proof: (i) It suffices to assume that μ is μ_1 because cross-ratio is preserved by projective collineations. Thus $\mu_1 = ((0) (\infty) (0,0), (0,1) (-1,0) (1))$ and $r(\mu_1) = -1$.

Conversely, if $r(\mu_1) = -1$; than $\mu_1 = ((0) (\infty) (0,0), (0,1) (-1,0) (1))$ and the points of $(0,1), (-1,0), (1)$ are collinear.

(ii) If μ_1 is a ceva 6-figure, than $\mu_1 = ((0) (\infty) (0,0), (0,1) (-1,0) (-1))$ and $r(\mu_1) = 1$.

Conversely, if $r(\mu_1) = 1$, μ_1 as above thus, the lines of $(0) (0,1), (\infty) (-1,0), (0,0) (-1)$ is concurrent.

THEOREM 2: For any 6-figure μ we have

$$(i) \quad r(\mu^{-1}) = (r(\mu))^{-1}$$

$$(ii) \quad r(-\mu) = -r(\mu)$$

$$(iii) \quad r(\mu^d) = (r(\mu))^2$$

$$(iv) \quad r(\mu^c) = -(r(\mu))^2$$

Proof: It is sufficient to assume that μ is μ_1 in figures 1,2,3 since cross ratio is preserved by projective collineations. With a simple calculation we have

$$(i) \quad r(\mu_1^{-1}) = -(\infty, 0; -1, m^{-1}) = \{m^{-1}\} = (r(\mu_1))^{-1};$$

$$(ii) \quad r(-\mu_1) = -(0, \infty; 1, m^{-1}) = \{-m\} = -r(\mu_1);$$

$$(iii) \quad r(\mu_1^d) = -(\infty, 0; -m^{-1}, m) = \{m^2\} = (r(\mu_1))^2$$

$$(iv) \quad r(\mu_1^c) = -(\infty, 0; m^{-1}, m) = \{-m^2\} = -(r(\mu_1))^2;$$

It follows immediately, from theorem 1 and 2 that conjugate of a menelaus 6-figure is ceva 6-figure and vice versa. In fact this has been shown in [6].

THEOREM 3. Let $m \in R$, $m \neq 0$. Then the equation $x^2 = m$ (or $x^2 = -m$) has a solution in R if and only if any 6-figure μ with ratio $\{m\}$ has a first ancestor (coancestor) in Π .

Proof: Without loss of generality we take $\mu = \mu_1$. Let $\lambda = ((0) (0,0) (\infty), DEF)$ be a first ancestor of μ_1 . Then we have $\{q^2\} = \{m\}$ by Theorem 1, and there exists an $x \in R$ satisfying $x^2 = m$.

Conversely suppose that there exist $q \in R$ such that $q^2 = m$. In this case it is easily computed that $\lambda_1 = ((0) (0,0) (\infty), (0, -q)(q)(q^{-1}, 0))$ is a first ancestor of μ_1 .

Now consider the equation $x^2 = -m$. Let $\lambda_1 = ((0)(0,0)(\infty), STU)$ be a first coancestor of μ_1 and let $r(\lambda_1) = \{q\}$. Then we have $\{m\} = -\{q^2\}$ by Theorem 1, and there exists an $x \in R$ satisfying $x^2 = -m$.

Suppose $q \in R$ and $q^2 = -m$. Then one can easily show that $\lambda_1 = ((0)(0,0)(\infty), (0,q)(q)(-q^{-1}, 0))$ is a first coancestor of μ_1 .

Clearly one can easily observe from the figures that first descendant of μ and $-\mu$ are same, and that first codescendant of μ and $-\mu$ are same. Namely $\mu^d = (-\mu)^d$ and $\mu^c = (-\mu)^c$.

Finally, it is worth to note that the construction which gives an algorithm for "squaring" a cross-ratio of points in a Desarguesian plane ([1]) can be also extended to a Moufang plane as follows:

Let A, B, C, D be any collinear points in Π . Choose any point E not on AB , and choose points $F \in AE$, $G \in BE$ such that F, G and C are collinear points.

Let $I = AE \cdot DG$, $H = BE \cdot DF$ and $J = AB \cdot HI$. For this project A, B, F, G to $(0), (0,0), (-1), (0,1)$ respectively. Thus $(A, B; C, D) = \{m\}$ and $(A, B; C, J) = \{m^2\}$ by Lemma 1, and consequently $(A, B; C, J) = (A, B; C, D)^2$.

Furthermore, if $K = AE \cdot CH$, $M = AB \cdot KG$ and $G' = BE \cdot MF$, $I' = AE \cdot G'D$, $N = AB \cdot HI'$ then $(A, B; C, N) = (A, B; C, D)^3$.

REFERENCES

1. CATER, F.S.: On Desarguesian projective planes, *Geom. Ded.* 7(1978) 433–441.
2. ÇİFTÇİ, S.: Moufang düzlemleri üzerine, Doktora Tezi, F.Ü. Fen Bilimleri Enstitüsü 1984, Elazığ.
3. ÇİFTÇİ, S., KAYA, R. and FERRAR, J.C.: On 4-Transitivity in the Moufang Plane, *Journal of Geometry* Vol. 31(1988) 65–68.
4. FAULKNER, J.R. and FERRAR, J.C.: Homomorphisms of Moufang planes and alternative places, *Geom. Ded.* 14(1983) 215–223.
5. FERRAR, J.C.: Cross-ratios in Projective and Affine Planes, in *Geometry: Von Staudt's Point of View*, Dordrecht (1980) 101–125.
6. KAYA, R. AND ÇİFTÇİ, S.: On Menelaus and Ceva 6-Figures in Moufang Projective Planes, *Geom. Ded.* 19(1985) 295–296.
7. STEVENSON, F.W.: *Projective Planes*, by W.H. Freeman and Company, San Francisco, U.S.A., 271–402, 1972.