PAPER DETAILS

TITLE: THREAT CATEGORIES OF ENDEMIC PLANTS OF SAKARAT MOUNTAIN (AMASYA/TURKEY) AUTHORS: M Umit BINGOL,Neslihan BALPINAR,Kerim GUNEY,Arzu CANSARAN,Fatmagul GEVEN,Omer Faruk KAYA,Munevver ARSLAN PAGES: 10-21

ORIGINAL PDF URL: https://dergipark.org.tr/tr/download/article-file/710744

Available online: April 28, 2019

Commun.Fac.Sci.Univ.Ank.Series C Volume 28, Number 1, Pages 10-21 (2019) ISSN 1303-6025 E-ISSN 2651-3749 https://dergipark.org.tr/tr/pub/communc/issue/45050/562256



THREAT CATEGORIES OF ENDEMIC PLANTS OF SAKARAT MOUNTAIN (AMASYA/TURKEY)

M. UMIT BINGOL, NESLIHAN BALPINAR, KERIM GUNEY, ARZU CANSARAN FATMAGUL GEVEN, OMER FARUK KAYA, MUNEVVER ARSLAN

ABSTRACT. This study aims to determine the threat categories of endemic plants in the vegetation of Sakarat Mountain (Amasya) on the basis of the vegetation field survey in 2004 and 2005 years. Plant samples were collected in their vegetation seasons. According to results, 78 families, 283 genera and 494 taxa are identified. As a result of field surveys, six different vegetation types forest (decidious, mixed coniferous and deciduous), subalpine, steppe, rocky, wet grassland and segetal were determined. 40 plant taxa (8.09% of all taxa) were confirmed as endemic. The families with the highest rate of endemic taxa are Fabaceae (15%) and Lamiaceae (15%). Phytogeographic regions (Chorotypes) among endemic taxa were listed as Irano-Turanian 13 (32.5%), Euro-Siberian 9 (22.5%), Mediterranean 1 (2.5%), while phytogeographic origin of (17 taxa) 42.5% of endemic taxa were Unknown. The threat categories regarding the endemic species were determined and analysed according to "Red Data Book of Turkish Plants" adopted 2001 IUCN criteria. In the evaluation phase of the endemic taxa and their threat categories, it was found that 1 species (2.5%) is in Endangered, 3 species (7.5%) in Near Threatened and 36 species (90%) in Least Concern according to the criteria of 2001 IUCN. When the life forms of plant taxa were analysed, it was determined that Hemicryptophytes have the most number of plant taxa with 82.5%, Therophytes and Geophytes have the less than others 2.5%.

1. INTRODUCTION

Turkey is a floristically rich and interesting country. It has over 12000 taxa of plants, of which around 3800 being endemics [1, 2]. Almost all plants are grown in Turkey is under negative pressure and are faced with various challenges in maintaining its generation. As long as the pressures continue, the plant faces the risk of extinction.

Due to presence of the risk factors threatening the plant species in Turkey, monitoring and finding solutions to determined problems of plants have become a necessity. Accordingly, "International Union for Conservation of Nature and Natural Resources" [1] was established to determine threat categories of endangered plants. Based on these criteria, "List of Rare,

2019 Ankara University Communications Faculty of Sciences University of Ankara Series C: Biology

Received by the editors: February 27, 2019; Accepted: March 30, 2019.

Key word and phrases: Sakarat mountain, A6, Amasya, Turkey, endemic plants, threat categories, IUCN

Threatened and Endemic Plants in Turkey" was prepared [3] and updated according to "IUCN Red List Categories" version 2.3 [4] in 2000 [5].

Sakarat Mountain, located Northeast of Central Anatolia and southeast of Amasya province, is in the transitional zone of Central and Northern Anatolia. It is also between Irano-Turanian and Euro-Siberian phytogeographical regions. The study area, located between 40°-39° latitudes and 35°-50° longitudes, is at the A6 square according to Davis's grid system. The areas between 500 m and 1956 m has a wide variety of habitats [6, 7] (Figure 1).

The data of the meteorological stations of Amasya, Tasova and Turhal (Tokat) in the work area [8] were analysed and summarized according to the method proposed by Akman and Daget [9]. The findings indicated that the region is under the influence of the Mediterranean climate [10-12]. The meteorological stations in the research area show the SWAS (Spring, Winter, Autumn, Summer) type of precipitation regime. This type of regime; The Eastern Mediterranean is characterized by the second type of precipitation regime. In addition, the least rainfall in the field of research in the Summer and Autumn seasons and all the stations in the upper cold Mediterranean climate is a common feature of the stations are seen.

In all of the Meteorology Stations, a minimum summer rainfall and a significant summer drought indicate that the region is under the influence of the Mediterranean climate.

January is the coldest month in the research area. Frost certainly occurs in Turhal in December, January and February, and in Amasya in January and February. Besides, frost is probable in Amasya in March, April, May, October, November and December, and in Tasova in November, December, January, February, March and April, and inTurhal in March, April, May, October and November. In addition, the presence of *Pinus sylvestris* forests, which are resistant to cold in these climatic conditions, is in full agreement with the climate. The vegetation in the research area shows an appropriate distribution for these climate types. The oak forests (especially *Quercus pubescens*) in the region indicate that the area is in the transition zone between Central Anatolia and Blacksea Region. Further, xerofil steppe formations are widespread in the western and southwestern parts of the region. There is a summer drought lasting for 4 or 5 months in the region. Similar studies by different researchers in different parts of Turkey were made [13-26].

In this study threat categories which apply to endemic plant species found in Sakarat Mountain (Amasya, Turkey) were determined and their assessments were made.

2. Material And Methods

The plant samples were collected at the different seasons of the years 2004 and 2005. The collected samples were deposited and saved in Herbarium of Biology Department, Faculty of Science, Ankara University (ANK). Identifications of the specimens were done according to flora studies [27-32].

12

The order of endemic taxa in Tables was given according to phylogenetical system used in Flora of Turkey. The books "2001 IUCN Red List Categories: version 3.1" [33, 34] and "Red Data Book of Turkish Plants" prepared by Ekim et al. [5] were utilized in determination of threat categories. In this study, in addition to the threat categories, phytogeographical regions/elements (chorotypes) [35, 36] and life forms [37] of the endemic plant taxa were also given.

$3. \ \mathrm{Results}$

In the result of study, totally 40 endemic taxa at the level of species, subspecies and variety, belonging to 16 families, were determined on the Sakarat Mountain. The number of endemic taxa in the families and their percentages were given in descending order in Table 1. The distribution of endemic taxa according to phytogeographic regions and their percentages were presented in Table 2, and a phylogenetic list of the endemic taxa distributing on Sakarat Mountain, threat categories, life forms and phytogeographical regions were given in Table 3. Also, the distributions of life forms of the taxa were shown in Table 4.

	Families	Number of Endemic Taxa	Percentage (%)
1	FABACEAE	6	15
2	LAMIACEAE	6	15
3	ASTERACEAE	4	10
4	BORAGINACEAE	4	10
5	PLANTAGINACEAE	4	10
6	CAMPANULACEAE	3	7.5
7	ROSACEAE	3	7.5
8	SCROPHULARIACEAE	2	5
9	BRASSICACEAE	1	2.5
10	CAPRIFOLIACEAE	1	2.5
11	CARYOPHYLLACEAE	1	2.5
12	CONVOLVULACEAE	1	2.5

TABLE 1. The distribution of endemic plant taxa into families.

13	FAGACEAE	1	2.5
14	GERANIACEAE	1	2.5
15	AMARYLLIDACEAE	1	2.5
16	IRIDACEAE	1	2.5
	Total	40	100

M. UMIT BINGOL, NESLIHAN BALPINAR, KERIM GUNEY, ARZU CANSARAN FATMAGUL GEVEN, OMER FARUK KAYA, MUNEVVER ARSLAN

TABLE 2. The distribution of endemic plant taxa according to phytogeographical regions/elements (chorotypes) [35, 36].

	Phytogeographical Regions	Number of Endemic Taxa	Percentage (%)
1	Irano-Turanian (IT)	13	32.5
2	Euro-Siberian (ES)	9	22.5
3	Mediterranean (M)	1	2.5
4	Unknown (U)	17	42.5
	Total	40	100

TABLE 3. Threat categories, phytogeographical regions/elements (chorotypes) and life forms of endemic plant taxa [33, 34].

	Families	Endemic Taxon	Chorotype	Life Form	Threat Category
1	BRASSICACEAE	Draba rigida Willd. var. rigida	U	Н	LC

2	CARYOPHYLLACE AE	Arenaria ledebouriana Fenzl var. ledebouriana	U	Th	LC
3	GERANIACEAE	Geranium cinereum Cav. subsp. subcaulescens (L'Hérit. ex DC.) Hayek var. subacutum (Boiss.) Davis & Roberts	IT	Н	LC
4	FABACEAE	Astragalus densifolius Lam. subsp. amasiensis (Freyn) Aytaç & Ekim	IT	Н	LC
5	FABACEAE	Astragalus campylosema Boiss. Subsp. campylosema	IT	Н	LC
6	FABACEAE	Lathyrus tukhtensis Czecz.	U	Н	LC
7	FABACEAE	Trifolium pannonicum Jacq. subsp. elongatum (Willd.) Zoh.	U	Н	LC
8	FABACEAE	Onobrychis bornmuelleri Freyn	U	Н	EN
9	FABACEAE	Ebenus laguroides Boiss. var. laguroides	IT	Н	LC
10	ROSACEAE	Potentilla cappadocica Boiss.	ES	Н	NT
11	ROSACEAE	Alchemilla holocycla Rothm.	IT	Н	LC
12	ROSACEAE	Crataegus tanacetifolia (Lam.) Pers.	U	Ph	LC
13	CAPRIFOLIACEAE	Lonicera caucasica Pallas subsp. orientalis (Lam.) Chamb. & Long	U	Ph	LC
14	ASTERACEAE	Helichrysum arenarium (L.) Moench subsp. aucheri (Boiss.) Davis & Kupicha	IT	Н	LC
15	ASTERACEAE	Cirsium pseudopersonata Boiss. & Bal. subsp. pseudopersonata	ES	Н	LC
16	ASTERACEAE	Jurinea pontica Hausskn. & Freyn ex Hausskn.	IT	Н	LC

M. UMIT BINGOL, NESLIHAN BALPINAR, KERIM GUNEY, ARZU CANSARAN FATMAGUL GEVEN, OMER FARUK KAYA, MUNEVVER ARSLAN

17	ASTERACEAE	Scorzonera eriophora DC.	U	Н	LC
18	CAMPANULACEAE	Campanula lyrata Lam. subsp. lyrata	U	Н	LC
19	CAMPANULACEAE	Campanula latiloba A.DC. subsp. latiloba	ES	Н	LC
20	CAMPANULACEAE	Asyneuma limonifolium (L.) Janchen subsp. pestalozzae (Boiss.) Damboldt	U	Н	LC
21	CONVOLVULACEA E	Convolvulus assyricus Griseb.	IT	Ch	LC
22	BORAGINACEAE	Onosma armenum DC.	U	Н	LC
23	BORAGINACEAE	Symphytum bornmuelleri Bucknall.	ES	Н	LC
24	BORAGINACEAE	<i>Cynoglottis chetikiana</i> Vural & Kit Tan subsp. <i>paphlagonica</i> (Hausskn. ex Bornm.) Vural & Kit Tan	U	Н	LC
25	BORAGINACEAE	Anchusa leptophylla subsp. incana Roemer & Schultes (Ledeb.) Chamb.	IT	Н	LC
26	SCROPHULARIACE AE	Verbascum ponticum (Boiss.) O. Kuntze	ES	Н	LC
27	SCROPHULARIACE AE	Verbascum abieticolum Bornm.	ES	Н	LC
28	LAMIACEAE	Scutellaria salviifolia Bentham	U	Н	LC
29	LAMIACEAE	Phlomis russeliana (Sims) Bentham	ES	Н	LC
30	LAMIACEAE	Phlomis armeniaca Willd.	IT	Н	LC
31	LAMIACEAE	Sideritis dichotoma Huter	U	Н	LC
32	LAMIACEAE	Sideritis amasiaca Bornm.	U	Н	NT
33	LAMIACEAE	Sideritis germanicopolitana Bornm. subsp. germanicopolitana	U	Н	LC
34	PLANTAGINACEAE	Chaenorhinum litorale (Bernh.) Fritch subsp. pterosporum (Fisch. & Mey.) Davis	М	Th	LC
35	PLANTAGINACEAE	Linaria corifolia Desf.	IT	Н	LC

36	PLANTAGINACEAE	Digitalis lamarckii Ivan.	IT	Н	LC
37	PLANTAGINACEAE	Veronica multifida L.	IT	Н	LC
38	FAGACEAE	Quercus macranthera Fisch. et Mey. Ex Hohen. subsp. syspirensis (C. Koch) Menitsky	U	Ph	LC
39	AMARYLLIDACEA E	Allium olympicum Boiss.	ES	G	LC
40	IRIDACEAE	Crocus speciosus Bieb. subsp. ilgazensis Mathew	ES	G	NT

16

TABLE 4. The distribution of endemic plant taxa according to life form [37].

	Life Forms	Number of Endemic Taxa	Percentage (%)
1	Hemicryptophyte (H)	32	80
2	Phanerophyte (Ph)	3	7.5
3	Geophyte (G)	2	5
4	Therophyte (Th)	2	5
5	Chamaephyte (Ch)	1	2.5
	Total	40	100

4. DISCUSSION AND CONCLUSION

There are 6 different vegetation types in Sakarat Mountain: Forest (deciduous, mixed coniferous and deciduous forest), subalpine, steppe, rocky, wet grassland and segetal vegetation. As a result of the evaluation of the samples collected from all of these vegetations, 283 genera and 494 taxa belonging to 78 families were determined. 40 of 494 taxa are endemic, and the rate of endemism is 8.09%. The number and percentage of endemic taxa according to risk categories were given in Table 5.

	Threat Categories	Number of Endemic Taxa	Percentage (%)	
1	Endangered (EN)	1	2.5	
2	Near Threatened (NT)	3	7.5	
3	Least Concern (LC)	36	90	
	Total	40	100	

TABLE 5. Number of endemic plant taxa determined in Sakarat Mountain and their threat categories.

The highest number of the endemic species was seen in Fabaceae (15%) and Lamiaceae (15%) families. When the phytogeographical distributions of the endemic taxa were examined, it was seen that Irano-Turanian elements have a rate of 32.5%, European-Siberian elements have a rate of 22.5%, Mediterranean elements have a rate of 2.5% and endemics with no phytogeographical regions have a rate of 42.5% (Table 2).

Also, the plant taxa were classified and analysed according to Raunkiaer's life forms [37]. It was shown that Hemicryptophytes are the most common (80%), and it is followed by Phanerophytes (7.5%), Geophytes (5%) and Therophytes (5%). Chamaephytes have the least number with 1 species (2.5%) (Table 4). The analysis of threat categories showed that 1 taxon (2.5%) was in Endangered, 3 taxa (7.5%) were in Near Threatened according to 2001 IUCN criteria while remaining 36 taxa (90%) were found to qualify for Least Concern category (Table 5).

We hope that the present study will contribute to future researches on rare and endemic plants of Turkey.



FIGURE 1. Map of the study area [6, 7, 26].

Acknowledgement. This study supported by TÜBİTAK (Scientific and Technical Research Council of Turkey, Project No: 1050018 TOVAG-HD) [6]. We thank the institution for their financial contributions.

References

- [1] IUCN Species Survival Commision, Draft IUCN Red List Categories. Gland, Switzerland, (1993).
- [2] A. Guner, S. Aslan, T. Ekim, M. Vural, M.T. Babac (Editors), Türkiye Bitkileri Listesi (Damarlı Bitkiler). *Nezahat Gokyigit Botanik Bahcesi ve Flora Arastırmaları Dernegi Yayını*, Istanbul, (2012).
- [3] T. Ekim, M. Koyuncu, S. Erik, R. Ilarslan, Türkiye'nin Tehlike Altındaki Nadir ve Endemik Bitkileri. *Türkiye Tabiatını Koruma Dernegi*, Ankara, 18, (1989).
- [4] IUCN Species Survival Commission, IUCN Red List Categories: Version 2.3. Gland, Switzerland, (1994).
- [5] T. Ekim, M. Koyuncu, M. Vural, H. Duman, Z. Aytac, N. Adıguzel, Türkiye Bitkileri Kırmızı Kitabı, Red Data Book of Turkish Plants (*Pteridophyta* and *Spermatophyta*). Turkish Association for the Conservation of Nature-Van Centennial University, Ankara, (2000) 246.
- [6] M.U. Bingol, F. Geven, K. Guney, Sakarat dağı (Amasya)'nın bitki ekolojisi ve bitki sosyolojisi yönünden araştırılması. TUBITAK, Proje No: TOVAG-105O018, (2005-2007) 132.

M. UMIT BINGOL, NESLIHAN BALPINAR, KERIM GUNEY, ARZU CANSARAN FATMAGUL GEVEN, OMER FARUK KAYA, MUNEVVER ARSLAN

- [7] A. Cansaran, M.U. Bingol, F. Geven, K. Guney, N. Erdogan, O.F. Kaya, Sakarat dağı florasına katkılar (Amasya-Türkiye). *BioDiCon (Biological Diversity and Conservation)*, 3(1), (2010) 103-120.
- [8] Anonim, Amasya, Taşova ve Turhal'a ait uzun yıllar rasat verileri. T.C. Çevre ve Orman Bakanlığı Devlet Meteoroloji İşleri Genel Müdürlüğü Elektronik Bilgi İşlem Merkezi (EBİM), Ankara, (2007).
- [9] Y. Akman, P.H. Daget, Quelques aspects synoptiques des climats de la Turquie. *Bull. Soc. Long. Georg.*, 5(3), (1971) 269-300.
- [10] Y. Akman, Climats et bioclimats en Turquie. *Ecologia Mediterranea*, 1(2), (1981) 73-87.
- [11] Y. Akman, O. Ketenoglu, The climate and vegetation of Turkey. In *Proceedings of the Royal Society of Edinburgh*, Section B, Biological Sciences, I.C. Hedge (Editor), Edinburgh, 89, (1986) 123-134.
- [12] Y. Akman, İklim ve biyoiklim (Biyoiklim metodları ve Türkiye iklimleri). *Palme Yayın Dağıtım*, Ankara, (1990) 319.
- [13] N. Erdogan, O. Ketenoglu, M.U. Bingol, F. Geven, M. Arslan, Evaluation of threat categories of the endemic plants of Sivrihisar mountains Eskişehir Turkey. *Anadolu Doğa Bilimleri Dergisi*, 5(1), (2014) 37-43.
- [14] M. Arslan, R. Karatas, S.T. Guner, K. Ozkan, Threat categories and endemism status of plants in the distibution areas of pulley oak in the Lakes District, *BioDiCon* (*Biological Diversity and Conservation*), 8(1), (2015) 7-15.
- [15] U. Budak, A.I. Ilbaş, Karanlıkdere vadisi (Şefaatli-Yerköy-Yozgat)'nde yayılış gösteren endemik bitkilerin tehlike kategorilerinin belirlenmesi, *Suleyman Demirel* Universitesi Fen Edebiyat Fakultesi Fen Dergisi, Konya, 24, (2004) 29-44.
- [16] E. Akciçcek, M. Vural, Kumalar dagı (Afyonkarahisar)' nın endemik ve nadir bitkileri, *Balikesir Universitesi Fen Bilimleri Enstitüsü Dergisi*, 9(2), (2007) 78-86.
- [17] R. Daskin, G. Kaynak, Conservation status of five endemic species distributed in Northwest Turkey. *Phytologia Balcanica*, 17(2), (2011) 213-219.
- [18] S.G. Senol, H. Yıldırım, A new distribution area of Asperula daphneola (Rubiaceae) in Western Turkey and it's new recommended IUCN threat category. BioDiCon (Biological Diversity and Conservation), 3(2), (2010) 123-127.
- [19] M.E. Uzunhisarcıklı, M. Vural, Taxonomy and IUCN categories of two Alcea L. (Malvaceae) species cited in the data deficient (DD) category. *BioDiCon (Biological Diversity and Conservation)*, 2(2), (2009) 90-95.
- [20] O. Yılmaz, R. Daskın, G. Kaynak, IUCN categories of three *Linum L. (Linaceae)* taxa endemic to Turkey. *BioDiCon (Biological Diversity and Conservation)*, 4(1), (2011) 144-149.
- [21] F. Celep, M. Dogan, A. Kahraman, Re-evaluated conservation status of *Salvia* (sage) in Turkey I: The Mediterranean and the Aegean geographic regions, *Turkish Journal* of Botany, 34, (2010) 201-214.
- [22] G.S. Erzurumlu, K.T. Yılmaz, Z. Sogut, A regional scale evaluation of conservation status of Orchid species recorded in The Eastern Mediterranean region of Turkey, *Turkish Journal of Agriculture-Food Science and Technology*, 5(9), (2017) 996-1001.
- [23] R. Daskin, G. Kaynak, Threat categories of three species endemic to Uludağ (Bursa/Turkey), *BioDiCon(Biological Diversity and Conservation*), 4(3),(2011) 8-13.

20

- [24] Z. Bahcecioglu, B. Yıldız, Five Critically Endangered Species in Malatya Province (Turkey), *Environment and Ecology Research*, 2(5), (2014) 206-208.
- [25] M.U. Bingol, O. Ketenoglu, F. Geven, K. Guney, N. Erdogan, M. Arslan Evaluation of threat categories of the endemic plants of Deveci mountains (Yozgat-Tokat/Turkey). XIII. Optima meeting, March 22-26 2010, Antalya, Turkey.
- [26] M.U. Bingol, A. Cansaran, F. Geven, K. Guney, N. Erdogan, O.F. Kaya, Evaluation of threat categories of the endemic plants of Sakarat mountain (Amasya/Turkey). XIII. Optima meeting, March 22-26 2010, Antalya, Turkey.
- [27] P.H. Davis, Flora of Turkey and The East Aegean Islands. Vol. 1-9. *Edinburgh* University Press, Edinburgh, (1965-1985).
- [28] P.H. Davis, R.R. Mill, K. Tan (Editors), Flora of Turkey and The East Aegean Islands (Supplement). Vol. 10. *Edinburgh University Press*, Edinburgh, (1988).
- [29] A. Guner, N. Ozhatay, T. Ekim, K.H.C. Baser (Editors), Flora of Turkey and the East Aegean Islands (Second Supplement). Vol.11, *Edinburgh University Press*, Edinburgh, (2000).
- [30] P.H. Davis, P.C. Harper, J.C. Hedge (Editors), Plant Life of South West Asia. *Botanical Society of Edinburgh*, Edinburgh, (1971) 335.
- [31] L.J. Donner, Distribution maps to P.H. Davis, "Flora of Turkey, Vol. 1-10". *Linzer Biol. Beitr.*, 22(2), (1990) 38-515.
- [32] A. Guner., S. Aslan, T. Ekim, M. Vural, M.T. Babac (Editors), Türkiye Bitkileri Listesi (Damarlı Bitkiler). *Nezahat Gökyigit Botanik Bahcesi ve Flora Arastırmaları Dernegi Yayını*, Istanbul, (2012).
- [33] IUCN Species Survival Commision, IUCN Red List Categories: Version 3.1. *Gland*, Switzerland, (2001).
- [34] IUCN Red List Categories and Criteria Version 3.1 (2001), Second edition 2012. The World Conservation Union, Switzerland and Cambridge, UK., https://www.iucn.org/knowledge/publications doc/, (2012) 32.
- [35] M. Zohary, Geobotanical Foundations of the Middle East, Vol. 1-2. *Gustav Fischer Verlag*, Stuttgart, (1973) 739.
- [36] A. Takhtajan, Floristic Region of The World. *Univ. of California Press*, Los Angeles, (1986). 102.
- [37] C. Raunkiaer, The Life Forms of Plants and Statistical Plant Geography. *Oxford University Press*, London, Great Britain, (1934) 632.

Current Address: M. UMIT BINGOL (Corresponding author): Ankara University, Science Faculty, Biology Department, 06100, Tandoğan, Ankara, Turkey. *E-mail: <u>mumit1111@hotmail.com</u>* ORCID: https://orcid.org/0000-0002-1342-3892

Current Address: NESLIHAN BALPINAR: Burdur Mehmet Akif Ersoy University, Faculty of Arts and Sciences, Department of Biology, 15030, Burdur, Turkey. *E-mail: <u>nerdogan@mehmetakif.edu.tr</u> ORCID: https://orcid.org/0000-0002-4469-8629* Current Address: KERIM GUNEY: Kastamonu University, Faculty of Forestry, Department of Forest Engineering, 37150, Kastamonu, Turkey. *E-mail: <u>kguney@kastamonu.edu.tr</u> ORCID: https://orcid.org/0000-0003-2305-790X*

Current Address: ARZU CANSARAN: Amasya University, Faculty of Education, Department of Mathematics and Science Education, Amasya, Turkey. *E-mail: arzu.cansaran@amasya.edu.tr ORCID: https://orcid.org/ 0000-0002-0912-147X*

Current Address: FATMAGUL GEVEN: Ankara University, Science Faculty, Biology Department, 06100, Tandoğan, Ankara, Turkey. *E-mail: geven@science.ankara.edu.tr ORCID: https://orcid.org/0000-0003-4231-406X*

Current Address: OMER FARUK KAYA: Harran University, Science and Arts Faculty, Department of Biology, 63190 Şanlıurfa, Turkey. *E-mail: phytosociologist@gmail.com ORCID: https://orcid.org/0000-0002-0851-0617*

Current Address: MUNEVVER ARSLAN: Ministry of Forestry and Water Works, Research Institute for Forest, Soils & Ecology, 61-26160, Eskişehir, Turkey. *E-mail:* <u>mune28@hotmail.com</u> ORCID: https://orcid.org/0000-0003-2645-1486