

PAPER DETAILS

TITLE: Spectroscopic, Electronic and Non-Linear Properties of
2-[3-Phenyl-4,5-dihydro-1H-1,2,4-triazol-5-one-4-yl]-phenoxyacetic Acide

AUTHORS: Murat BEYTUR, Haydar YUKSEK

PAGES: 162-170

ORIGINAL PDF URL: <http://www.epstem.net/tr/download/article-file/802744>

Spectroscopic, Electronic and Non-Linear Properties of 2-[3-Phenyl-4,5-dihydro-1*H*-1,2,4-triazol-5-one-4-yl]-phenoxyacetic Acide

Murat BEYTUR
Kafkas University

Haydar YUKSEK
Kafkas University

Abstract: In this study, 2-[3-phenyl-4,5-dihydro-1*H*-1,2,4-triazol-5-one-4-yl]-phenoxyacetic acid was optimized by using B3LYP/6-311G+(d,p) and HF/6-311G+(d,p) basis sets. Firstly, calculated IR data of compound were calculated in gas phase by using of 6-311G+(d,p) basis sets of B3LYP and HF methods and are multiplied with appropriate adjustment factors. Theoretical infrared spectrums are formed from the data obtained according to B3LYP and HF methods. Experimental IR values and theoretical values were compared. In the identification of calculated IR data was used the veda4f program. Also, ¹H-NMR and ¹³C-NMR spectral data values were calculated according to the method of GIAO using the program package Gaussian G09W Software. Experimental and theoretical values were inserted into the graphic according to equation of $\delta_{exp} = a + b \cdot \delta_{calc}$. The standard error values were found via SigmaPlot program with regression coefficient of a and b constants. Furthermore, molecular structure, HOMO and LUMO energy analysis, total static dipol moment (μ), the mean polarizability ($\langle\alpha\rangle$), the anisotropy of the polarizability ($\Delta\alpha$), the mean first-order hyperpolarizability ($\langle\beta\rangle$), electronegativity (χ), hardness (η), molecular electrostatic potential maps (MEP) and Mulliken charges of 2-[3-phenyl-4,5-dihydro-1*H*-1,2,4-triazol-5-one-4-yl]-phenoxyacetic acid molecule have been investigated by using B3LYP and HF levels with the 6-311G+(d,p) basis set.

Keywords: 1,2,4-Triazol-5-one, GIAO, Non-Linear Properties, Electronegativity HOMO and LUMO

Introduction

Triazole is an unsymmetrical heterocyclic organic compound having three nitrogen atoms in the five-membered ring. 1,2,4-Triazole and 4,5-dihydro-1*H*-1,2,4-triazol-5-one derivatives are reported to possess a broad spectrum of biological activities such as antimicrobial, antifungal, antitumor, anti-HIV, antiviral, anticancer, anti-inflammatory, analgesic and antioxidant properties (Alkan et al., 2007; Hashem et al., 2007; Aytac et al., 2009; Bayrak et al., 2010; Gürsoy Kol and Yüksek, 2010; Güzel demirci and Kucukbasmaci, 2010; Sancak et al., 2010; Beytur et al., 2019; Turhan Irak and Beytur, 2019). Also, several articles reporting the synthesis of some *N*-arylideneamino-4,5-dihydro-1*H*-1,2,4-triazol-5-one compounds and derivatives have been published (İkizler and Yüksek, 1994; Bahçeci et al., 2002; Yüksek et al., 2004; Yüksek et al., 2005; Yüksek et al., 2006; Turhan Irak and Gümüş, 2017).

In this paper, the optimized molecular structure, vibrational frequencies, spectroscopic parameters, atomic charges and frontier molecule orbitals (HOMO and LUMO) of the 2-[3-phenyl-4,5-dihydro-1*H*-1,2,4-triazol-5-one-4-yl]-phenoxyacetic acid have been calculated by using DFT/B3LYP and HF method with 6-311G+(d,p) basis set. All quantum chemical calculations were carried out by using Gaussian 09W (Frisch et al., 2009; Wolinski, Hilton & Pulay, 1990) program package and the GaussView molecular visualization program (Frisch, Nielson & Holder, 2003). The molecular structure and vibrational calculations of the molecule were computed by using Becke-3-Lee Yang Parr (B3LYP) (Becke, 1993; Lee et al., 1998) density functional method and Hatree-Fock with 6-311G+(d,p) basis set in ground state. IR absorption frequencies of analyzed molecule were calculated by two methods. Then, they were compared with experimental data (Çiftçi et al., 2018), which are shown to be accurate. The assignments of fundamental vibrational modes of the title molecule were performed

- This is an Open Access article distributed under the terms of the Creative Commons Attribution-Noncommercial 4.0 Unported License, permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

- Selection and peer-review under responsibility of the Organizing Committee of the Conference

on the basis of total energy distribution (TED) analysis by using VEDA 4f program (Jamroz, 2004). Furthermore, molecular structure, HOMO and LUMO energy analysis, total static dipole moment (μ), the mean polarizability ($\langle\alpha\rangle$), the anisotropy of the polarizability ($\Delta\alpha$), the mean first-order hyperpolarizability ($\langle\beta\rangle$), electronegativity (χ), hardness (η), molecular electrostatic potential maps (MEP) and Mulliken charges of titled compound have been investigated by using B3LYP and HF levels with the 6-311G+(d,p) basis set.




Figure 1. The optimized molecular structure (Gaussview Appearance) of 2-[3-phenyl-4,5-dihydro-1H-1,2,4-triazol-5-one-4-yl]-phenoxyacetic acid with DFT/B3LYP 6-311G+(d,p) level.

Method

The molecular structure of the title compound in the ground state is computed by performing both the density functional theory (DFT) and Hartree-Fock (HF) (Becke, 1993; Lee, 1998) at 6-311G+(d,p) level. Density functionals for all studies reported in this paper have been in the following form

$$E_{XC} = (1 - a_0)E_X^{LSDA} + a_0E_X^{HF} + a_X\Delta E_X^{B88} + a_CE_C^{LYP} + (1 - a_C)E_C^{VWN}$$

where the energy terms are the Slater exchange, the Hartree-Fock exchange, Becke's exchange functional correction, the gradient corrected correlation functional of Lee, Yang and Parr, and the local correlation functional of Vosko, Wilk and Nusair (Vosko et al., 1980). The theoretical geometric structure of the title compound is given in Figure 1. Molecular geometry is restricted and the optimized geometrical parameters of the title compound in this study are carried out by using Gaussian 09W program package (Frisch et al., 2009) and the visualization parts were done with GaussView program (Dennington et al., 2009) on personal computer employing 6-311G+(d,p) basis set. Additionally, harmonic vibrational frequencies for the title compound are calculated with these selected methods and then scaled by 0.9516 and 0.9905, respectively (Avcı and Atalay, 2008) and these results were compared with the experimental data (Çiftçi et al., 2018).

Results and Discussion

Analysis of vibrational modes

In spectroscopic field, the vibrational spectra of substituted benzene derivatives have been greatly investigated by various spectroscopic, since the single substitution can have a tendency to put greater changes in vibrational wavenumbers of benzene (Tereci et al., 2012; Pir et al., 2013). In other words, molecular system of benzene is greatly affected by the nature of substituents. The number of potentially active fundamentals of non-linear molecule which have N atoms is equal to (3N-6) apart from three translational and three rotational degrees of freedom. The title molecule contains 39 atoms and 111 normal vibration modes have C1 symmetry (Table 1).

Experimentally (Çiftçi et al., 2018), the investigated titled compound, as expected the IR spectra data, the N-H stretching vibration at 3179 cm^{-1} and two C=O peak at 1708 cm^{-1} range was observed. In addition, C=N stretching vibration at 1597 cm^{-1} and COO stretching vibrations at 1229 cm^{-1} are occurred. Theoretically and experimentally (Çiftçi et al., 2018), the calculated vibrational frequencies for the 2-[3-phenyl-4,5-dihydro-1H-1,2,4-triazol-5-one-4-yl]-phenoxyacetic acid are summarized in Table 1. Furthermore the experimental IR (Çiftçi et al., 2018) and simulated spectra by using B3LYP/6-311G+(d,p) and HF/6-311G+(d,p) levels of the titled compound under investigation are given in Figure 2.

Table 1. The calculated frequencies values of the titled compound

Selected Vibrational Types	Experim.	Scaled DFT	Scaled HF
$\delta\text{ O}_{37}\text{C}_5\text{C}_6$ (23)	423	439	447
$\tau\text{ C}_1\text{C}_{13}\text{C}_{17}\text{C}_{12}$ (21)	484	502	507
$\delta\text{ C}_{12}\text{O}_{37}\text{C}_5$ (12), $\tau\text{ C}_5\text{C}_6\text{C}_7\text{C}_8$ (32), $\tau\text{ O}_{37}\text{C}_4\text{C}_6\text{C}_5$ (10), $\tau\text{ H}_{20}\text{C}_6\text{C}_7\text{C}_8$ (12)	544	572	582
$\delta\text{ O}_{39}\text{C}_{11}\text{O}_{38}$ (34), $\delta\text{ O}_{37}\text{C}_5\text{C}_6$ (10), $\delta\text{ C}_{11}\text{C}_{10}\text{O}_{37}$ (20)	576	587	602
$\delta\text{ C}_{15}\text{C}_{14}\text{C}_{13}$ (30), $\delta\text{ O}_{36}\text{C}_2\text{N}_{33}$ (18)	596	614	623
$\tau\text{ O}_{36}\text{N}_{33}\text{N}_{34}\text{C}_2$ (73)	678	714	734
$\tau\text{ H}_{27}\text{C}_{13}\text{C}_{14}\text{C}_{15}$ (39), $\tau\text{ N}_{32}\text{N}_{34}\text{C}_{12}\text{C}_1$ (10), $\tau\text{ C}_1\text{C}_{13}\text{C}_{17}\text{C}_{12}$ (19)	758	760	787
$\tau\text{ H}_{20}\text{C}_6\text{C}_7\text{C}_8$ (36)	802	831	859
$\delta\text{ C}_5\text{C}_6\text{C}_7$ (38), $\delta\text{ C}_4\text{C}_3\text{N}_{35}$ (11)	839	877	899
$\delta\text{ H}_{27}\text{C}_{13}\text{C}_{14}$ (23), $\tau\text{ H}_{27}\text{C}_{13}\text{C}_{14}\text{C}_{15}$ (35), $\tau\text{ H}_{31}\text{C}_{17}\text{C}_{16}\text{C}_{15}$ (28)	880	913	948
$\tau\text{ H}_{19}\text{C}_3\text{N}_{35}\text{N}_{34}$ (79)	943	980	1013
$\nu\text{ O}_{37}\text{C}_{10}$ (40)	1028	1039	1061
$\nu\text{ C}_{14}\text{C}_{13}$ (24), $\delta\text{ H}_{24}\text{C}_{13}\text{C}_{14}$ (22), $\nu\text{ N}_{33}\text{N}_{32}$ (14)	1066	1079	1082
O_{37}C_5 (25), $\text{C}_{10}\text{C}_{11}$ (10), $\delta\text{ H}_{26}\text{O}_{38}\text{C}_{11}$ (41)	1111	1141	1106
$\delta\text{ H}_{24}\text{C}_{10}\text{O}_{37}$ (12), $\nu\text{ O}_{37}\text{C}_5$ (19)	1168	1199	1201
$\delta\text{ H}_{24}\text{C}_{10}\text{O}_{37}$ (51)	1229	1267	1265
$\delta\text{ H}_{19}\text{C}_3\text{N}_{35}$ (18), $\delta\text{ H}_{18}\text{N}_{33}\text{N}_{32}$ (40)	1354	1367	1405
$\delta\text{ H}_{20}\text{C}_6\text{C}_7$ (11), $\delta\text{ H}_{27}\text{C}_{13}\text{C}_{14}$ (11)	1423	1434	1457
$\nu\text{ N}_{32}\text{C}_1$ (29), $\nu\text{ C}_{14}\text{C}_{13}$ (12)	1488	1542	1588
$\nu\text{ N}_{32}\text{C}_1$ (40), $\nu\text{ C}_5\text{C}_6$ (12)	1597	1602	1704
$\nu\text{ O}_{39}\text{C}_{11}$ (86), $\nu\text{ O}_{36}\text{C}_2$ (74), $\nu\text{ N}_{33}\text{C}_2$ (11)	1708	1753	1805
$\nu\text{ C}_{10}\text{H}_{24}$ (99)	2791	2973	2935
$\nu\text{ C}_{10}\text{H}_{24}$ (99)	2916	3016	2982
$\nu\text{ C}_3\text{H}_{27}$ (77)	2936	3068	3013
$\nu\text{ C}_{13}\text{H}_{27}$ (53)	3011	3089	3036
$\nu\text{ C}_{17}\text{H}_{31}$ (83)	3076	3102	3048
$\nu\text{ N}_{33}\text{H}_{18}$ (100)	3179	3558	3556
$\nu\text{ O}_{38}\text{H}_{26}$ (100)	3208	3632	3722




Figure 2. IR spectra simulated with DFT/B3LYP/6-311G+(d,p) and HF/6-311G+(d,p) levels of the titled compound

NMR spectral analysis

the isotropic chemical shift analysis allows us to identify relative ionic species and to calculate reliable magnetic properties in nuclear magnetic resonance (NMR) spectroscopy which provide the accurate predictions of molecular geometries, (Wade, 2006; Rani, et al., 2010; Subramanian et al., 2010). For this purpose, the optimized molecular geometry of the 2-[3-phenyl-4,5-dihydro-1*H*-1,2,4-triazol-5-one-4-yl]-phenoxyacetic acid was obtained by using B3LYP and HF methods with 6-311G+(d,p) basis level in DMSO solvent. By considering the optimized molecular geometry of the titled compound. The ^1H and ^{13}C NMR chemical shift values were calculated at the same level by using Gauge-Independent Atomic Orbital method (Table 2). Theoretically and experimentally values (Çiftçi et al., 2018) were plotted according to $\delta_{\text{exp}} = a \cdot \delta_{\text{calc}} + b$, Eq. a and b constants regression coefficients with a standard error values were found using the SigmaPlot program. The correlation graphics are given Figure 3 and the linear correlation data of the titled compound by considering the results are given in Table 2. Therefore the (R^2) values (DFT/HF) for ^1H NMR (DMSO) and ^{13}C NMR (DMSO) chemical shifts in different solvents has been found as 0.2228 and 0.98849 for the 2-[3-phenyl-4,5-dihydro-1*H*-1,2,4-triazol-5-one-4-yl]-phenoxyacetic acid (Figure 3). In our study, the ^1H -NMR spectrum of the titled compound was observed belong to H181 proton peak at 12.38 ppm because acidic show feature (Yüksek, 1992; Yüksek et al., 2005; Yüksek et al., 2006, Çiftçi et al., 2018). H19 protons were observed at 10.01 ppm. Therotically, DMSO solvent these values for the mentioned proton atoms were found as 8.05/7.15 and 10.54/9.53 ppm, respectively. In Table 2, the biggest ^{13}C chemical shift value of the molecule are observed at 170.38 ppm for the C11 carbon atom double bounded to the oxygen in carbonyl group (Anderson et al., 2004). DMSO solvent the calculated ppm values (DFT/HF) for C11 carbon atom were theoretically found as 179.70/163.08 ppm. Additionally, due to the electronegative property of nitrogen atoms in molecule, the experimental NMR chemical shift values for C1 and C2 carbon atom the bounded to nitrogen atoms in 1,2,4-triazol ring and C3 carbon atom with sp^2 hybride are observed at 145.16, 151.83 and 152.17 ppm, respectively.

Table 2. The calculated and experimental ^1H and ^{13}C NMR isotropic chemical shifts of the titled compound

	Experiment.	DFT/631d/DMSO	Diff./DMSO	HF/631d/DMSO	Diff./DMSO
C1	145,16	153,48	-8,32	143,66	1,50
C2	151,83	155,64	-3,81	144,63	7,20
C3	152,17	158,19	-6,02	149,49	2,68
C4	122,19	132,27	-10,08	116,63	5,56
C5	157,78	165,80	-8,02	150,23	7,55
C6	121,89	130,72	-8,83	116,79	5,10
C7	130,53	137,77	-7,24	128,76	1,77
C8	113,43	130,09	-16,66	116,97	-3,54
C9	133,55	141,78	-8,23	131,50	2,05
C10	65,36	73,69	-8,33	52,62	12,74
C11	170,38	179,70	-9,32	163,08	7,30
C12	126,30	132,98	-6,68	119,42	6,88
C13	127,19	133,01	-5,82	123,99	3,20
C14	128,48	133,44	-4,96	121,29	7,19
C15	128,97	135,74	-6,77	126,46	2,51
C16	126,30	132,83	-6,53	120,21	6,09
C17	128,48	134,80	-6,32	126,34	2,14
H18	12,38	8,05	4,33	7,15	5,23
H19	10,01	10,54	-0,53	9,53	0,48
H20	7,56	7,88	-0,32	7,45	0,11
H21	7,49	7,68	-0,19	7,48	0,01
H22	7,53	7,57	-0,04	7,18	0,35
H23	7,06	7,85	-0,79	7,61	-0,55
H24	4,88	3,95	0,93	2,92	1,96
H25	4,88	4,19	0,69	3,01	1,87
H26	13,15	6,60	6,55	5,68	7,47
H27	7,85	8,26	-0,41	7,91	-0,06
H28	7,55	7,85	-0,30	7,45	0,10
H29	7,54	7,72	-0,18	7,61	-0,07
H30	7,07	7,68	-0,61	7,48	-0,41
H31	7,91	8,56	-0,65	8,00	-0,09




Figure 3. The correlation graphics for ^{13}C -NMR (a), ^1H -NMR (b), ^{13}C -NMR (DMSO) (c) and ^1H -NMR (DMSO) (d), chemical shifts of the titled compound

Frontier molecular orbital analysis

The energies of two important molecular orbitals of the title molecule; the second highest and highest occupied MO's (HOMO), the lowest and the second lowest unoccupied MO's (LUMO) were calculated by using DFT/B3LYP and HF methods with 6-311G+(d,p) level and are presented in Table 3. The energy gap of the title molecule was calculated at DFT/B3LYP and HF level, which reveals the chemical reactivity and proves the occurrence of eventual charge transfer. The HOMO is located almost over the carbon atoms, oxygen atoms and also slightly delocalized in hydrogen atom and the LUMO is mainly delocalized in carbon atoms of benzene ring. The energy gap (energy difference between HOMO and LUMO orbital) is a critical parameter in determining molecular electrical transport properties (Fukui, 1982). The HOMO-LUMO energy gap of the title molecule is found to 4.272/2.752 eV. obtained at DFT/HF method with 6-311G+(d,p) basis set.

Table 3. The calculated HOMO-LUMO energies of the titled compound (3) according to DFT/B3LYP/6-31G(d) and HF/B3LYP/6-31G(d) levels

Electronic properties	B3LYP	HF
I; Ionization Potential (eV)	6.145	8.430
A; Electron Affinity (eV)	1.873	5.679
η ; Chemical Hardness (eV)	2.136	1.376
S; Molecular Softness (eV)	1.068	0.689
χ ; Electronegativity (eV)	4.009	7.05
Total Energy (a.u.)	-1497	-1492
ΔE ; Energy Gap (eV)	4.272	2.752

Mulliken's atomic charges

The Mulliken atomic charges at the HF/6-311G+(d,p) and B3LYP/6-311G+(d,p) level of 2-[3-phenyl-4,5-dihydro-1H-1,2,4-triazol-5-one-4-yl]-phenoxyacetic acid in gas phase are given in Table 4 (Mulliken, 1955). The electronegative N32, N33, N34, N35, O36, O37, O38 and O39 atoms of the titled compound have negative atomic charge values. The Mulliken atomic charges (B3LYP/HF) of the mentioned atoms were calculated as -0,18358/-0,06376, -0,18470/-0,38072, -0,18497/-0,19111, 0,04705/-0,03712, -0,36908/-0,49686, 0,07053/0,05755, -0,18642/-0,30668 and -0,26420/-0,36914 a.u., respectively. The C2, C3, C4, C6, C13 and C16 carbon atoms bounded to the mentioned electronegative atoms in the molecule under study have positive atomic charge values. The values of the positive charges of the mentioned carbon atoms were found as 0,54508/0,77855, 0,13796/0,47904, 0,88901/1,20224, 0,03142/0,20002, 0,58601/0,74340 and 0,02243/0,47360 a.u., respectively. Therefore the C1 atom surrounded with two electronegative N32 and N34 atoms. In the titled compound the atomic charges of all hydrogen atoms have positive values.

Table 4. Mulliken atomic charges of the titled compound

	B3LYP	HF		B3LYP	HF		B3LYP	HF
1C	-0,08604	-0,04616	14C	-0,02262	-	27H	0,31520	0,17005
	0,54508	0,77855		-0,21346	-		0,13673	0,14209
2C	0,13796	0,47904	15C	-0,52235	-	28H	0,12864	0,14123
3C	0,88901	1,20224	16C	-0,45032	-	29H	0,13096	0,14147
4C	-0,47927	-1,03838	17C	0,00277	0,00277	30H	0,14235	0,15549
5C	-0,20818	-0,30140	18H	0,33533	0,37704	31H	-	-
6C	-0,23333	-0,32372		0,17444	0,20726	32N	0,18358	0,06376
			19H	0,14732	0,16557		-	-
7C	-0,20602	-0,38056	20H	0,13554	0,14630	33N	0,18470	0,38072
8C	-0,60935	-0,47661	21H	0,13182	0,14150		-	-
9C	-0,54042	-0,41960	22H	0,13945	0,14896	34N	0,18497	0,19111
10C	0,03142	0,20002	23H	0,22609	0,24619	35N	0,04705	-
11C	0,58601	0,74340	24H	0,19323	0,19155		-	-
12C	0,02243	-0,47360	25H	0,27772	0,30292	36O	0,18642	0,30668
13C			26H			37O	-	-
						38O	0,26420	0,36914
						39O		

Nonlinear Optic Properties

The materials having nonlinear activity possess a nonlinear response to the electric fields associated with the light of a laser beam. It is well known that the higher values of dipole moment, polarizability, and hyperpolarizability are important for more active NLO properties. In this study dipole moment, polarizability and first hyperpolarizability of the titled molecule were investigated by using B3LYP method at 6-311G+(d,p) basis set. The following formulas are used for calculating the magnitude of total static dipole moment (μ), polarizability (α) and first hyperpolarizability (β):

$$\mu = \left(\mu_x^2 + \mu_y^2 + \mu_z^2 \right)^{\frac{1}{2}}$$

$$\alpha = \frac{1}{3} (\alpha_{xx} + \alpha_{yy} + \alpha_{zz})$$

$$\beta = \sqrt{(\beta_{xxx} + \beta_{yyy} + \beta_{zzz})^2 + (\beta_{yyy} + \beta_{xxy} + \beta_{yzz})^2 + (\beta_{zzz} + \beta_{xxz} + \beta_{yyz})^2}$$

Where, the total static dipole moment (μ), linear polarizability (α) and the first hyperpolarizability (β) using the x, y, z components are defined as (Abraham et al., 2008; Karamanis et al., 2008). The energy gap ΔE_g , dipole moment (μ), linear polarizability (α) and the first hyperpolarizability (β) values of conformer ct of the titled molecule are investigated as a function of the two torsional angle using B3LYP/6-311G+(d,p) level of theory (Govindarajan et al., 2012) (Table 5).

Tablo 5. Calculated polarization and hyperpolarizability values of the molecule (B3LYP/6-311G+(d,p) and HF/6-311G+(d,p))

	B3LYP	HF		B3LYP	HF
μ_x	-2.2250 debye	-3.8219 debye	β_{xxx}	7774.61 a.u.	3821.35 a.u.
μ_y	6.7852 debye	-0.8012 debye	β_{xyy}	1053.82 a.u.	538.44 a.u.
μ_z	2.3422 debye	2.0819 debye	β_{yyy}	823.08 a.u.	495.47 a.u.
μ_{Toplam}	7.2818 debye	4.4253 debye	β_{yyz}	-340.62 a.u.	161.42 a.u.
α_{xx}	74.364 a.u.	60.185 a.u.	β_{xzz}	-1001.26 a.u.	-903.63 a.u.
α_{yy}	41.696 a.u.	37.522 a.u.	B_{xyz}	-154.56 a.u.	-181.90 a.u.
α_{zz}	33.325 a.u.	30.965 a.u.	B_{yyz}	-309.57 a.u.	-202.32 a.u.
α	49.795x10 ⁻²⁴ esu	42.891x10 ⁻²⁴ esu	β_{zzz}	-295.06 a.u.	87.33 a.u.
$\Delta\alpha$	37.560x10 ⁻²⁴ esu	26.556x10 ⁻²⁴ esu	β_{yzz}	669.31 a.u.	523.13 a.u.
β_x	-9651.51 a.u.	4855.26 a.u.	B_{zzz}	1193.83 a.u.	866.30 a.u.
β_y	-1496.43 a.u.	-944.53 a.u.	B	8.567 ⁻³⁰ esu	5.006x10 ⁻³⁰ esu
β_z	1568.10 a.u.	1476.75 a.u.			

Conclusion

In this paper, the structure of the titled compound is characterized by using ¹H, ¹³C NMR and FT-IR spectroscopic methods. The molecular structures, vibrational frequencies, ¹H and ¹³C NMR chemicals shifts, UV-vis spectroscopies, HOMO and LUMO analyses and atomic charges of 2-[3-phenyl-4,5-dihydro-1H-1,2,4-triazol-5-one-4-yl]-phenoxyacetic acid obtained have been calculated by using DFT/B3LYP and HF methods. By considering the results of experimental works it can be easily stated that the ¹H and ¹³C NMR chemical shifts, and vibrational frequencies spectroscopic parameters obtained theoretically are in a very good agreement with the experimental data. Also, the electronic structure of titled compound are determined electronic structure identifiers such as the Energy of the Highest Occupied Molecular Orbital, Energy of the Lowest Unoccupied Molecular Orbital, molecular hardness, chemical softness, electronegativity, chemical potential, electrophilicity index, nucleophilicity index and dipole moment. Finally, in this study, The nonlinear optical properties of the compound were calculated theoretically. It was found that the molecule concerned had a higher hyperpolarizability value than urine (0.77x10⁻³⁰esu).

References

- Abraham, J. P., Sajan, D., Hubert, Joe I. H., & Jayakumar, V. S. (2008). Molecular structure, spectroscopic studies and first-order molecular hyperpolarizabilities of p-amino acetanilide, *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 71 (2), 355-367.
- Alkan, M., Yuksek, H., Islamoglu, F., Bahceci, S., Calapoglu, M., Elmastas, M., Aksit, H., & Ozdemir, M. (2007). A study on 4-acylamino-4,5-dihydro-1H-1,2,4-triazol-5-ones, *Molecules*, 12 (8), 1805-1816.
- Anderson, R. J., Bendell, D. J., & Groundwater, P. W. (2004). Organic Spectroscopic Analysis, *The Royal Society of Chemistry*: Sanderland, UK.
- Aytac, S. P., Tozkoparan, B., Kaynak, F. B., Aktay, G., Goktas, O., & Unuvar, S. (2009). Synthesis of 3,6-disubstituted 7H-1,2,4-triazolo[3,4-b]-1,3,4-thiadiazines as novel analgesic/anti-inflammatory compounds, *European Journal of Medicinal Chemistry*, 44 (11), 4528-4538.
- Avci, D., & Atalay, Y. (2008). Theoretical analysis of vibrational spectra and scaling-factor of 2-aryl-1,3,4-oxadiazole derivatives, *International Journal of Quantum Chemistry*, 109, 328-341.
- Bahceci, S., Yuksek, H., Ocak, Z., Koksall, C., & Ozdemir, M. (2002). Synthesis and non-aqueous medium titrations of some new 4,5-dihydro-1H-1,2,4-triazol-5-one derivatives. *Acta Chimica Slovenica*, 49, 783-794.
- Bayrak, H., Demirbas, A., Demirbas, N., & Karaoglu, S. A. (2010). Cyclization of some carbothioamide derivatives containing antipyrine and triazole moieties and investigation of their antimicrobial activities, *European Journal of Medicinal Chemistry*, 45 (11), 4726-4732.
- Becke A. D. (1993). Density functional thermochemistry. III. The role of exact Exchange, *The Journal of Chemical Physics*, 98, 5648-5652.

- Beytur, M., Turhan Irak, Z., Manap, S., and Yüksek, H. (2019). Synthesis, Characterization and Theoretical Determination of Corrosion Inhibitor Activities of Some New 4,5-Dihydro-1H-1,2,4-Triazol-5-one Derivatives, *Heliyon*, 5 (6), e01809, 1-8.
- Çiftçi, E., Beytur, M., Calapoğlu, M., Gürsoy Kol, Ö., Alkan, M., Toğay, V. A., Manap, S., & Yüksek, H. (2018). Synthesis, Characterization, Antioxidant and Antimicrobial Activities and DNA Damage of Some Novel 2-[3-alkyl (aryl)-4,5-dihydro-1H-1,2,4-triazol-5-one-4-yl]-phenoxyacetic acids in Human Lymphocytes, *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 9 (5), 1760-1771.
- Dennington, R., Keith, T., & Millam, J. (2009). GaussView, Version 5, Semiche Inc., Shawnee Mission KS.
- Frisch, A., Nielson, A.B., & Holder, A.J. (2003). Gaussview User Manual, Gaussian, Inc., Wallingford, CT,
- Frisch, M. J., Trucks, G., Schlegel, H. B., Scuseria, G. E., Robb, M. A., Cheeseman, J. R., Scalmani, G., Barone, V., Mennucci, B., Petersson, G. A., Nakatsuji, H., Caricato, M., Li, X., Hratchian, H. P., Izmaylov, A. F., Bloino, J., Zheng, G., Sonnenberg, J. L., Hada, M., Ehara, M., Toyota, K., Fukuda, R., Hasegawa, J., Ishida, M., Nakajima, T., Honda, Y., Kitao, O., Nakai, H., Vreven, T., Montgomery, J. A., Jr. Vreven, T., Peralta, J. E., Ogliaro, F., Bearpark, M., Heyd, J. J., Brothers, E., Kudin, N., Staroverov, V. N., Kobayashi, R., Normand, J., Raghavachari, K., Rendell, A., Burant, J. C., Iyengar, S. S., Tomasi, J., Cossi, M., Rega, N., Millam, J. M., Klene, M., Knox, J. E., Cross, J. B., Bakken, V., Adamo, C., Jaramillo, J., Gomperts, R., Stratmann, R. E., Yazyev, O., Austin, A. J., Cammi, R., Pomelli, C., Ochterski, J. W., Martin, L. R., Morokuma, K., Zakrzewski, V. G., Voth, G. A., Salvador, P., Dannenberg, J. J., Dapprich, S., Daniels, A. D., Farkas, O., Foresman, J. B., Ortiz, J. V., Cioslowski, J. & Fox, D. J. (2009). Gaussian Inc., Wallingford, CT.
- Govindarajan, M., Periandy, S., & Carthigayen, K. (2012). FT-IR and FT-Raman spectra, thermo dynamical behavior, HOMO and LUMO, UV, NLO properties, computed frequency estimation analysis and electronic structure calculations on α -bromotoluene. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 97, 411-422.
- Gürsoy Kol, Ö., and Yüksek, H. (2010). Synthesis and invitro antioxidant evaluation of some novel 4,5-dihydro-1H-1,2,4-triazol-5-one derivatives, *E-Journal of Chemistry*, 7 (1), 123-136.
- Guzeldemirci, N. U., & Kucukbasmaci, O. (2010). Synthesis and antimicrobial activity evaluation of new 1,2,4-triazoles and 1,3,4-thiadiazoles bearing imidazo[2,1-b]thiazole moiety, *European Journal of Medicinal Chemistry*, 45 (1), 63-68.
- İkizler, A. A., & Yüksek, H. (1994). Reaction of 4-amino-4,5-dihydro-1H-1,2,4-triazol-5-ones with 2,5-dimetoxytetrahydrofuran, *Collection of Czechoslovak Chemical Communications*, 59 (3), 731-735.
- Jamróz, M. H. (2004). Vibrational Energy Distribution Analysis: VEDA 4 program, Warsaw.
- Karamanis, P, Pouchan, C., & Maroulis, G., (2008). Structure, stability, dipole polarizability and differential polarizability in small gallium arsenide clusters from all-electron ab initio and density-functional-theory calculations, *Physical Review A*, 77, 013201-013208.
- Lee, S. Y. (1998). Molecular Structure and Vibrational Spectra of Biphenyl in the Ground and the lowest Triplet States. Density Functional Theory Study, *Bulletin of the Korean Chemical Society*, 19 (1), 93-98.
- Mulliken, R. S., (1955). Electronic population analysis on LCAO–MO molecular wave functions. *Journal of Chemical Physics*, 23, 1833–1840.
- Pir, H., Günay, N., Tamer, Ö., Avcı, D., & Atalay, Y. (2013). Theoretical investigation of 5-(2-Acetoxyethyl)-6-methylpyrimidin-2, 4-dione: Conformational study, NBO and NLO analysis, molecular structure and NMR spectra, *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 112, 331-342.
- Rani, A. U., Sundaraganesan, N., Kurt, M., Çınar, M., & Karabacak, M. (2010). FT-IR, FT-Raman, NMR spectra and DFT calculations on 4-chloro-N-methylaniline, *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 75, 1523-1529.
- Sancak, K., Unver, Y., Kazak, C., Dugdu, E., and Arslan, B. (2010). Synthesis and characterizations of some new 2,4-dihydro-[1,2,4]-triazol-3-one derivatives and X-ray crystal structures of 4-(3-phenylallylideneamino)-5-thiophen-2-yl-methyl-2,4-dihydro-[1,2,4]triazol-3-one, *Turkish Journal of Chemistry*, 34 (5), 771-780.
- Subramanian, N., Sundaraganesan, N., & Jayabharathi, J. (2010). Molecular structure, spectroscopic (FT-IR, FT-Raman, NMR, UV) studies and first-order molecular hyperpolarizabilities of 1,2-bis(3-methoxy-4-hydroxybenzylidene)hydrazine by density functional method, *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 76, 259-269.
- Turhan Irak, Z., & Beytur, M. (2019). 4-Benzilidenamino-4,5-dihidro-1H-1,2,4-triazol-5-on Türevlerinin Antioksidan Aktivitelerinin Teorik Olarak İncelenmesi, *Iğdır University Journal of the Institute of Science and Technology*, 9 (1), 512-521.
- Yüksek, H. (1992). 3-Alkil(aril)-4-amino-4,5-dihidro-1,2,4-triazol-5-on'ların bazı reaksiyonlarının incelenmesi. Doktora Tezi, KTÜ Fen Bilimleri Enstitüsü, Trabzon.

- Yüksek, H., Bahceci, S., Ocak, Z., Alkan, M., Ermis, B., Mutlu, T., Ocak, M., & Ozdemir, M. (2004). Synthesis of some 4,5-dihydro-1H-1,2,4-triazol-5-ones, *Indian journal of heterocyclic chemistry*, 13 (4), 369-372.
- Yüksek, H., Üçüncü, O., Alkan, M., Ocak, Z., Bahçeci, Ş., & Özdemir, M. (2005). Synthesis and Determination of pKa Values of Some New 3,4-Disubstituted-4,5-Dihydro-1H-1,2,4-triazol-5-one Derivatives in Non-aqueous Solvents, *Molecules* 10, 961-970.
- Yüksek, H., Küçük, M., Alkan, M., Bahçeci, S., Kolaylı, S., Ocak, Z., Ocak, U., Şahinbas, E., & Ocak, M. (2006). Synthesis and antioxidant activities of some new 4-(4-hydroxybenzylidenamino)-4,5-dihydro-1H-1,2,4-triazol-5-one derivatives with their acidic properties, *Asian Journal of Chemistry*, 18 (1), 539-550.
- Tereci, H., Askeroğlu, İ., Akdemir, N., Uçar, İ., & Büyükgüngör, O. (2012). Combined experimental and theoretical approaches to the molecular structure of 4-(1-formylnaphthalen-2-yloxy)phthalonitrile, *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 96, 569-577.
- Turhan Irak, Z., & Gümüş, S. (2017). Heterotricyclic compounds via click reaction: A computational study. *Noble International Journal of Scientific Research*, 1(7), 80-89.
- Vosko, S. H., Wilk, L. :& Nusair, M. (1980). Accurate spin-dependent electron liquid correlation energies for local spin density calculations: a critical analysis, *Canadian Journal of Physics*. 58, 1200.
- Wade, Jr. L.G. (2006). Organic Chemistry, 6nd ed.; Pearson Prentice Hall: New Jersey.
- Wolinski, K., Hinton, J. F. & Pulay, P. (1990). Efficient implementation of the gauge-independent atomic orbital method for NMR chemical shift calculations. *Journal of the American Chemical Society*, 112, 8251-8260.

Author Information

Murat Beytur

Kafkas University
Faculty of Science and Letters, Department of Chemistry,
Kars, Turkey
Contact E-mail: muratbeytur83@gmail.com

Haydar Yuksek

Kafkas University
Faculty of Science and Letters, Department of Chemistry,
Kars, Turkey
