PAPER DETAILS

TITLE: Investigation of Vitamin D Receptor Gene rs731236 Polymorphism in Turkish Rugby Players AUTHORS: Basak Funda EKEN,Zeyneb Nur SAHIN,Kate LLEWELLYNWATERS,Gökhan TUNA,Ajlan SAÇ,Sehkar OKTAY,Korkut ULUCAN PAGES: 33-40

ORIGINAL PDF URL: https://dergipark.org.tr/tr/download/article-file/3137690

ARAŞTIRMA MAKALESİ / RESEARCH ARTICLE

Investigation of Vitamin D Receptor Gene rs731236 Polymorphism in Turkish Rugby Players

Türk Ragbi Oyuncularında D Vitamini Reseptör Geni rs731236 Polimorfizminin Araştırılması

> Başak Funda EKEN^{*} Zeyneb Nur ŞAHIN^{**} Kate Llewellyn-WATERS^{***} Gökhan TUNA^{****} Ajlan SAÇ^{*****} N. Şehkâr OKTAY^{******} Korkut ULUCAN^{********}

Abstract

Vitamin D is an important molecule responsible for the development, contraction and regeneration of muscles, and activates its biological functions by means of its receptors, the vitamin D receptor (VDR), which is a member of the nuclear receptor group. In the present study, we aimed to analyze the rs731236 distribution in Turkish rugby players. A total of 22 male players and 30 sedentary individuals (as a control group) participated in the study. Following DNA isolation, rs731236 polymorphisms of VDR was determined by real-time polymerase chain reaction (RT – PCR). Statistical analysis was carried out by chi-square test. CC, CT and TT genotype and percentage distributions for rs731236 polymorphism in players

Makale Gönderim Tarihi: 12.05.2023 Yayına Kabul Tarihi: 28.05.2023

^{*} PhD, Marmara University, Faculty of Dentistry, Medical Biology and Genetics, Istanbul, Turkey, basak.funda@marun.edu.tr, ORCID: 0000-0002-2690-8071

^{**} PhD, Marmara University, Faculty of Dentistry, Medical Biology and Genetics, Istanbul, Turkey, zeynebnursahin@gmail.com, ORCID: 0000-0002-1956-0206

^{***} MSc, Lansdown Clinic, Bath, UK, katellewellynwaters@icloud.com, ORCID: 0000-0002-9340-106X

^{****} Ass. Dr. Trakya University, Kırkpınar Physical Education and School, Edirne, Turkey, tunagokhan@yandex.com, ORCID: 0000-0002-3120-5602

^{*****} Assoc. Prof., Trakya University, Kırkpınar Physical Education and School, Edirne, Turkey, ajlansac@trakya.edu.tr, ORCID: 0000-0002-8645-8826

^{******} Assoc. Prof., Marmara University, Faculty of Dentistry Department of Biochemistry, Istanbul, Turkey, nsehkar@yahoo.com, ORCID: 0000-0002-2878-288X

^{*******} Prof. Dr., Marmara University, Faculty of Dentistry, Medical Biology and Genetics, Istanbul, Turkey, korkutulucan@hotmail.com, ORCID: 0000-0002-1304-9386

were as 5 (23%), 10 (45%) and 7 (32%), respectively. Allelic distributions were found to be 20 (45%) for C allele, and 24 (55%) for T allele. For the control group; CC, CT and TT genotypes and percentages were as 4 (13%), 17 (57%) and 9 (30%), respectively. C allele was counted as 25 (42%) and 35 (58%) for controls. When we compare athletes and control groups in the terms of genotypes, we detected no statistically significant difference (p=0.12). For the allele comparison, we detected no significant difference between groups (p=0.76). Determination of gene combinations in athletes is of great importance for carrying out of the optimal training programs. Therefore, we aimed to determine the distribution of the alleles related with athletic performance on successful athletes. In our cohort, the CT genotype and T allele were more intensely determined in players.

Keywords: VDR, polymorphism, athletes, rugby, athletic performance

Öz

D vitamini kasların gelişmesinden, kasılmasından ve yenilenmesinden sorumlu önemli bir moleküldür ve nükleer reseptör grubunun bir üyesi olan vitamin D reseptörü (VDR) reseptörleri aracılığıyla biyolojik fonksiyonlarını aktive eder. Bu çalışmada, Türk rugby oyuncularında rs731236 dağılımını analiz etmeyi amaçladık. Çalışmaya toplam 22 erkek oyuncu ve 30 sedanter birey (kontrol grubu olarak) katıldı. DNA izolasyonunu takiben, VDR'nin rs731236 polimorfizmleri gerçek zamanlı polimeraz zincir reaksiyonu (Rt-PCR) ile belirlendi. İstatistiksel analiz ki-kare testi ile yapıldı. Sporcularda rs731236 polimorfizmi için CC, CT ve TT genotip ve yüzde dağılımları sırasıyla 5 (%23), 10 (%45) ve 7 (%32) olarak bulundu. Alel dağılımları C aleli için 20 (%45) ve T aleli için 24 (%55) olarak bulundu. Kontrol grubu için; CC, CT ve TT genotipleri ve yüzdeleri sırasıyla 4 (%13), 17 (%57) ve 9 (%30)'dur. C aleli kontrol grubu için 25 (%42) ve 35 (%58) olarak belirlendi. Sporcular ve kontrol grubunu genotip açısından karşılaştırdığımızda istatistiksel olarak anlamlı bir fark tespit edilmedi (p= 0,12). Alel karşılaştırması için gruplar arasında anlamlı fark tespit edilmedi (p= 0,12). Alel karşılaştırması için gruplar arasında anlamlı fark tespit edilmedi için büyük önem taşımaktadır. Bu nedenle atletik performansla ilgili alellerin başarılı sporcular üzerindeki dağılımını belirleneyi amaçladık. Bizim kohortumuzda CT genotipi ve T aleli oyuncularda daha yoğun olarak belirlendi.

Anahtar Kelimeler: VDR, polimorfizm, sporcu, ragbi, atletik performans

INTRODUCTION

Sport is a physical and cognitive contest in which multiple factors effect. In both individual or team sports, individuals' athletic performance is very important and the limit of the player's ability/ capacity is influenced by several factors, including physical, physiological, environmental and mental factors (Ulucan, 2016; Eken et al., 2021). All factors are directly or indirectly related with genetic endowment (Wang et al., 2018). This why the intrigue of association between sports and genes or continues to fascinate both athletes and researchers, despite current findings.

In the terms of athletic performance, muscle metabolism is expected to be optimal. Rugby, unlike other sports, is a team sport that significantly needs to be well-structured and according to the positions in terms of anthropometric diversity and physiological differences of players (Heffernan et al., 2015). Therefore, analyzing athletes' performance and defining physical boundaries is an important step to uncovered a successful game. In the simplest term, while Props (tight and loosehead, number 1 & 2) need lifting strength and endurance to scrum, Lock players (number 4 & 5) key characteristic is height and mobility. On the other hand, while the expectation from Flankers (number 6 & 7) is no fear and handling ability along with mental strength and fearlessness, Fly half (Number 10) are

required to have precise and accurate kicking ability, with excellent communication skills. When we look at the marker-assisted strategy studies done specifically on rugby athletes, there are some studies that have been conducted with the genes, which are significant in the pool of sports-related genes.

Vitamin D is a fat-soluble sterol molecule that plays a direct or indirect role in musculoskeletal development such as muscle growth (Dawson-Hughes et al., 2017) and bone mineralization (Orr, Holt, Wilkins, & Boone, 1923). In particular, the active form of vitamin D (25OHD3), which is 1,25(OH)2D3 and also known as calcitriol, which acts as a key regulator in many systems such as immune response (Murdaca et al.,2019), cardiovascular events (Wang, Manson, Song, & Sesso 2010) apoptosis and cancer mechanisms (Deeb, Trump, & Johnson, 2017) as well as the central nervous system (Häusler & Weber, 2019). In activates its biological functions by binding its receptor, vitamin D receptor (VDR). Any mutation on the pathways may affect the activity of gene regulation or change the entire system that can cause mortal diseases as previously mentioned (Kim et al., 2017). The vitamin D receptor is a member of the nuclear receptor group. The gene encoding the vitamin D receptor (VDR) is localized at 12q13.11. It is 100 kb long and has more than 100 polymorphic regions identified before (Norman et al., 2006). The rs731236 polymorphism in exon 9 of *VDR* is caused by T>C transition, due to the change ATT codon converts to ATC and encodes the isolate form (Ulucan et al., 2012).

The aim of this study is to investigate the distribution of *VDR* rs731236 polymorphism in rugby players. To date, there are no studies on Turkish rugby players in the terms of *VDR* rs731236 polymorphism. Therefore, it is aimed to determine the distribution of this gene allele in rugby players. According to the obtained data, it is aimed to be a pioneer for future studies and to contribute to the literature.

METHODS

Participants

A total of 22 male professional rugby players and 30 sedentary controls, all with Turkish ancestry, were recruited for the study.

Ethical Statement

Our study and study protocol were prepared in accordance with the Helsinki Declaration-2 (2015) guidelines and approved by Usküdar University Non-Interventional Ethics Committee (Protocol code: B.08.06,YÖK.2.ÜS.0.05.06/2013/09). The volunteers participating in the study were given detailed information about the analyzes and outputs before the study and their consent forms were obtained from them.

VDR Genotyping

DNA Isolation: Oral epithelium cells were collected by DNA collection swabs from the volunteers who participated in the study, and DNA isolation was completed by using PureLink DNA isolation kit (Invitrogen, Van Allen Way Carlsbad, CA, USA). Briefly, 20 μ L proteinase K was vortexed by adding 10 μ L of RNAase to 200 μ L of DNA isolation. After 2 min at room temperature, 200 μ L of binding buffer was added and homogenized with stirring. After incubation for 10 minutes in a 55 ° C water bath, 200 μ L of ethanol was added and vortexed for 5 seconds. It was taken to the filtered tube and centrifuged at 10000g for 1 minute. The supernatant was discarded and 500 μ L of washing buffer was added and centrifuged at 10000 g for 1.15 seconds. 80 μ L of elution buffer was added and incubated and centrifuged at maximum speed for 1 minute. The DNA samples obtained were stored at – 20 °C until the analysis of the respective gene regions was completed.

Genotyping of VDR rs731236

Genotyping of *VDR* rs731236 was performed from the isolated DNAs by using 7500 Fast Real-Time PCR System (Applied Biosystems). Taqman Genotyping Assays (Applied Biosystems Foster City, CA, USA) genotyping kit was used for allelic determination. C and T alleles were determined using VIC and FAM primers, respectively (Table 1). Genotyping was completed using 5 μ L master mix, 3.75 μ L H2O, 0.25 μ L assay and 1 μ L (10 ng) DNA.

Table 1. VIC/FAM labelled times primer use to rs731236 polymorphism

qPZR	qPZR Sequence (5^3
VIC/FAM	GGACGCCGRGCTGAT [C/T] GAGGCCATCCAGGAC

Statistical Analysis

All data were analyzed by using SPSS 20,0 for windows (SPSS Inc., Chicago, IL, USA). Statistical analysis was conducted by using chi-square test. Relationships yielding p – values less than 0.05 (p<0.05) were considered to be significant.

RESULTS

22 rugby athlete and 30 control groups participated in the study. Genotypes and alleles for the VDR rs731236 polymorphism are summarized in the table. CC, CT and TT genotypes and percentages for rs731236 polymorphism were analyzed as 5 (23%), 10 (45%) and 7 (32%); respectively. C allele was counted as 20 (45%) and T allele as 24 (55%). CC, CT and TT genotypes and percentages for rs731236 polymorphism in controls were as 4 (13%), 17 (57%) and 9 (30%); respectively. For allelic distributions, C allele was counted as 25 (42%) and T allele as 35 (58%). When we compare athletes and controls in the terms of genotypes, we detected no statistically significant difference (p=0.12).

For the allele comparison, we detected no significant difference between groups (p= 0.76). Table 2 summarizes the genotype, allele comparison of our cohort and their statistical analysis

	VDR Genotype			Allele Frequency			
	CC	СТ	ТТ	p-value	С	Т	p-value
Players (n=22)	5	10	7	0.12	20	24	0.76
Percentage	23%	45%	32%		45%	55%	
Controls(n=30)	4	17	9		25	35	
Percentage	13%	57%	30%		42%	58%	

Table 2. Comparison of the VDR rs731236 polymorphism of the study cohort

DISCUSSION and CONCLUSION

Rugby is a high-impact sport that is played with two teams of 15 players, one with two halves of 40 minutes each. Therefore, optimal athletic performance should be supplied for successful players. In the formation and development of athletic performance, muscle metabolism is expected to be optimal. The effect of genetic variations on metabolism on and cellular basis of activities, such as exercise, is important a lot through article crucial, therefore optimal to more favourable forms are expected to be included in the genotypes of athletes (Ulucan, Göle, Altindas, & Güney, 2013).

Recent studies demonstrated the significance of genetic parameters in athletic performance and physical fitness. In the studies carried out to date, more than 200 gene variants were associated with athletic performance, some related with structural and, some with functional factors like mediators or enzymes (Arica et al., 2018). The comparison works between environmental and genetic factors to determine the athletic performance and which have more effect on sports exercise metabolism has not been fully identified and warrants further research is a point of research (Bryık et al., 2018).

CT genotype and T allele were more dominant compared with other genotypes and C allele in terms of *VDR* rs731236 polymorphism in our cohort. According to the best of our knowledge, this report is the first which investigates reflects the genotype distribution of rs731236 polymorphism in Turkish rugby players.

Rabon-Stith et al. (2005) stated that rs2228570 polymorphisms influences bone mineral density of endurance training in healthy individuals. In the same study, they reported that they were not associated with aerobic training in a similar study. In another study, the effect of *Apa1* and *Bsm1* polymorphisms on muscle endurance was stated the same effect is not associated with *Taq1* polymorphism (Wang et al., 2006).

Bahat et al. (2010) reported that extensor strength of the knee was associated with rs1544410, but no significant association was found in rs2228570 and rs731236 haplotypes. In the study conducted with rs2228570 polymorphism on young Italian male footballers, they found the CC genotype as 52%, CT genotype as 34% and TT genotype as 14%, and demonstrated that the obtained genotype

results may be an important marker in football susceptibility (Micheli et al., 2011). Eken et al. 2018 analyzed *Fok1* (rs 2228570) and *Bsm1* (rs1544410) polymorphisms in 27 athletes and stated that CC genotype and C allele for *Fok1* polymorphism and AG genotype and A allele for *Bsm 1* were superior to other genotypes and alleles in the given cohort.

Studies investigating the role of rs731236 and athletic performance are not enough. Some previous studies in the literature reported that the VDR rs2228570 polymorphism increases the risk of stress fractures in athletes who exercise heavily, and the genetic variant in question should be analyzed to create healthy training programs. Varley et al. (2018) reported the important role of VDR rs10735810 and rs731236 in the pathophysiology of stress fractures in a study with 518 elite athletes. They stated that homozygotes of the rare allele of rs731236 polymorphism showed an association with increased stress fracture risk when compared to those homozygotes for the common allele combined with heterozygotes. This demonstrates the important functions of rs731236 in athletic performance, considering the role of this polymorphism in stress fracture condition. Another study on 385 Israeli soldiers, 17 genes, including VDR gene, were analyzed and it was determined that VDR could create the risk of stress fracture (Yanovich et al., 2012). Polymorphisms in different genes, including VDR, can impact muscle strength and alter responses to training stimuli. Lazary et al. (2013) conducted a study on 788 Hungarian school children and analyzed three functional SNPs (rs4516035, rs1544410 and rs731236) in their cohort. They constructed a haplogroup of SNPs (rs1544410-rs731236-rs10783215), and reported that haplogroups were significantly associated with increased grip strength of the dominant hand. As a result of their study, they concluded that VDR polymorphisms are significantly associated with different muscle-related phenotypes.

Genes are very important in determining athletic performance. As athletic performance involves multifactorial elements, analyzing the gene groups instead of single genes may provide precise results. In the rugby cohort, we detected no statistically significant difference in the terms of genotype and in the terms of alleles. The low numbers of the players and lacking the information of serum 25-hidroxy D vitamin levels may be considered as the major limitations of the present study. However, we believe that the effect of *VDR* gene on athletic performance should be determined with similar studies involving a greater number of subjects.

Conflict of interest: There is no conflict of interest between the authors regarding the publication of this article.

Authors' contributions: Design of the study: 1. Author % 40, 2. Author % 10, 3. Author % 10, 4. Author % 5, 5. Author % 5, 6. Author % 15, 7. Author 15 contributed. All authors have read and approved the final manuscript

Ethics Committee: Uskudar University Non-Interventional Ethics Committee, Date: 07.03.2013. Protocol number: B.08.06,YÖK.2.ÜS.0.05.06/2013/09

REFERENCES

- Arica, S., Solgun, HA, Dengi, A.S., Bakmaz, E., Akçay, T., Kapici, S., Sercan, C., Ulucan, K. (2018).–174 G/C polymorphism of interleukin 6 gene is not significantly different in Turkish professional short and long distance runners. *Cellular and Molecular Biology*, 64 (11),85-87
- Bahat, G., Saka, B., Erten, N., Ozbek, U., Coskunpinar, E., Yildiz, S., Sahinkaya, T.,& Karan M.A. (2010).BsmI polymorphism in the vitamin D receptor gene is associated with leg extensor muscle strength in elderly men. Aging Clinical and Experimental Research, 22(3), 198-205. doi: 10.1007/BF03324797.
- Biyik B, Kapici S, Sercan C, Kulaksiz H, Yüksel İ, & Ulucan K. (2018). Angiontensin Converting Enzyme Insertion
 / Deletion Polymorphism Profile Of Turkish Professional Hip-Hop And Latin Dancers. *Pamukkale Journal of Sport Sciences*, 20 9(3), 49-54. https://dergipark.org.tr/en/pub/psbd/issue/39562/335969
- Dawson-Hughes, B. (2017). Vitamin D and muscle function. The Journal of Steroid Biochemistry and Molecular Biology, 173, 313-316. doi: 10.1016/j.jsbmb.2017.03.018.
- Deeb, K.K., Trump, D.L., & Johnson, C.S. (2017). Vitamin D signalling pathways in cancer: potential for anticancer therapeutics. *Nature Reviews Cancer*, 7(9), 684-700. doi: 10.1038/nrc2196.
- Eken, B.F., Gezmiş, H., Sercan, C., Kapıcı, S., Chousein, Ö.M., Kıraç, D., & Ulucan K. (2018). Analysis of Vitamin D Receptor Fok1 (rs2228570) and Bsm1 (rs1544410) Polymorphism in Turkish Athletes. *IGUSABDER*, 6, 561-572. DOI: 10.38079/igusabder.453412
- Eken, B. F., Savci, A. A., Polat, T, Yılmaz, Ö. Ö., Şılar, Ç., Tacal Aslan, B., Oktay, Ş. & Ulucan, K. (2021). Preliminary Findings of the Distribution of Alpha-2-Adrenergic Receptor (ADRA2A) rs1800544 Polymorphisms in Kickboxing Players. *Clinical and Experimental Health Sciences*, 11 (4), 742-745. DOI: 10.33808/ clinexphealthsci.863591
- Häusler, D., &Weber MS. (2019). Vitamin D Supplementation in Central Nervous System Demyelinating Disease-Enough Is Enough. *International Journal of Molecular Sciences*, 20(1), 218. doi: 10.3390/ ijms20010218
- Heffernan, S.M, Kilduff, L.P., Day, S.H., Pitsiladis, Y.P., & Williams, A.G. (2015) Genomics in rugby union: A review and future prospects. *European Journal of Sport Science*, 15(6), 460-468. doi: 10.1080/17461.391.2015.1023222.
- Kim, D. (2017). The role of vitamin D in thyroid diseases. International Journal of Molecular Sciences, 18(9),1949. doi: 10.3390/ijms18091949
- Lazary, A., Bozsodi, A., Somhegyi, A., &Varga, PP. (2013).Vitamin D Receptor Polymorphisms Are Associated with Muscle Performance. *The Spine Journal*, 9,22-23. doi.org/10.1016/j.spinee.2013.07.085
- Micheli, M.L., Gulisano, M., Morucci, G., Punzi, T., Ruggiero, M., Castellini, E, & Pacini S. (2011). Angiotensinconverting enzyme vitamin D receptor gene polymorphisms and bioelectrical impedance analysis in predicting athletic performances of Italian young soccer players. *The Journal of Strength & Conditioning Research*, 25(8), 2084-2091. doi: 10.1519/JSC.0b013e31820238aa
- Murdaca, G., Tonacci, A., Negrini, S., Greco, M., Borro, M., Puppo F, &Gangemi S. (2019). Emerging role of vitamin D in autoimmune diseases: An update on evidence and therapeutic implications. *Autoimmunity Reviews*, 18(9), doi: 10.1016/j.autrev.2019.102350.
- Norman, A.W. (2006). Vitamin D receptor: new assignments for an already busy receptor. *Endocrinology*, 147, 5542-5548. doi: 10.1210/en.2006-0946.
- Orr, W.J., Holt, L.E., Wilkins, L., & Boone, FH. (1923). The calcium and phosphorus metabolism in rickets, with special reference to ultraviolet ray therapy. *The American Journal of Diseases of Children*, 26 (4), 362-372.
- Rabon-Stith, K.M., Hagberg, J.M., Phares, D.A., Kostek, M.C., Delmonico, M.J, Roth, S.M., Ferrell, R.E, Conway, J.M., Ryan, A.S., &Hurley B.F. (2005). Vitamin D receptor FokI genotype influences bone mineral

density response to strength training, but not aerobic training. *Experimental Physiol*ogy, 90(4), 653-661. doi: 10.1113/expphysiol.2005.030197.

- Ulucan, K. (2016). Literature review of Turkish sportsmen in terms of ACTN3 R577X polymorphism. *Clinical and Experimental Health Science*, 6, 44-47. doi: 10.5152/clinexphealthsci.2016.059
- Ulucan, K., Akyüz, S., Özbay, G., Pekiner, F.N., &Güney, A.İ. (2012). Evaluation of Vitamin D Receptor (VDR) Gene Polymorphisms (FokI, TaqI and ApaI) in a Family with Dentinogenesis Imperfecta. *Cytology and Genetics*, 47(5), 282 – 286
- Ulucan, K., Göle, S., Altindas, N., & Güney, AI. (2013).Preliminary findings of alpha-actinin-3 gene distribution in Turkish elite wind surfers. *Balkan Journal Medical Genetics*, 16 (1), 69 – 72. doi: 10.2478/bjmg-2013-0020.
- Varley, I., Hughes, D.C., Greeves. J.P, Stellingwerff, T., Ranson, C., Fraser, W.D., & Sale C. (2018). The association of novel polymorphisms with stress fracture injury in Elite Athletes: Further insights from the SFEA cohort., *Journal of Science and Medicine in Sport*,21(6), 564-568. doi: 10.1016/j.jsams.2017.10.038.
- Wang, L., Manson, J.E., Song, Y., & Sesso, HD. (2010). Systematic review: vitamin D and calcium supplementation in prevention of cardiovascular events. *Annals Of Internal Medicine*, 152(5),315-323. doi: 10.7326/0003-4819-152-5-201003.020.00010.
- Wang, T., Heianza, Y., Sun, D., Huang, T., Ma, W., Rimm, E.B., & Qi L. (2018). Improving adherence to healthy dietary patterns, genetic risk, and long term weight gain: gene-diet interaction analysis in two prospective cohort studies. *British Medical journal*, 360, 5644. doi: 10.113 6/bmj.j5644.
- Wang, P., Ma, L.H., Wang, H.Y., Zhang, W., Tian, Q., Cao, D.N., Zheng, G.X., &Sun Y.L. (2006). Association between polymorhisms of vitamin D receptor gene ApaI, BsmI and TaqI and muscular strength in young Chinese women. *International Journal of Sports Medicine*, 27(3), 182-186. doi: 10.1055/s-2005-865626.
- Yanovich, R., Friedman, E., Milgrom, R., Oberman, B., Freedman, L., & Moran, Ds. (2012). Candidate gene analysis in Israeli soldiers within stress fractures. *Journal of Sports Science and Medicine*, 11, 147-155.