## PAPER DETAILS

TITLE: Accumulation of Heavy Metals in Cladophora fracta and Chaetomorpha ligustica Species

AUTHORS: Öznur Yazilan Çamlik, Ergün Taskin

PAGES: 70-78

ORIGINAL PDF URL: https://dergipark.org.tr/tr/download/article-file/3907716



Mediterranean Fisheries and Aquaculture Research

https://dergipark.org.tr/medfar



**ORIGINAL RESEARCH PAPER** 

Received: 5 May 2024 | Accepted: 25 Jul. 2024

### Accumulation of Heavy Metals in Cladophora fracta and Chaetomorpha ligustica Species

# Öznur Yazılan Çamlık<sup>1</sup>\*<sup>1</sup>, Ergün Taşkın<sup>1</sup>

<sup>1</sup> Department of Biology, Faculty of Engineering and Natural Sciences, Manisa Celal Bayar University, Manisa, Türkiye

\*Corresponding author e-mail: oznuryazilan@gmail.com

### ABSTRACT

Besides playing a crucial ecological role, certain types of algae have also become significant indicators of chemical pollutants. In this research, marine algae were examined to evaluate their suitability as biomonitoring instruments for detecting heavy metal pollution in coastal regions. The levels of cadmium (Cd), nickel (Ni), copper (Cu), lead (Pb), and zinc (Zn) were assessed in two types of green algae: *Cladophora fracta* and *Chaetomorpha ligustica*, which were both gathered from the same natural habitat. Samples were collected from the coast of Bostanlı, Izmir (Aegean Sea, Turkey) in August 2023. The collected samples were first subjected to morphological species identification. Inductively Coupled Plasma-Mass Spectro-metry (ICP-MS) analysis was used to measure the accumulated concentrations of heavy metals (Zn Ni, Cu, Pb, Cd) of the separated species. As a result of the analyses, it was determined that the accumulation of heavy metals was higher in *Cladophora fracta* compared to *Chaetomorpha ligustica*. Both species have particularly high accumulation potentials for Zn in their tissues. The heavy metal accumulation concentrations for both species are as follows, respectively: Zn > Ni > Cu > Pb > Cd.

KEYWORDS: Heavy metal, Aegean Sea, pollution, green algae

How to cite this article: Yazılan Çamlık, Ö., Taşkın, E. (2024) Accumulation of Heavy Metals in *Cladophora Fracta* and *Chaetomorpha Ligustica* Species. *MedFAR.*, 7(2):70-78. https://doi.org/10.63039/medfar.1478935.

### 1. Introduction

Heavy metal pollution is one of the most significant environmental issues of our time. Various industries such as mining, energy and fuel production, fertilizer and pesticide production and use, iron and steel industry, electro-osmosis, electrolysis, leather industry, electronics manufacturing, aviation, and atomic energy installation produce and discharge wastes containing various heavy metals into the environment. Due to all these activities, severe environmental pollution erupts, which ultimately threatens human health and the ecosystem. There are three types of heavy metals that are a cause for concern: toxic metals (Hg, Cr, Pb, Zn, Cu, Ni, Cd, As, Co, Sn, etc.), precious metals (Pd, Pt, Ag, Au, Ru, etc.), and radionuclides (U, Th, Ra, Am, etc.) (Wang and Chen, 2006).

The increasing human population and coastal expansion contribute to the rise in anthropogenic pollution burden, posing a serious threat to marine and aquatic environments. Polluted water can degrade water quality, thus limiting the use of water bodies for various purposes. It is well known that heavy metal pollution in aquatic environments is a growing problem worldwide and has currently reached an alarming level. Levels of heavy metal pollution in marine ecosystems can be estimated through analysis of aquatic organisms (Morillo et al., 2005). While certain trace elements such as manganese, copper, and chromium are necessary for biological functions, they can become toxic when present in excess amounts. Conversely, elements like arsenic, cadmium, lead, and mercury can be highly toxic even at very low concentrations. (Domingo, 1994).

From both economic and ecological standpoints, there is growing interest in utilizing plants as phytoremediators to purify water in diverse aquatic ecosystems (Eid et al., 2020; Ahmad et al., 2014; Lytle and Lytle, 2001). Many aquatic plant species have demonstrated the capability to uptake zinc and other heavy metals from water, thereby accumulating higher concentrations

internally compared to their surrounding environment (Whitton, Various 1981). species are commonly utilized to monitor concentrations different metal in geographical areas, often selected based on their heavy taxonomic distribution (Ho, 1990; Stratis et al. 1996). Metals of interest, such as copper, zinc, cadmium, and lead, have been highlighted (West, 1973). Organisms investigated as potential monitors for trace metals in water include clams (Lord, 1974), as well as clams, mussels, and shrimp (Bertine and Goldberg, 1972), groupers (Taylor and Bright, 1973), and algae (Stokes et al., 1973). It is true that algae primarily bind free metal ions, and the concentrations of these ions are influenced by the nature of suspended particulate matter. This matter is formed by a combination of organic and inorganic complexes, as noted by Luoma (1983), Seeliger and Edwards (1977), and Volterra and Conti (2000).

within the Chlorophyta Macroalgae phylum, an ecologically significant group in aquatic systems, are commonly utilized as indicators of heavy metal pollution, as noted McCormick and Cairns by (1994). Cladophora species, as a filamentous green algae, have been proposed as effective biological monitors for species present in natural waters (Whitton, 1970). It is indeed a hardy plant, thriving in diverse climatic regions, and it has been observed that substances are concentrated in the tissues of this plant to a significant extent (Keeney et al., 1976).

The uptake and bioaccumulation of heavy metals by algae have primarily been investigated using atomic absorption spectrophotometry (Stokes et al.1973; Stokes, 1975).

The aim of this study is to present the obtained data and compare them with the results found in the literature. Additionally, by comparing the heavy metal accumulation potentials of two species belonging to the genera *Cladophora* and *Chaetomorpha*, both of which are pollution indicators within the Cladophorales order, their potential applications and futures have been estimated.

These species have the ability to easily adapt to areas contaminated with heavy metals. In this study, heavy metal levels of toxic metals such as Zn, Pb, Ni, Cu, Cd were measured in *Cladophora* and *Chaetomorpha* species. The samples were collected from Bostanlı, İzmir, a region with high terrestrial pressure. The main reasons for selecting these species include their tendency to reach alarming levels due to cultural eutrophication, their wide distribution ranges, and their high biological accumulation capacities attributed to their adaptation to various climatic regions.

### 2. Material and Methods

### 2.1. Materials

The materials of this study consist of species belonging to the genera *Cladophora* and *Chaetomorpha* within the order Cladophorales.

### Cladophora fracta (O.F.Müller ex Vahl) Kützing

The apical cell of the algae has a width of  $30 \ \mu\text{m}$  and a length of  $125 \ \mu\text{m}$ . Median cells range in width from  $50 \ \mu\text{m}$  to  $100 \ \mu\text{m}$  and in length from  $350 \ \mu\text{m}$  to  $250 \ \mu\text{m}$ . Apical cells have oval tips, and the branching pattern is pseudodichotomous. The thallus is slender and filamentous, with a curved shape, and apical branching is prevalent. Various cell shapes were observed, with round cells being predominant. The cells contain numerous chloroplasts arranged reticularly (Figure 1).

### Chaetomorpha ligustica (Kützing) Kützing

This species forms a soft and intertwined, bright to light green mass composed of fine filaments, which consist of a series of nonbranching cells (Figure 2). The cells are 90  $\mu$ m in diameter, multinucleate, and the length is approximately 4 times the width.



Figure 1. Cladophora fracta. a) Habit, b) Filament and cell structure



Figure 2. Chaetomorpha ligustica. a) Habit, b) Filament and cell structure

### 2.2. Study Area and Sampling

Samples were collected from the coast of Bostanlı, Izmir (38.456731°N, 27.090075°E) (Aegean Sea, Turkey) in August 2023 (Figure 3). The sampling area within İzmir Bay is influenced by the heavy ship traffic in İzmir Port, the discharge of treated water into the bay, and the water carried by the Gediz River.

The samples were directly collected by hand from the sublittoral zone. The collected samples were washed with clean water to separate them from waste and epiphytes. The washed samples were then brought to the laboratory. Samples were examined at the Department of Biology, Manisa Celal Bayar University (Türkiye). Cladophorales species initially were examined under а stereomicroscope and systematically categorized. Subsequently, a more detailed examination was conducted under а microscope, and the apical cell length, width, thallus cell length, branching patterns, and cell shapes of each species were measured. The identification of the samples was made on the basis of the descriptions by Cormaci et al. (2014) and Sfriso (2010).

The species identified were first dried in the sun, and then 0.5 grams of each species were weighed for heavy metal analysis. Approximately 0.5g of homogenized samples were weighed precisely and digested using a mixture of nitric acid, hydrochloric acid, hydrogen peroxide, and hydrofluoric acid in a 3000 Multiwave microwave oven under an appropriate temperature/pressure program. The amounts of Zn, Ni, Cu, Cd, and Pb in the solutions prepared using the microwave digestion system were measured with ICP-MS. The values presented for the samples represent the results of a single measurement.



Figure 3. General view of the research area

#### 3. Results

We compared the heavy metal accumulation potentials of two different species belonging to two different genera, both of which serve as pollution indicators, collected from the same habitat. The levels of cadmium (Cd), copper (Cu), lead (Pb), nickel (Ni), and zinc (Zn) in the collected samples were investigated, and the obtained results were evaluated. Based on the ICP-MS analysis, we found that both species accumulated the highest levels of Zn.

Heavy metal accumulation concentrations in the thallus of *Chaetomorpha ligustica*; Zn: 197,496  $\mu$ g/g Ni: 36,694  $\mu$ g/g Cu: 12,452  $\mu$ g/g Pb:11,514  $\mu$ g/g Cd:0,097  $\mu$ g/g (Figure 4).

The order of accumulation potentials and concentrations of other metals was as follows: Zn > Ni > Cu > Pb > Cd.

Heavy metal accumulation concentrations in the thallus of *Cladophora fracta*; Zn: 117,153  $\mu$ g/g Ni: 80,467  $\mu$ g/g Cu: 25,054  $\mu$ g/g Pb:20,076  $\mu$ g/g Cd:0,613  $\mu$ g/g (Figure 4).

It has been determined that, generally, *Cladophora fracta* has a higher heavy metal accumulation potential than *Chaetomorpha ligustica*, except for zinc.



Figure 4. Heavy metals concentrations in *Cladophora fracta* and *Chaetomorpha ligustica* samples

#### 4. Discussion

Chemical pollution in aquatic ecosystems can cause destructive effects. Heavy metals, which are transferred through the food chain, not only contribute to the degradation of aquatic ecosystems but also have adverse effects on human health. It is known that utilizing aquatic organisms for remediation and monitoring of pollution in aquatic ecosystems is possible. Particularly, due to immobility compared their to other organisms, wide distribution ranges, and ease of collection, aquatic plants are more suitable for use in biological remediation and monitoring studies.

In recent years, many studies have been published using different macroalgae as biomonitors and for water remediation purposes (Agarwal et al., 2020).

For many years, algae have been recognized for their role in wastewater treatment and their ability to bind heavy metal cations. This is primarily due to their high negative surface charge, which allows them to attract and absorb positively charged heavy metal ions from water. This property makes algae effective in removing pollutants like heavy metals from wastewater, contributing significantly to water purification processes (Rao, 1986).

The concentrations of heavy metals accumulated in macroalgae vary depending on the geological structure of the sampling area, the season of sampling, morphological characteristics of the algae, their lifespan, physicochemical parameters of their habitat, and interactions with other heavy metals (Sawidis et al., 2001).

Previous studies conducted in the Aegean Sea have reported Pb values for *Cystoseira* species ranging between 0.02-2.5  $\mu$ g g<sup>-1</sup> (Sawidis et al., 2001), while Akçalı and Küçüksezgin (2009) reported a value of 0.083  $\mu$ g g<sup>-1</sup>. When these values are compared with the results obtained in this study, it is evident that the Pb value found in this study is considerably higher.

Wang and Chen (2009) reported that the capacity of algae to absorb heavy metals is quite high. Vymazal (1987) demonstrated in a study that *Cladophora* species have a rapid ability to absorb Zn.

In benthic macrophytes, Zn concentrations below 100  $\mu$ g g<sup>-1</sup> are considered to indicate unpolluted areas, as evaluated by Storelli et al. (2001), while higher concentrations are indicative of anthropogenic pollution. The highest Zn value obtained in this study is 197.496  $\mu$ g/g, indicating a high anthropogenic impact.

Malea and Haritonidis (1999) stated that algae accumulate essential metals such as Zn at high levels while maintaining toxic metals like Cd, Cu, and Pb at low levels necessary for metabolic activities.

In a study conducted in 2011 to measure the levels of some heavy metals in *Cladophora glomerata* seasonally in Lake Hazar, the Cu concentration was found to be 1.31 mg/g and 1.77 mg/g, and Zn was found to be 9.61 mg/g and 11.8 mg/g (Alp and Ozbay, 2011).

In a study conducted by Akçalı and Küçüksezgin (2009) along the Aegean Sea coast, the order of heavy metal concentrations in *Cystoseira* species was determined to be Pb < Cd < Cu < Zn. In this study, the highest Zn accumulation value for

Cystoseira species was reported to be 62.48 µg g<sup>-1</sup>. According to Akçalı and Küçüksezgin (2011). the highest heavy metal concentration measurements in macroalgae species Enterophmorpha sp., Gracilaria gracilis, and Ulva sp. collected from the shores of Bostanlı (Izmir) were reported as follows: Cd (0.49 µg/l), Pb (3.20 µg/l), Cu (5.57  $\mu$ g/l), and Zn (10.36  $\mu$ g/l). Cetingül et al. (2000) reported the heavy metal accumulation concentrations of Cladophora dalmatica species as Cu 5.93 µg g<sup>-1</sup> and Zn 136.61  $\mu$ g g<sup>-1</sup> in their study conducted in Izmir Bay. Sawidis et al. (2001), their study on Cystoseria sp. in the Aegean Sea, reported Pb values ranging from 0.02 to 2.5  $\mu$ g g<sup>-1</sup>, whereas the Pb value obtained in this study is significantly higher.

In this study. the heavy metal concentration levels of Chaetomorpha ligustica species are as follows: Zn: 197.496 μg/g, Ni: 36.694 μg/g, Cu: 12.452 μg/g, Pb: 11.514 µg/g, Cd: 0.097 µg/g. The heavy metal concentrations in Cladophora fracta species are as follows: Zn: 117.153 µg/g, Ni: 80.467 µg/g, Cu: 25.054 µg/g, Pb: 20.076  $\mu g/g$ , Cd: 0.613  $\mu g/g$ . When looking at the values from previous studies, it is seen that the special Zn, Pb and Cu values obtained in this study are higher.

Considering the conducted studies, İzmir Bay has significantly high concentrations of heavy metals due to industrial activities, İzmir port, and the inflow of rivers into the bay.

It is understood that *Cladophora* and *Chaetomorpha*, which adapt to the high anthropogenic influence in this region, have high biological accumulation potentials. Considering their wide distribution ranges, ability to thrive in different climatic regions, and dominance in aquatic environments with high chemical pollution, it is thought that these species are more suitable to be used as biomonitoring organisms compared to other species.

### 5. Conclusion

Rincon et al. (2005) indicated in their study on *Cladophora* species that they have the potential to be used as biological material for remediation purposes, particularly in the removal of heavy metal pollution. In addition to being used as a biomonitoring, their high biological accumulation potentials make them suitable for use in biological purification purposes as well. Brinza et al. examined metal (2007)the heavy accumulation capacities of Cladophora species and reported that they have the ability to adsorb a high level of various heavy metal ions, including K, Mg, Ca, Fe, Sr, Co, Cu, Mn, Ni, V, Zn, As, Cd, Mo, Pb, Se, and Al.

Chmielewská and Medved (2001)confirmed the high bioaccumulation capabilities of Cladophora sp. for Pb, Cd, Ni, Cr, and V metals in their study aimed at accumulation determining their metal abilities. As a result, it was found that algae contribute to improved water quality through wastewater treatments.

As a result, it has been determined that among the species we have been studying, *Cladophora fracta*, in particular, has a high capacity for accumulating heavy metals. Utilizing the bioaccumulation abilities of macroalgae, it is possible to observe the extent of pollution in aquatic ecosystems resulting from human activities and take parallel measures accordingly.

Due to their tolerance to metals, simple morphology, provision of sufficient tissue for analysis, and sedentary lifestyle, *Cladophora* species can be successfully used in biomonitoring studies. Moreover, their ease of collection and relatively widespread distribution across various coastal regions ensure reliable outcomes (Zbikowski et al., 2007).

### **Compliance with Ethical Standards**

### **Conflict of interest**

The authors declare that they have no competing interests.

### Author contribution

All authors' contributions are equal for the preparation of research in the manuscript. All the authors verify that the Text, Figures, and Tables are original and that they have not been published before.

### **Ethical approval**

Ethics committee approval is not required.

### Acknowledgements

This study is supported by TÜBİTAK (The Scientific and Technological Research Council of Türkiye) under grant number 121Y215.

### References

Agarwal, A., Mhatre, A., Pandit, R., Lali, A.M. (2020) Synergistic biorefinery of Scenedesmus obliquus and Ulva lactuca in poultry manure towards sustainable bioproduct generation. Bioresource Technology 297: 122462.

Ahmad, S.S., Reshi, Z.A., Shah, M.A., Rashid, I., Ara, R., Andrabi, S.M.A. (2014) Phytoremediation Potential of Phragmites australis in Hokersar Wetland-a Ramsar Site of Kashmir Himalaya. Int. J. Phytoremediat 16: 1183–1191.

Akçalı, İ., Küçüksezgin, F. (2009) Ağır Metal Birikimi Ege Denizi Kıyılarında Görülen Kahverengi Alg Cystoseira sp.'de. E.Ü. Su Ürünleri Dergisi 3: 159-163. Akçalı, İ., Küçüksezgin, F. (2011) A biomonitoring study: Heavy metals in macroalgae from eastern Aegean coastal areas. Marine Pollution Bulletin 62(3): 637-645.

Alp, M.T., Özbay, Ö. (2011) Seasonal Levels of Certain Heavy Metals in Cladophora glomerata in Lake Hazar. Ekoloji 20(78): 13-17.

Bertine, K.K., Goidberg, E.D. (1972) Trace elements in clams, mussels and shrimp. Limnology Oceanogrophy 17: 877-84.

Brinza, L., Dring, M.J., Gavrilescu, M. (2007) Marine micro and macro algal species as biosorbents for heavy metals. Environmental Engineering and Management Journal 6: 237–251.

Chmielewská, E., Medved, J. (2001) Bioaccumulation of Heavy Metals by Green Algae Cladophora glomerata in a Refinery Sewage Lagoon. Croatica Chemica Acta 74(1): 135-145.

Cormaci, M., Furnari, G., Alongi, G. (2014) Flora marina bentonica del Mediterraneo: Chlorophyta. Bollettino dell'Accademia Gioenia di Scienze Naturali di Catania 47: 11-436.

Çetingül, V., Aysel V., Kurumlu-Kuran, Y. (2000) Biochemical Investigation and Heavy Metal Contents of Cladophora dalmatica Kütz. and Ceramium ciliatum (Ellis) Duel. var. robustum (J.Ag.) from Aegean Sea (Turkish Coast) Turkish Journal Marine Sciences 6(1): 9-22.

Domingo, J.L. (1994) Metal-induced developmental toxicity in mammals: a review. Journal of Toxicology Environmental Health 42:123–141.

Eid, E.M., Galal, T.M., Sewelam, N.A., Talha, N.I., Abdallah, S.M. (2020) Phytoremediation of Heavy Metals by Four Aquatic Macrophytes and Their Potential Use as Contamination Indicators: A Comparative Assessment. Environmental Science Pollution Research 27: 12138– 12151.

Ho, Y. (1990) Ulva lactuca as bioindicator of metal contamination in intertidal waters in Hong Kong. Hydrobiology 203: 73-81. Keeney, W. L., Breck, W. G., Vanloon, G. W. Page, J. A. (1976) The determination of trace metals in Cladophora glomerata C. Glomerata as a potential biological monitor. Water Research 10(11): 981-984.

Lord, D.A. (1974) Trace elements in mussels and seston in the Kingston Basin of Lake Ontario, Ph.D. Thesis, Kington, Canada.

Lytle, J.S., Lytle, T.F. (2001) Use of Plants for Toxicity Assessment of Estuarine Ecosystems. Environmental Toxicology Chemistry 20: 68–83.

McCormick, P.V., Cairns, J.J., (1994) Algae as indicators of environmental change. Journal of Applied Phycology 6(5-6): 509-526.

Messyasz, B., Leska, B., Fabrowska, J., Pikosz, M., Roj, E., Cieslak, A., Schroeder, G. (2015) Biomass of freshwater Cladophora as a raw material for agriculture and the cosmetic industry. Open Chem 13: 1108– 1118.

Morillo, J., Usero, J., Gracia, I. (2005) Biomonitoring of Trace Metals in a Mine-Polluted Estuarine System (Spain). Chemosphere 58: 1421-1430.

Rao, S.V.R. (1986) Cadmium accumulation in fiddler crabs Uca annulipes: Uptake of lead, chromium, cadmium, and cobalt by Cladophora glomerata. International Journal of Environmental Studies 27: 219-223.

Rincon, J., Gonzalez, F., Ballester, A., Blazquez, M.L., Munoz, J.A. (2005) Biosorption of heavy metals by chemicallyactivated alga Fucus vesiculosus. Journal of Chemical Technology and Biotechnology 80: 1403-1407.

Sawidis, T., Brown, M.T., Zachariadis, G., Sratis, I. (2001) Trace metal concentrations in marine algae from different biotopes in the Aegean Sea. Environment International 27: 43-47.

Seeliger, U., Edwards, P. (1977) Correlation coefficients and concentration factors of copper and lead in seawater and benthic algae. Marine Pollution Bulletin 8 (1): 16–19. Sfriso, A. (2010) Chlorophyta multicellulari e fanerogame acquatiche, Arpa Emilia-Romagna, Bologna.

Stokes P.M., Hutchinson T.C., Krauter, K. (1973) Heavy-metal tolerance in algae isolated from contaminated lakes near Sudbury, Ontario. Canadian Journal of Botany 51: 2155-2168.

Stokes, P.M. (1975) Uptake and accumulation of copper and nickel by metaltolerant strains of scenedesmus. Verhandlungen International Vereinigung Limnologie 19: 2128-2137.

Storelli, M.M., Storelli, A., Marcotrigiano, G.O. (2001) Heavy metals in the aquatic environment of the Southern Adriatic Sea, Italy: Macroalgae, sediments and benthic species. Environment International 26: 505-509.

Stratis, I., Simeonov, V., Zachariadis, G., Sawidis, T., Madjukov, P., Tsakovski, S. (1996) Chemometrical approaches to treat analytical data from aquatic macrophytes and marine algae. Fresenius Journal Analytical Chemistry 355: 65-70.

Taylor, D.D., Bright, T.J. (1973) The Distribution of Heavy Metals in Reef-Dwelling Groupers in the Gulf of Mexico and Bahamas Islands. Department of Marine Resources, Information Center for Marine Resources. Texas A&M Univ. Tech. Rep. TAMU-SG:73-208.

Volterra, L., Conti, M.E. (2000) Algae as biomarkers, bioaccumulators and toxin producers, in: Conti, M.E., Botre, F. (Eds.), The Control of Marine Pollution: Current Status and Future Trends; Int. J.Environ. Pollut. Inderscience Enterprises Ltd., Milton Keynes, UK, pp. 92–125,

Vymazal, J. (1987) Zn uptake by Cladophora glomerata. Hydrobiologia 148: 97-101.

Wang, J.L., Chen, C. (2009) Biosorbents for heavy metals removal and their future. Biotechnology Advances 27: 195-226.

Wang, J.L., Chen, C. (2006) Biosorption of heavy metals by Saccharomyces cerevisiae: a review. Biotechnol Advances 24: 427–51. West P.W. (1973) Analytical studies of the environment. Pure Applied Chemistry 34: 163-170.

Whitton, B.A., Say, P.J., Wehr, J.D. (1981) Use of plants to monitor heavy metals in rivers, in: P. J. Say and B. A. Whitton (Eds.), Heavy metals in northern England: Environmental and biological aspects, Durham, pp. 135–145.

Whitton, B.A. (1970) Biology of Cladophora in freshwaters. Water Reserach 4: 457-476.

Zbikowski, R., Szefer, P., Latala, A. (2007) Comparison of Green Algae Cladophora sp. and Enteromorpha sp. as Potential Biomonitors of Chemical Elements in the Southern Baltic. Science of the Total Environment 387: 320-332.