

PAPER DETAILS

TITLE: Compulsive Water Drinking Resulting in Hyponatremia: A Pimozide Case

AUTHORS: Mehmet Hamdi ORUM,Oguzhan Bekir EGILMEZ

PAGES: 48-49

ORIGINAL PDF URL: <https://dergipark.org.tr/tr/download/article-file/876783>



Compulsive Water Drinking Resulting in Hyponatremia: A Pimozide Case

Hiponatremi ile Sonuçlanan Kompulsif Su İçme: Bir Pimozid Olgusu

Mehmet Hamdi Orum¹, Oguzhan Bekir Egilmez²

¹Kahta State Hospital, Clinic of Psychiatry, Malatya, Turkey

²Department of Psychiatry, Adiyaman University, Faculty of Medicine, Adiyaman, Turkey

Copyright © 2019 by authors and Medical Records Publishing Inc.

Abstract

Pimozide is an antipsychotic can cause side effects such as tremor, rigidity, dystonia, akathisia, gynecomastia. Another rare side effect is psychogenic polydipsia and subsequent hyponatremia. In this case report, we discussed the polydipsia-hyponatremia syndrome associated with pimozide, which is not currently preferred due to cardiac side effects. This case illustrates the risk of water intoxication in antipsychotic-treated patients with non-psychiatric care and emphasizes the interest of regular ionic control in such patients.

Keywords: Pimozide, hyponatremia, side effect, antipsychotic

Oz

Pimozid, tremor, rijidite, distoni, akatizi, jinekomasti gibi yan etkilere neden olabilen bir antipsikotiktir. Diğer bir nadir yan etkisi psikojenik polidipsiyi takiben ortaya çıkan hiponatremi tablosudur. Bu olgu sunumunda, günümüzde kardiyak yan etkileri nedeniyle fazla tercih edilmeyen pimozid ile ilişkili polidipsi-hiponatremi sendromunu tartıştık. Bu olgu, psikiyatrik bakımı yeterli olmayan ve antipsikotik kullanan hastalardaki su zehirlenmesi riskini göstermekte ve bu tür hastalarda düzenli iyon dengesi kontrolünün önemini vurgulamaktadır.

Anahtar Kelimeler: Pimozid, hiponatremi, yan etki, antipsikotik

INTRODUCTION

Pimozide is an antipsychotic diphenylbutyl-piperidine derivative used in the treatment of schizophrenia patients with blocking the dopamine 2 receptors in the nigrostriatal and mesolimbic dopamine pathways. It causes side effects such as tremor, rigidity, dystonia, akathisia, extrapyramidal symptoms, headache, galactorrhea, gynecomastia. Dose-dependent QTc prolongation, risks for ventricular arrhythmias, and sudden death syndrome are some serious side effects of pimozide. Edema, itching, skin rashes are rare side effects. Another rare side effect is psychogenic polydipsia and subsequent hyponatremia (1). In this case report, we discussed the polydipsia-hyponatremia syndrome associated with pimozide, which is not currently preferred due to cardiac side effects.

CASE PRESENTATION

A, 71-year-old male patient was followed up for more than 50 years with the diagnosis of schizophrenia according to Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5) (2). In the past, the drugs he used were haloperidol, chlorpromazine, quetiapine, risperidone, olanzapine, amisulpride, escitalopram, biperidene, alprazolam. She was admitted to the emergency department with vomiting, abdominal pain, dizziness, self-talk, and agitation. Laboratory findings were as follows: serum sodium (Na) level 117 mEq/L, serum chlorine (Cl) level 92 mEq/L, serum potassium (K) level 4.0 mEq/L, urinary osmolality 1300 mosmol/kg, urine sodium level 130 mEq/L. It was determined that 8 mg/day pimozide was used before the application. Drug-induced syndrome of

Geliş Tarihi / Received: 17.06.2019 Kabul Tarihi / Accepted: 08.08.2019

Sorumlu Yazar /Corresponding Author: Mehmet Hamdi Kahta State Hospital, Clinic of Psychiatric, Kahta, Turkey, E-mail: ftrmine@hotmail.com

inappropriate antidiuretic hormone (SIADH) was thought to develop. No medication was used, supportive treatment was given, and clinical symptoms improved after the third day. Serum Na 136 mEq/L was determined in his laboratory. Clozapine was started at 12.5 mg/day. Slowly titrated 200 mg/day dose was reached. White blood cell and electrolytes were in normal limits before discharge. Written informed consent was taken from the patient in order to publish her data. Naranjo Adverse Drug Reaction Probability Scale (NADPRS) score of the patient was 6 (3).

DISCUSSION

Hyponatremia due to drug use is a condition encountered in psychiatric practice. Various cases of polydipsia-hyponatremia due to antidepressants, mood stabilizers and antipsychotic use have been reported (4-6). Potomania, in the words of Koide (7), is either inherited in the psychiatric condition or induced by antipsychotic therapeutics. The author stated that this drug could cause a sensation of thirst and contribute to the development of a potomania, polydipsia-hyponatremia syndrome. Later, Nishimura et al. (8), Leclercq et al. (9) made notifications. There are not many studies about pimozone which is not used frequently because of its cardiac side effects. The mechanism invoked is not yet formally elucidated. Essentially, three hypotheses corresponding to three action sites are identified: action on the pituitary gland with increased synthesis and/or secretion of ADH; action on the kidney, either by direct action or by potentiating the activity of ADH at this level; central action on osmoreceptors with modifications of the threshold of activity leading to a secretion of ADH for an abnormally low level of blood osmolality. High urinary sodium (in our patient: 130 mEq/L) can be found in both etiologies and is certainly not a discriminating factor. The key element of the differential diagnosis appears to be urinary osmolality (1, 7-9). Ethically, it is not considered lawful to continue the administration of pimozone for fear of running the risk of a new episode of hyponatremia in a patient traumatized cranial.

As a result, we report a case of water intoxication with severe hyponatremia resulting in various symptoms in a patient diagnosed with schizophrenia recently treated by pimozone. This case illustrates the risk of water intoxication in antipsychotic-treated patients with non-psychiatric care and emphasizes the interest of regular ionic control in such patients.

NOT: This case report was presented as a poster in "11th International Congress on Psychopharmacology & 7th International Symposium on Child and Adolescent Psychopharmacology [Abstract:0020] [Psychopharmacology]"

Financial disclosures: All authors report no financial interests or potential conflicts of interest.

Conflict of Interest: The authors declare that they have no competing interest.

Acknowledgments: The funding entities had no role in the design of the study, the collection and analysis of data, the decision to publish, or preparation of the manuscript.

ORCID ID

Mehmet Hamdi Orum, ORCID: 0000-0002-4154-0738

Oguzhan Bekir Egilmez, ORCID: 0000-0002-6303-1758

REFERENCES

1. Smyj R, Wang XP, Han F. Pimozone. Profiles Drug Subst Excip Relat Methodol 2012;37:287-311.
2. American Psychiatric Association. (2013). Diagnostic and statistical manual of mental disorders (5th ed.). Washington, DC: Author.
3. Kose S, Akin E, Cetin M. Adverse drug reactions and causality: The Turkish version of Naranjo Adverse Drug Reactions Probability Scale. Psychiatry and Clinical Psychopharmacology 2017;27(2):205-6.
4. Orum MH, Kalenderoglu A, Egilmez OB, Ozen ME, Kapici Y. Hyponatremia associated with repeated use of sodium valproate. Psychiatry and Behavioral Sciences 2018;8(2):93-4.
5. Orum MH, Han-Almis B. Hyponatremia during treatment with the clozapine-amisulpride combination: A suspected association and improvement with dose reduction. Dusunen Adam The Journal of Psychiatry and Neurological Sciences Doi: 10.14744/DAJPNS.2019.00024.
6. Ozen ME, Orum MH, Kalenderoglu A. Escitalopram-induced hyponatremia: A case report. Psychiatry and Behavioral Sciences. Doi: 10.5455/PBS.20171230091731.
7. Koide H. Three cases of hyponatremia during administration of pimozone. No To Hattatsu 1991;23(5):502-5.
8. Nishimura H, Machida K, Nakano H, Shinozaki T, Kasahara H, Ushijima S, et al. A female case of schizophrenia who presented hyponatremia induced by pimozone. Sishin Igaku 1995;37(6):657-60. Doi: 10.11477/mf.1405903894.
9. Leclercq P, Canivet JL, Damas P, Lamy M. A case of severe hyponatremia under pimozone (Orap) administration. Rev Med Liege 1995;50(4):151-2.