# PAPER DETAILS

TITLE: Traditional medicinal plants used for the treatment of viral infections: A short review

AUTHORS: Ismail UZAR, Fatima UZAR

PAGES: 29-48

ORIGINAL PDF URL: https://dergipark.org.tr/tr/download/article-file/1096234

Yayıncı / Publisher Sağlık Bilimleri Üniversitesi University of Health Sciences

Derleme Makale / Review Article, 1(2): 29-48, 2020

# Traditional medicinal plants used for the treatment of viral infections: A short review

Fatıma Uzar<sup>1\*</sup> (D İsmail Uzar<sup>2</sup> (D

<sup>1</sup>University of Health Sciences, Institute of Health Sciences, Istanbul, Turkey <sup>2</sup> Bezmialem Foundation University, Faculty of Medicine, Istanbul, Turkey

\*Sorumlu Yazar / Corresponding Author: Fatima Uzar, e-mail: fatimauzar@gmail.com

#### ABSTRACT

Viruses are among the most infectious microorganisms and have been present alongside humans throughout history. Viruses infecting humans can be divided into DNA or RNA viruses. It is important to be familiar with the viral replication cycles and the infection mechanisms of each class of viruses sn order to produce efficient vaccines and develop effective treatment regimens, which will help mankind prevent or minimize pandemics such as the current COVID-19 pandemic. Many antiviral drugs are used as broad spectrum to attack common factors of viral infections. However, absence of equal accessibility worldwide and cost of treatments and vaccines, together with the undesirable side effects of these drugs encourages us to venture into alternative treatment modalities. This short review will focus on plants used in traditional medical systems worldwide and their corresponding pharmacological trials, both *in-vitro* and *in-vivo*. Medicinal plants used for Herpesviridae and Hepatitis B virus was addressed when considering DNA viruses, while plants used to treat HIV, influenza, and the Coronaviridae family was discussed under the scope of RNA viruses. Our results indicate that many drugs were developed from natural sources and many plants are yet to be explored for their potential specific and broad-spectrum effects against common viral infections.

Key words: viral infection, medicinal plants, traditional medicine, ethnopharmacology, ethnobotany

# ÖZET

İnsanlik tarihi boyunca varlığını muhafaza eden virülser, en bulaşıcı mikroorganizmalardan sayılabilir. İnsanları enfekte eden virüsler DNA veya RNA virüslerine ayrılabilir. İnsanlığın, mevcut COVID-19'a benzer pandemikleri önlemesine veya en aza indirmesine yardımcı olacak etkili tedavi yöntemleri geliştirmesi için, her bir virüs sınıfının viral replikasyon döngülerine ve enfeksiyon mekanizmalarına aşina olması önemlidir. Çoğu antiviral ilaçlar, geniş spektrumlu olup ortak viral enfeksiyon faktörlerini hedef alır. Ancak dünya genelinde eşit erişilebilirlik olmaması ve yüksek maliyet ile istenmeyen yan etkiler, alternatif tedavi yöntemlerine girmemizi teşvik etmektedir. Bu kısa inceleme, dünya çapında viral enfeksiyonları tedavi etmek için geleneksel tıbbi sistemlerde kullanılan bitkiler ve bunların *in-vitro* ve *invivo* farmakolojik çalışmalarına odaklanacaktır. DNA virüslerinden Herpesviridae ve Hepatit B virüsü için kullanılan tıbbi bitkiler ele alınırken, HIV, influenza ve Coronaviridae ailesini tedavi etmek için kullanılan bitkiler, RNA virüsleri kapsamında tartışıldı. Sonuçlarımız, doğal kaynaklardan birçok ilacın geliştirildiğini, ve birçok bitkinin yaygın viral enfeksiyonlara karşı potansiyel spesifik ila geniş spektrumlu etkileri için henüz yeterince araştırılmadığını göstermektedir.

Anahtar kelimeler: viral enfeksiyon,tibbi bitkiler,geleneksel tıp,etnofarmakoloji,etnobotani

Cilt/ Volume: 1 Sayı / Issue: 2 Yıl/Year: 2020 Yayıncı / Publisher Sağlık Bilimleri Üniversitesi University of Health Sciences

#### **INTRODUCTION**

Viruses are the most abundant biological entities on Earth and are widely present as parasites in cellular life. They date back to well before humankind's history. They are important agents of many infectious diseases most of which are transmitted from animals. Viruses depend entirely on living cells as hosts for all aspects of their life cycle (Waterson & Wilkinson, 1978).

Apart from their certain benefits, viruses cause a wide range of diseases in humans like the common colds and the rabies. They also play a role in several types of cancer (F. Huang et al., 2016). Viruses do not only cause harm for individuals they can also affect the well-being of Human societies. Smallpox and Immunodeficiency Virus (HIV) are two of the many examples that can be given. For this reason, understanding the nature of viruses, the mechanism of their replication and their pathogenesis is important for taking counteractive measures in the prevention, diagnosis, and treatment of viral ailments through the development of vaccines, diagnostic reagents and anti-viral drugs. Vaccines can save the lives of millions and bring about the end of viral diseases just like in the case of Smallpox virus which was eradicated (WHO, 2014).

Viruses can only replicate and produce new infectious viruses through the attachment and entry into the host cell. Production of viral mRNA and its translation by host ribosomes, genome replication and assembly and release of new viruses containing the genome, all takes place inside the host. The newly formed viruses can then spread and infect other cells (Wagner & Krug, 2020)

Viruses may have DNA or RNA as their genetic material. DNA viruses, such as chickenpox, Hepatitis B and Herpesvirus, can direct the host cell's replication proteins to synthesize new copies of the viral genome and to transcribe and translate that genome into viral proteins. While RNA viruses, such as rabies, hepatitis C and the Coronavirus, encode enzymes that can replicate RNA into DNA, which cannot be done by the host cell. These RNA polymerase enzymes are more likely to make copying errors

than DNA polymerases and, therefore, often make mistakes during transcription. For this reason, mutations in RNA viruses occur more frequently than in DNA viruses. This causes them to change and adapt more rapidly to their host (Lodish et al., 2000).

The culturing of the Herpes simplex virus was first carried out in 1925, which paved the way for the development of treatment and possibly prevention methods (Parker Jr & Nye, 1925). Acyclovir is a drug developed and patented in the 70's from nucleosides obtained from the Caribbean sponge, Cryptotethya crypta, and together with its derivates forms the basis for herpes treatment (Schaeffer, Gurwara, Vince, & Bittner, 1971). The general idea behind antiviral drug design is identifying targets such as proteins or parts of proteins, preferably those that are common among many viruses. Nowadays, it is possible to develop newer drugs suitable for new targets at the molecular level using computeraided design programs.

Antiviral drugs usually cause undesired side-effects vomiting, such as nausea, hallucinations, and worsening of asthma conditions (Stiver, 2003). As an example, the use of nucleoside reverse transcriptase inhibitors (NRTIs) in the treatment of AIDS can lead to hepatic failure and lactic acidosis (Lewis, Day, & Copeland, 2003). This, together with the emergence of resistant strains (Sheu et al., 2008; Wyles, 2013) urges the scientific community to explore and research more options for the discovery and development of antiviral drugs for the treatment of diseases and production of possible vaccines (Birkmann & Zimmermann, 2016; Treml et al., 2020). Drug discovery from natural resources is a practice adopted by many traditional medicine systems throughout mankind's history. Some of the major known system include Ayurveda, Unani, traditional Chinese medicine (TCM), Kampo, and Jamu. Although most of medicinal plant usage is recorded in pharmacopoeias and ancient texts, it is possible to access numerous ethnobotanical studies concerning the use of plants as medicines and food in different regions of the world and use

Cilt/ Volume: 1 Sayı / Issue: 2 Yıl/Year: 2020 Yayıncı / Publisher Sağlık Bilimleri Üniversitesi University of Health Sciences

these studies as a starting point for drug discovery.

Due to their low toxicity, renewable resources, biodegradability and in most cases low isolation of biologically the cost. active compounds from natural resources has gained more attention (Brahmachari, 2012). With the advent of new technologies in the fields of biology and chemistry – mostly the 'omics' – the pharmacological effects of potential drug substances can now be analyzed more thoroughly. Moreover, the more accurate establishment of synergic effects between natural products enables the development of newer methods for disease treatment and hopefully prevention. Bioactive compounds from medicinal plants were proven to be active against a wide range of virally infectious species, some of these compounds are flavonoids, coumarins, polyphenols, tannins, terpenoids, essential oils, alkaloids, polysaccharides and proteins (Abad, Bedoya, Apaza, & Bermejo, 2012; Bekut et al., 2018; Mukhtar et al., 2008; Naithani et al., 2008).

In light of the abundance of information available to us, this short review will attempt to summarize the available literature in relation to the most infectious viral diseases and the treatment strategies employed in this regard. The focus will be on plants used in traditional medical systems throughout the world, that were documented through ethnobtanical analysis. The active constituents and the related pharmacological studies on biological activity were also included in this study.

#### 1. METHODOLOGY:

Ethnopharmacology is defined as "the interdisciplinary scientific exploration of biologically active agents traditionally employed or observed by man" (Bruhn & Helmstedt, 1981; Rivera et al., 2014). Based on this, the present article will explore the treatment methods employed worldwide in traditional medicine systems. The most known databases were researched, namely PubMed, Science Direct, and Web of Science, using the keywords: "plant name", "disease name", "disease family", ethnobotany, ethnopharmacology, antiviral. herbal medicine, traditional, herb, extract. In

traditional medical records, plants are recorded based on the symptom they treat. Hence, symptoms of each disease as mentioned in the WHO fact sheets were used as search keywords as well.

The records can be subdivided according to the type of viral infection it addresses. In this article we will be focusing on the pandemic and endemic viral infections mentioned in the World Health Organization (WHO)'s website. Both review articles and scientific articles, based on recorded traditional uses and pharmacological evaluation, were summarized here. Articles were selected based on their metric, coverage, H index, and quartile. Moreover, the general quality and authenticity of the information based on other more reliable literature was taken into consideration.

#### 2. RESULTS AND DISCUSSION:

The results were discussed as shown in the sub sections below. Most research revolved around plants with potential activity against some of the most common viral infections namely those caused by Herpes Simplex virus (HSV), HIV, Hepatitis, A, B, and C viruses (HAV, HBV, HCV), and Influenza virus (IV). The most common DNA viruses were those belonging to the family Herpesviridae and the HBV. As for RNA viruses, the most common ones were IV, HIV, the viruses belonging to the Flaviviridae family, and last but not least, based on current events, the viruses from the Coronaviridae family, which will be discussed in the final part of the article, together with current treatment regimens being adopted during the newest COVID-19 pandemic.

#### 2.1. DNA viruses: Herpesviridae and Hepatitis B

The family Herpesviridae contains viruses such as Herpes Simplex (HSV), chicken pox, cytomegalovirus, Epstein-Barr virus, and Varicella zoster virus. It has a double stranded DNA enclosed in an icosahedral envelope (Denaro et al., 2019). Concurrently to its vast spread in the world, the number of studies on possible cures based on traditional medical systems is quite high. Two of the most common

Yayıncı / Publisher Sağlık Bilimleri Üniversitesi University of Health Sciences

diseases of the Herpesviridae family are HSV-1 and HSV2, and are recorded in WHO fact sheets as a lifelong infection, with an estimate HSV-1 infection for 67% of people under the age of 50 worldwide (WHO, 2017).

Hepatitis B (HB) is an infectious disease caused by the hepatitis B virus (HBV) that affects the liver (Logan & Rice, 1987; WHO, 2019a). It can cause both acute and chronic infections with over 750,000 deaths occurring from hepatitis B each year. HBV is a partially double-stranded DNA virus. а species of the genus Orthohepadnavirus, and a member of the Hepadnaviridae family of viruses (W.-S. Ryu, 2016). The virus particle (virion) consists of an outer lipid envelope and an icosahedral capsid (Locarnini, 2004). The virus is one of the smallest enveloped animal viruses. Vaccines for the prevention of hepatitis B have been routinely recommended for babies since 1991 in the United States (Schillie et al., 2013) and the first dose is generally recommended within a day of birth (Diseases, 2017).

Euphorbia spp. are known in many folk medicinal systems possess antiviral. to antifungal, and antibacterial activities. While some records, like the Rwandan folk medicine, show antiviral related use of its leaves and stems against poliomyelitis, aphtha, and diarrhea (Vlietinck et al., 1995) other medical systems, like that of the Amazonian people, have records of its use in skin ailments or other seemingly unrelated conditions (MACRAE, HUDSON, & TOWERS, 1988). Dioscorides's De Materia Medica contains a depiction of the collection of the latex of a Euphorbia plant (Appendino & Szallasi, 1997).

The triterpenes and steroid content of *E. denticulata* Lam., especially betulin, was effective against HSV-1, and it was indicated that the effects were exerted at 2.0 hours post-infection, where it is early on in the replication stage of the virus (Shamsabadipour et al., 2013). Another species named *E. spinidens* Bornm was also able to inhibit the earlier stages of HSV-1 replication (Karimi, Mohammadi-Kamalabadi, Rafieian-Kopaei, Amjad, & Salimzadeh, 2016). A study on the diterpenoids obtained from the

acetone extract of *E. milii* Linn. showed promising results against the Epstein-Barr virus's lytic replication (S. N. Liu et al., 2017). Kim and colleagues also recorded effectiveness of *Euphorbia spp.* against the Epstein-Barr virus (D. eun Kim et al., 2018). The tannin isolate named putranjivain A from *E. jolkini* was shown to have antiviral effect against HSV-2 at the later stages of viral replication (H. Y. Cheng et al., 2004).

Belonging to the family of Phyllanthaceae, another plant worth mentioning due to its frequent appearance in the literature is Phyllanthus urinaria L.. This plant was used in China's folk medicine as a remedy for hepatitis B, nephrolithiasis, pain, and jaundice (Wei et al., 2005), while in Trinidad and Tobago, it is used to treat diarrhea (Lans, 2007). In the Peruvian Amazon, decoctions of the plant are drunk to treat hepatitis (Roumy et al., 2020). The roots and stem of this plant were also used against HBV among other problems in Korea (Bae, Kim, & Choi, 2009). Recorded pharmacological studies exist on its activity against HSV-1 and HSV-2 due to its tannin content (Hua-Yew Cheng et al., 2011; C.-M. Yang, Cheng, Lin, Chiang, & Lin, 2005; C. M. Yang, Cheng, Lin, Chiang, & Lin, 2007). Other species in the genus Phyllanthus were reviewed recently for their active constituents and biological effects (Qi, Hua, & Gao, 2014). According to a review, there is recorded use of *P*. orbicularis against bovine herpesvirus-1(BHV-1), HSV-2, and adenovirus-7 (AV-7) (Del Barrio & Parra, 2000). Additionally, in another study, the plant was shown to have inhibitory effect against the early replication and DNA synthesis of HSV-2, and this effect was attributed to its epicatechin and procyanidin content (Álvarez et al., 2012). A research was carried out to assess the in-vitro and in-vivo effects of P. niruri L. against HBV. The results indicate that the lignan content named niranthin, could be effective in reducing the viral load and offer hepatoprotective effects (S. Liu et al., 2014). However, one clinical trial was conducted to test the plant on patients with chronic HBV and no significant differences were observed in the viral loads of the treatment and placebo groups (Baiguera et al., 2018).



Yayıncı / Publisher Sağlık Bilimleri Üniversitesi University of Health Sciences

*Morus alba*, a plant mentioned in the Chinese pharmacopoeia (Pharmacopoeia Commission of People's Republic of China, 2000) to remove lung heat and relieve asthma, has been studied for the biological activity of its various parts. Of interest is a research conducted in 2003 that aimed to characterize the root bark extract's active constituents and assess its antiviral activity. The results showed potent activity of the flavonoid Leachianone G against HSV-1 (Du et.al, 2003).

The use of Quercus brantii's against diarrhea and stomach ulcers were recorded (Mosaddegh, Naghibi, Moazzeni, Pirani, & Esmaeili, 2012). Although stomach ulcers maybe unrelated to the underlying viral infectious diseases mechanisms, it is well known that diarrhea (acute gastroenteritis) is one of the symptoms in some virally infectious diseases such as norovirus, cytomegalovirus, viral hepatitis, and rotavirus (Grytdal et al., 2016). Thus, it is natural to infer the possible presence of antiviral activity of this plant. Indeed, the activity of Q. brantii extract against the herpes simplex virus was studied to reveal its prevention of the viral entry into the host cell due to its condensed tannin content (Karimi, Rafieian-Kopaei, Moradi, & Alidadi, 2017). Additional details can be found in Table 2.1.

# 2.2: RNA viruses

#### 2.2.1: HIV

Belonging to the Retroviridae family, HIV is a virus that attacks the body's immune system, specifically the white blood cells knows as CD4. HIV is significant due to its major role in Acquired Immunodeficiency Syndrome (AIDS) (Douek, Roederer, & Koup, 2009; Weiss, 1993), which enables the virus to thrive through progressive failure of the immune system via lifethreatening opportunistic infections and cancers (Powell et al., 2016).

HIV has a spherical shape around 60 times smaller than a red blood cell, with a diameter of about 120 nm (Harvey, 2007), and two copies of positive-sense single-stranded RNA enclosed in a conical capsid (Kuiken et al., 2009). The treatment regimens include the use of multiple

antiretroviral drugs known as antiretroviral therapy (ART) which will reduce the viral load in the blood to an undetectable level and prevent transmission by 96%. Due to the advances, it is now considered a chronic condition with increasingly rarer instances of progression to AIDS. However, the ART should be taken every day in a patient's lifetime, hence with loss of access to ART, alternative medicines must be considered (WHO, 2019b).

One of the methods of suppressing HIV is by inhibiting the virus's reverse transcriptase (RT) enzyme. According to Chinsembu's research, most RT inhibitors were present in plants belonging to the Lamiaceae (13.7%), Fabaceae (10.7%), Euphorbiaceae (9.9%), Clusiaceae (6.1%),Asteraceae (4.6%), Combretaceae (4.6%),Moraceae (3.0%)families and (Chinsembu, 2019). Among the Lamiaceae family, the subfamilies Neptoideae and Lamiodieae containing rosmarinic acid and iridoids respectively have been studied for the antibacterial, antifungal, and antioxidant properties of their essential oils. Among plants that possess anti-RT activity, anti-HIV protease and integrase activities, and DNA copying inhibitory activity for HIV are Thymus sp., Mentha sp., Rosmarinus sp., Melissa sp., Origanum sp., Ocimum sp., Prunella sp., Hyssopus sp., Salvia sp., and Saturja sp. (Bekut et al., 2018).

During the early middle ages, it was common for emperors to establish medicinal herb gardens either in their palaces or in monasteries. One such garden was ordered by King Charles the Great (747-814). The garden named Capitulare de villis contained majority of the plant species of the family Lamiaceae mentioned above, due to their medicinal value (Heinrich et.al, 2018). Mentha sp. were used for headache, bronchitis, lung inflammation, and for improving the general health among Albanians and Serbians from South Kosovo (Mustafa, Hajdari, Pulaj, Quave, & Pieroni, 2020). On the Javor Mountain of Bosnia and Herzegovina Lamiacaea family plants are used for cough, asthma, respiratory infections, fever, and cold (Savić, Mačukanović-Jocić, & Jarić, 2019)

## Table 2.1: Plants used traditionally against DNA viruses

Diant Family	Scientific	Active	Traditional use	References
	name	ingredient(s)	<b>Biological activity</b>	
Aloaceae	Aloe spp.	Aloin B and aloe- emodin	Cold, fever (Zulu medicine) Anticatarrhal (Catalonia)	(Hutchings, 1996) (Bonet, Parada, Selga, & Vallès, 1999)
			Chest complaints (Namibia) Fever, lung disorders (Pakistan) Fever/malaria (Uganda)	(Koenen, 2001) (Yaseen et al., 2015) (Anywar et al., 2020)
			Anti HBV	(Parvez et al., 2019)
Anacardiaceae	Rhus spp.	Gallic acid	Arthritis (Thailand)	(Salguero, 2003)
			Purgative, lung rejuvenator,	(Djakpo & Yao, 2010)
			anti-asthma (Bhutan)	
			Anti HSV-1 and HSV-2 activity	(Reichling, Neuner, Sharaf, Harkenthal, & Schnitzler, 2009)
Araliaceae	Hydrocotyle sibthorpioides	Asiaticoside	Hepatitis, herpes (TCM)	(Au et al., 2008)
			Anti-HBV	(Q. Huang et al., 2013)
Asteraceae	Artemisia capillaris Thunb.	Pumilaside A and enediyne glucosides	Liver disease, indigestion,	(H. Kim & Song, 2011)
			Liver disease hepatitis (India)	(Thakur et al. 2016)
			Jaundice, hepatitis (Mauritius)	(Suroowan, Pynee, &
				Mahomoodally, 2019)
			Anti-HBV	(Zhao et al., 2014)
			Anti-HBV	(Geng et al., 2018)
	Euphorbia spp. Quercus brantii Lindl	4,4 dimethyl steroids Condensed tannins	Sore throats (China)	(Zheng & Xing, 2009)
			Cough, bronchial infections,	(Gairola, Sharma, & Bedi,
			Kashmir and India)	2014)
Euphorbiaceae			Fungal infections, sores (Jammu	(Shah, Bharati, Ahmad, &
			and Kashmir)	Sharma, 2015)
			Anti HSV-1	(Shamsabadipour et al., 2013)
				(B. Wang et al., 2018)
			Anti EV/1 and anti HRV	
Fagaceae			Stomach ulcers, diarrhea (Iran)	(Mosaddegh et al., 2012)
			Anti HSV-1	(Karimi et al., 2017)
	<i>Prunus dulcis</i> (Mill.) D.A. Webb	Quercetin, epicatechin, and catechin	(Catalonia)	(Agelet & Valles, 2003)
Rosaceae			Sore throat (southern Italy)	(Pieroni, Quave, & Santoro,
			Cough (southern Italy)	(Scherrer, Motti, & Weckerle, 2005)
			Antitussive (eastern Mallorca)	(Carrió & Vallès, 2012)
			Anti HSV-1	(Bisignano et al., 2017)
			Anti HSV-1	(Musarra-Pizzo et al., 2019)
Saururaceae	<i>Houttuynia</i> <i>cordata</i> Thunb.	Quercetin, quercitrin or isoquercitrin and houttuynoid A	Herpes, smallpox (China)	(S. Li et al., 2006)
			Cough four infant four	(Au et al., 2008)
			(Laos)	(De Boer, Lamxay, & Bjork, $2012$ )
			Common cold (Thailand)	(Panyadee, Balslev.
			· · · · · · · · · · · · · · · · · · ·	Wangpakapattanawong, & Inta,
				2019)
			Anti HSV-2	(X. Chen et al., 2011)
			Anti HSV-1	(T. Li et al., 2017)

HSV-1: Herpes Simplex virus type 1; HSV-2: Herpes Simplex virus type 2; EV71: Enterovirus 71; HRV: Human rhinovirus; HBV: Hepatitis B virus.

Yayıncı / Publisher Sağlık Bilimleri Üniversitesi University of Health Sciences

According to Ibn al-Baytar's (1197-1248) *Kitab al-Yami' li-mufradat al-adwiya wa-l-aghdiya* (the *Compendium of Simple Medicaments and Foods*), most members of the Lamiaceae family were used for digestive followed by respiratory illnesses, while the most mentioned plant is *Ocimum basilicum* (El-Gharbaoui, Benítez, González-Tejero, Molero-Mesa, & Merzouki, 2017). The Rasta bush doctors in the Western Cape of South Africa use species of this family for hypertension, cancer, ulcers, gout, general health tonic, respiratory ailments, asthma, bronchitis, cold, fever, cough, and diarrhea (Aston Philander, 2011).

Hyssopus officinalis was shown to inhibit HIV replication due to its polysaccharide type compound content (Kreis, Kaplan, Freeman, Sun, & Sarin, 1990). The use of the whole plant of Melissa officinalis L. was recorded as an anti-HIV agent. The extracts of M. officinalis L., Mentha  $\times$  piperita L., and Salvia officinalis L. exhibited strong antirviral effects by inhibiting the viral replication at its earlier stages prior to entry into host (Geuenich et al., 2008). Some studies conducted address the latent HIV which may later resurface and lead to infections. One such drug being considered for human clinical trials is prostratin, which is an ester isolated from the tropical plant Homalanthus nutans of the Euphorbiaceae family. This compound can block HIV infections as well as induce the latent virus's expression (Kulkosky et al., 2001; Rullas et al., 2004; Salehi et al., 2018). Triterpenes isolated from Ocimum labiatum was shown to have a similar effect on activating latent HIV, and thus having the potential to be used as an adjuvant in ART (Kapewangolo, Omolo, Fonteh, Kandawa-Schulz, & Meyer, 2017).

Artemisinin, a compound isolated from the plant *Artemisia annua* L., which is a famous traditional plant in the Chinese folk systems, has gained popularity, together with its semisynthetic forms, for its major role in the treatment of malaria disease caused by a parasitic worm. Apart from being an approved medicine by the

Food and Drug Administration (FDA) of the United States, it has produced a Nobel prize in 2015 in medicine for its discoverers. When analyzing the local uses of this plant, a survey conducted by Willcox in 2011 found that most local people of Kenya and Uganda used this plant for conditions other than malaria as well, such as diarrhea, abdominal pain, and HIV/AIDS (Willcox et al., 2011). *A. annua* tea infusions were shown to possess anti-HIV activity *in-vitro* with low toxicity (Lubbe, Seibert, Klimkait, & Van Der Kooy, 2012). A few more plants were included in table 2.2 below.

#### 2.2.2: Influenza viruses

Influenza viruses have originated from the Spanish flu in 1918, and swine flu in 2009 (Sriwilaijaroen et al., 2012). These viruses belong to the Orthomyxoviridae family and have single stranded negative sense RNA enclosed in capsids (Denaro et al., 2019). Due to the high mutation rates of these virus they reoccur seasonally as flu and cold and are an important category of viruses that need continuous attention for cure and vaccine development. Drug development for influenza viruses targets the neuroaminidase (NA) protein present on the virus which is essential for its replication (Denaro et al., 2019). Some of the plants used traditionally and researched for their pharmacological properties will be discussed here.

An interesting study on Aloe species was conducted Glatthaar-Saalmüller by and colleagues on the use of Biaron  $C^{\mathbb{R}}$  (a product containing Aloe arborescens Mill., Vitamin C, and Aronia melanocarpa Elliot. succus) against viruses causing upper respiratory tract infections. The results showed excellent inhibitory effect against the replicatory ability of orthomyxoviridae - influenza A and influenza B viruses, as well as two non-enveloped RNA viruses belonging to the Picornaviridae family (Glatthaar-Saalmüller et al., 2015).

Plant family	Scientific name	Active	Traditional use	References
		ingredient(s)	Biological activity	
Acanthaceae	Andrographis paniculate Nees	Andrographolide	Liver disease (Ayurveda) Rid the body of heat in fevers and dispel toxins (TCM) Common colds (Scandinavian	(Chturvedi, Tomar, Tiwari, & Singh, 1983) (Reddy et al., 2005) (Caceres, Hancke, Burgos,
			countries)	& Wikman, 1997)
			Increase in the level of CD4 <sup>+</sup> lymphocytes	(Calabrese et al., 2000)
			Anti-HIV activity	(Reddy et al., 2005)
Acanthaceae	Avicenna marina	Iridoid glycoside	Snakebites, wounds, abscess, blotch (Iran) Ulcers, rheumatism, smallpox	(Safa et al., 2013) (Duke, 1991)
			(Australasian countries) Skin parasites and gangrenous wounds (India)	(Mohan, Rao, & Pragada, 2014)
			Anti-HIV-1 replication	(Behbahani, 2014)
Clusiaceae	Calophyllum brasiliense	Tricyclic coumarin	Warts (Brazil) General infections, indigestion (Brazil)	(Lago et al., 2016) (Ribeiro, Bieski, Balogun, & Martins, 2017)
			Inhibits viral replication by suppressing NF-kB activation Inhibits HIV-1 RT*	(Kudo et al., 2013) (César et al., 2011)
Fabaceae	Acacia spp.	Acaconin protein, catechins, kaempferol, rutin, ferulic acid and caeffic acid	Flu/cold, cough, headache, chicken pox (Argentina) Diarrhea, chest problems, cold, flu (South Africa)	(Suárez, 2019) (Mhlongo & Van Wyk, 2019)
			Inhibits HIV RI* Inhibits viral protease	(Lam & Ng, 2010) (Modi et al., 2013)

\*Reverse transcriptase

Many other studies were conducted on these species and the active constituents were analyzed. For example, the polysaccharide section rich in mannose unit, namely acemamman, was shown to improve wound healing, modulate immunity and exert antiviral effects (Chandegara & Varshney, 2013). The mechanisms studied prevention include of virus adsorption, attachment, and entry into the cell, while immune function is thought to be modulated via its direct effect on the effectivity of T helper cells (Sánchez-Machado, López-Cervantes, Sendón, & Sanches-Silva, 2017)Kampo medicine which was mostly practiced in ancient China and was introduced into Japan in the fifth or sixth century has taken its current form towards the seventeenth century. Kampo drugs consist of a mixture of plants, fungi, minerals, and insects (TSUMURA & CO., 2016). A study conducted

on several Kampo drugs (maoto, kakkonto, jinkokato, senkyuchachosan, and bakumondoto) used for common cold and influenza, and the crude drugs Glycyrrhizae radix, Atractylodis lanceae rhizome, Rehmanniae radix, Citri unshiu pericarpium, Cnidii rhizome, and Saposhnikoviae radix. The Kampo drugs were shown to inhibit replication and spread of the virus. Glycyrrhizae radix, which was present in all the Kampo drugs tested, showed cytotoxicity at very high concentrations only but had the strongest viral inhibitory effects due to the chalcone isoliquiritigenin and the flavanone liquiritigenin (Nomura et al., 2019). Triterpenoid saponins of the plant were previously reported to have antiinfluenza effects (Ji et al., 2016).

A commercially available essential oil blend consisting of oils of orange, clove, cinnamon, eucalyptus, and rosemary were tested to be

effective against the influenza virus's proteins (Wu et al., 2010). Eucalyptus was recorded in Zulu medical system as a remedy for headache, diarrhea, cold, and flu (Mhlongo & Van Wyk, 2019) and the leaves are used to treat flu, fever, and common cold as well as an antitussive in Colombian medicine folk (Ceuterick, Vandebroek, Torry, & Pieroni, 2008). Rosemary (Rosmarinus officinalis L.) was recorded in the north-eastern Iberian region for its use against colds, to enhance circulation, for pain in muscles, bones and joints, and for laziness (Parada, Carrió, Bonet, & Vallès, 2009).

High throughput screening was conducted on 50 plants used traditionally in different parts of Sarawak, Malaysia. The plants had recorded uses against flu, cold, fever, cough, headache, conjunctivitis, sore throat, and some possible effects against AIDS. Based on the results of the study, 11 plants had significant inhibitory effects against two strains of influenza by inhibiting both the hemagglutinin and NA indication activity against the virus in the early stages of replication. Synergistic effects due to the rich chemical constituents of the extracts were thought to be a reason for the multiple inhibition mechanisms of the extracts towards the viral strains (Rajasekaran et al., 2013).

In southern mountainous regions of Korea, Thuja orientalis L. and Aster sphathulifolius Maxim. were used in common cold, fever, and generally the respiratory system disorders (H. Kim & Song, 2011). A study conducted on the antiviral activities against these plants showed activity, probably through blockage of the virus attachment to the host cell or the inhibition of viral replication (Won, Lee, Song, & Poo, 2013). Finally, it is worth mentioning that some plants show broad range of activity towards several viruses. One such plant is mentioned in table 2.1 above, namely Houttuynia cordata Thunb., which showed anti influenza virus activity due to its polysaccharide and flavonoid contents (Ling et al., 2020).

# 2.2.3: SARS, MERS, and the novel Coronavirus (COVID-19)

Coronavirus is in the Coronaviridae family causing primarily infections in birds and animals (zoonotic infections) but can also be transmitted to humans and lead to infections in them too. This crossing of the species barrier was evident in the outbreaks of both severe acute respiratory syndrome (SARS) in 2002 and Middle East respiratory syndrome (MERS) in 2012 which caused thousands of deaths (Schoeman & Fielding, 2019).

Both SARS-CoV and MERS-Cov are in the  $\beta$ -coronavirus family (Alimuddin Zumla, David S Hui, 2020). The recent outbreak of COVID-19 at the end of 2019 which happened in Wuhan city, China (Cohen & Normile, 2020), is a result of a highly contagious infection by the SARS-CoV-2 virus among humans (Lu, 2020), has a higher mortality rate than its predecessors, and poses a great problem for humanity as new treatment modalities and a vaccine are trying to be developed to end its killing spree.

The genetic material of the new coronavirus (COVID-19) is a single-stranded positive-sense RNA and belongs to the  $\beta$ -coronavirus family (N. Chen et al., 2020) showing great similarity (%82) to the SARS-CoV (Lu, 2020). Currently, there is no specific treatment for COVID-19. Until the specific antiviral drugs are available, broadspectrum antivirals like the Nucleoside analogues and HIV-protease inhibitors are used in the hopes to relieve the course of the disease (Lu, 2020). The treatments that have so far been attempted showed that patients were administrated existing antiviral drugs with a course of treatment including twice a day of oral administration of 75 mg oseltamivir, 500 mg lopinavir, 500 mg ritonavir and the intravenous administration of 0.25 g ganciclovir for 3–14 days (N. Chen et al., 2020). Another report showed that the broadspectrum antiviral remdesivir and chloroquine are highly effective in the control of 2019-nCoV infection in vitro (M. Wang et al., 2020). However, more research needs to be carried out to find a new specific treatment for the COVID-19. The most recent development was the FDA approval of the anti-parasitic drug ivermectin



Yayıncı / Publisher Sağlık Bilimleri Üniversitesi University of Health Sciences

based on the promising results obtained in *in-vitro* trials. The drug was shown to reduce the viral RNA levels greatly 2 hours post-infection. It was shown to be effective against other viruses such as HIV, influenza A, and dengue (Caly, Druce, Catton, Jans, & Wagstaff, 2020)

saikosaponins Previously (triterpene glycosides) from medicinal plants were shown to be active against the activity of HCoV-22E9 by preventing viral attachment and penetration, as well as earlier stages of replication (P. W. Cheng, Ng, Chiang, & Lin, 2006). Extracts obtained from Lycoris radiata, A.annua, Pyrrosia lingua, and *Lindera aggregate* which are medicinal plants used in TCM have been documented for their activity against SARS-CoV (S. Y. Li et al., 2005). Helicase and protease enzymes together with myricetin, scuttellarein, and other phenolic compounds from Isatis indigotica and Torreya mucifera were active against the SARS-CoV enzymes (Lin et al., 2005; Y. B. Ryu et al., 2010; Yu et al., 2012). The water extract of *H. cordata* mentioned in table 3.1 has shown inhibition against SARS-CoV by inhibiting the viral protease and RNA-dependent RNA polymerase activities (Lau et al., 2008).

According to a recent review on the use of Chinese traditional medicine for the treatment of the new COVID-19, different herbal drugs were used for the treatment of different stages of the disease (Ang, Lee, Choi, Zhang, & Lee, 2020). For the mild stages, two drugs were mentioned namely Magnolia officinalis cortex and Platycodonis cortex. These two plants were normally prescribed for gastrointestinal problems and coughs respectively (Xi & Gong, 2017). The *in-vivo* effects of these two plants were previously established for their antiinflammatory activity in acute lung injury models (Guo et al., 2018; Hu et al., 2017).

It is important to mention *Sambucus spp.* (elderberry) during these times. The species was used in southern Kosovo for respiratory diseases, asthma, and for general health (Mustafa et al., 2020), and in Bosnia and Herzegovina its use for fevers, colds, and high temperatures was recorded (Savić et al., 2019). *S. nigra* was used for pain, bronchitis, stomach infections,

headache, diarrhea, and flu in the high river Ter valley of Catalonia (Rigat, Bonet, Garcia, Garnatje, & Vallès, 2007). The plant has been extensively reviewed thanks to the efforts of Porter and Bode, where its active constituents, the related in-vitro and in-vivo studies, and the approved products that can be accessed currently were reviewed in the research (Porter & Bode, 2017). The plant is mainly used for influenza like conditions, however, a thesis that was recently published mentions the potential use of the plant for viruses like the novel SARS-CoV-2. According to the thesis, the use of S. nigra L. extract on chicken coronaviruses with promising results on the inhibition of the earlier replication stages of the virus may indicate its potential effectiveness against human coronaviruses too, if further studied (Wermig-morgan, 2020).

A severe respiratory illness outbreak during the winter of 2017, in a long-term care facility in Louisiana has been reported and was associated with a human coronavirus named HCoV-NL63 (Hand et al., 2018). A study was conducted to assess the effectiveness of another elderberry species named *S. Formosona*Nakai against this virally infectious outbreak (Weng et al., 2019). The study results indicate a significant reduction of the viral yield, plaque formation, and viral attachment to the host cell. The effects were attributed to the caffeic acid, chlorogenic acid, and gallic acid contents of the plant.

The current COVID-19 pandemic is the focus point of the scientific community as of April 2020. On February 17, the Chinese State Council announced its endorsement of the possible beneficial effects of chloroquine phosphate, which is a structural analogue of quinine, the famous anti-malarial drug. The compound is isolated from barks of Cinchona *spp.*. Ancient uses of the tree parts mainly revolve around its use against fevers, such as those tales told in the Peruvian amazon (Urdang, 1945)), and its use in indigenous communities of South America (Thompson, 1928). The antiviral effects of chloroquine phosphate have been tested in more than 10 hospitals in China and has shown to alleviate symptoms and to speed up the virus Ì

Journal of Integrative and Anatolian Medicine Bütünleyici ve Anadolu Tibbı Dergisi Yayıncı / Publisher Sağlık Bilimleri Üniversitesi University of Health Sciences

seroconversion ("Redeploying plant defences," 2020).

## CONCLUSION

This review focused on some of the most infectious viral disease that were endorsed by the WHO, and to date, are potential sources of epidemics or pandemics if not studied carefully for the potential treatment and prevention options. We have tried to compile as many plants as we could find that were adopted in traditional medical systems used worldwide, and that were additionally studied for their inhibition of the viruses mentioned.

Due to the vast amount of data available, it is important to apply high throughput screening techniques and make use of the up-to-date technological facilities in order to narrow down and weed out the wealth of information we have on hand in this 21<sup>st</sup> century. On this note, our suggesting for researchers would be to focus on broad range therapeutics such as Aloe sp., Hottuynia sp., and Phyllanthus sp., just to name a few. The advantage would be to gain a better understanding of the general mechanisms of plant metabolite functions, as well as to produce possible treatment modalities that remind its importance and urgency in times like the current COVID-19 outbreak. The study can be improved by carrying out a more comprehensive and systematic review on more plants and with easier access to ancient traditional medical texts.

#### REFERENCES

- Abad, M. J., Bedoya, L. M., Apaza, L., & Bermejo, P. (2012). Anti-infective flavonoids: an overview. In *Bioactive Natural Products: Opportunities and Challenges in Medicinal Chemistry* (pp. 443–473). World Scientific.
- Agelet, A., & Vallès, J. (2003). Studies on pharmaceutical ethnobotany in the region of Pallars (Pyrenees, Catalonia, Iberian Peninsula). Part III. Medicinal uses of nonvascular plants. *Journal of Ethnopharmacology*, 84(2–3), 229–234.

- Alimuddin Zumla, David S Hui, S. P. (2020). Middle East respiratory syndrome. *The Lancet*, 386(9997), 995–1007.
- Álvarez, Á. L., Dalton, K. P., Nicieza, I., Diñeiro, Y., Picinelli, A., Melón, S., ... Parra, F. (2012). Bioactivity-guided fractionation of phyllanthus orbicularis and identification of the principal anti HSV-2 compounds. *Phytotherapy Research*, 26(10), 1513–1520.
- Ang, L., Lee, H. W., Choi, J. Y., Zhang, J., & Lee, M. S. (2020). Herbal medicine and pattern identification for treating COVID-19: a rapid review of guidelines. *Integrative Medicine Research*, 9(2), 100407.
- Anywar, G., Kakudidi, E., Byamukama, R., Mukonzo, J., Schubert, A., & Oryem-Origa, H. (2020). Indigenous traditional knowledge of medicinal plants used by herbalists in treating opportunistic infections among people living with HIV/AIDS in Uganda. *Journal of Ethnopharmacology*, 246(August 2019), 112205.
- Appendino, G., & Szallasi, A. (1997). Euphorbium: Modern research on its active principle, resiniferatoxin, revives an ancient medicine. *Life Sciences*, 60(10), 681–696.
- Aston Philander, L. (2011). An ethnobotany of Western Cape Rasta bush medicine. *Journal* of Ethnopharmacology, 138(2), 578–594.
- Au, D. T., Wu, J., Jiang, Z., Chen, H., Lu, G., & Zhao, Z. (2008). Ethnobotanical study of medicinal plants used by Hakka in Guangdong, China. Journal of Ethnopharmacology, 117(1), 41–50.
- Bae, K.-H., Kim, J.-A., & Choi, Y.-E. (2009). Induction and in vitro proliferation of adventitious roots in Dendropanax morbifera. *Journal of Plant Biotechnology*, 36(2), 163– 169.
- Baiguera, C., Boschetti, A., Raffetti, E., Zanini,
  B., Puoti, M., & Donato, F. (2018).
  Phyllanthus niruri versus Placebo for Chronic Hepatitis B Virus Infection: A Randomized Controlled Trial. *Complementary Medicine Research*, 25(6), 376–382.



- Behbahani, M. (2014). Evaluation of anti-HIV-1 activity of a new iridoid glycoside isolated from Avicenna marina, in vitro. *International Immunopharmacology*, *23*(1), 262–266.
- Bekut, M., Brkić, S., Kladar, N., Dragović, G., Gavarić, N., & Božin, B. (2018). Potential of selected Lamiaceae plants in anti(retro)viral therapy. *Pharmacological Research*, 133, 301–314.
- Birkmann, A., & Zimmermann, H. (2016). HSV antivirals–current and future treatment options. *Current Opinion in Virology*, *18*, 9– 13.
- Bisignano, C., Mandalari, G., Smeriglio, A., Trombetta, D., Pizzo, M. M., Pennisi, R., & Sciortino, M. T. (2017). Almond skin extracts abrogate HSV-1 replication by blocking virus binding to the cell. *Viruses*, 9(7), 1–15.
- Bonet, M. À., Parada, M., Selga, A., & Vallès, J. (1999). Studies on pharmaceutical ethnobotany in the regions of L'Alt Emporda and Les Guilleries (Catalonia, Iberian Peninsula). *Journal of Ethnopharmacology*, 68(1–3), 145–168.
- Brahmachari, G. (2012). *Bioactive natural products: opportunities and challenges in medicinal chemistry*. World Scientific.
- Bruhn, J. G., & Helmstedt, B. (1981). Ethnopharmacology: objectives, principles and perspectives. *Natural Products as Medicinal Agents*.
- Caceres, D. D., Hancke, J. L., Burgos, R. A., & Wikman, G. K. (1997). Prevention of common colds with Andrographis paniculata dried extract. A pilot double blind trial. *Phytomedicine*, 4(2), 101–104.
- Calabrese, C., Berman, S. H., Babish, J. G., Xinfang, M., Shinto, L., Dorr, M., ... Standish, L. J. (2000). A phase I trial of andrographolide in HIV positive patients and normal volunteers. *Phytotherapy Research*, 14(5), 333–338.
- Caly, L., Druce, J. D., Catton, M. G., Jans, D. A., & Wagstaff, K. M. (2020). The FDAapproved Drug Ivermectin inhibits the

replication of SARS-CoV-2 in vitro. *Antiviral Research*, *178*(March), 104787.

- Carrió, E., & Vallès, J. (2012). Ethnobotany of medicinal plants used in Eastern Mallorca (Balearic Islands, Mediterranean Sea). *Journal of Ethnopharmacology*, 141(3), 1021–1040.
- César, G. Z. J., Alfonso, M. G. G., Marius, M. M., Elizabeth, E. M., Ángel, C. B. M., Maira, H. R., ... Ricardo, R. C. (2011). Inhibition of HIV-1 reverse transcriptase, toxicological and chemical profile of Calophyllum brasiliense extracts from Chiapas, Mexico. *Fitoterapia*, 82(7), 1027–1034.
- Ceuterick, M., Vandebroek, I., Torry, B., & Pieroni, A. (2008). Cross-cultural adaptation in urban ethnobotany: The Colombian folk pharmacopoeia in London. *Journal of Ethnopharmacology*, *120*(3), 342–359.
- Chandegara, V. K., & Varshney, A. K. (2013). Aloe vera L. processing and products: A review. *Int. J. Med. Arom. Plants*, *3*(4), 2249– 4340. Retrieved from http://www.openaccessscience.com
- Chen, N., Zhou, M., Dong, X., Qu, J., Gong, F., Han, Y., ... Zhang, L. (2020).
  Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *The Lancet*, 395(10223), 507–513.
- Chen, X., Wang, Z., Yang, Z., Wang, J., Xu, Y., Tan, R. xiang, & Li, E. (2011). Houttuynia cordata blocks HSV infection through inhibition of NF-κB activation. *Antiviral Research*, 92(2), 341–345.
- Cheng, H. Y., Lin, T. C., Yang, C. M., Wang, K.
  C., Lin, L. T., & Lin, C. C. (2004).
  Putranjivain A from Euphorbia jolkini inhibits both virus entry and late stage replication of herpes simplex virus type 2 in vitro. *Journal of Antimicrobial Chemotherapy*, 53(4), 577–583.
- Cheng, P. W., Ng, L. T., Chiang, L. C., & Lin, C. C. (2006). Antiviral effects of saikosaponins on human coronavirus 229E in vitro. *Clinical*

and Experimental Pharmacology and Physiology, 33(7), 612–616.

- Chinsembu, K. C. (2019). Chemical diversity and activity profiles of HIV-1 reverse transcriptase inhibitors from plants. *Brazilian Journal of Pharmacognosy*, 29(4), 504–528.
- Chturvedi, G. N., Tomar, G. S., Tiwari, S. K., & Singh, K. P. (1983). Clinical studies on kalmegh (andrographis paniculata nees) in infective hepatitis. *Ancient Science of Life*.
- Cohen, J., & Normile, D. (2020). New SARS-like virus in China triggers alarm. *Science*.
- De Boer, H. J., Lamxay, V., & Björk, L. (2012). Comparing medicinal plant knowledge using similarity indices: A case of the Brou, Saek and Kry in Lao PDR. *Journal of Ethnopharmacology*, 141(1), 481–500.
- Del Barrio, G., & Parra, F. (2000). Evaluation of the antiviral activity of an aqueous extract from Phyllanthus orbicularis. *Journal of Ethnopharmacology*, 72(1–2), 317–322.
- Denaro, M., Smeriglio, A., Barreca, D., De Francesco, C., Occhiuto, C., Milano, G., & Trombetta, D. (2019). Antiviral activity of plants and their isolated bioactive compounds: An update. *Phytotherapy Research*, (May), 1–27.
- Diseases, C. on I. (2017). Elimination of perinatal hepatitis B: providing the first vaccine dose within 24 hours of birth. *Pediatrics*, *140*(3), e20171870.
- Djakpo, O., & Yao, W. (2010). Rhus chinensis and Galla Chinensis - Folklore to modern evidence: Review. *Phytotherapy Research*, *24*(12), 1739–1747.
- Douek, D. C., Roederer, M., & Koup, R. A. (2009). Emerging Concepts in the Immunopathogenesis of AIDS. *Annual Review of Medicine*, 60(1), 471–484.
- Duke, N. C. (1991). A systematic revision of the mangrove genus avicennia (Avicenniaceae) in Australasia. *Australian Systematic Botany*, *4*(2), 299–324.
- El-Gharbaoui, A., Benítez, G., González-Tejero, M. R., Molero-Mesa, J., & Merzouki, A.

(2017). Comparison of Lamiaceae medicinal uses in eastern Morocco and eastern Andalusia and in Ibn al-Baytar's Compendium of Simple Medicaments (13th century CE). *Journal of Ethnopharmacology*, *202*(March), 208–224.

- Gairola, S., Sharma, J., & Bedi, Y. S. (2014). A cross-cultural analysis of Jammu, Kashmir and Ladakh (India) medicinal plant use. *Journal of Ethnopharmacology*, 155(2), 925–986.
- Geng, C. A., Yang, T. H., Huang, X. Y., Yang, J., Ma, Y. B., Li, T. Z., ... Chen, J. J. (2018).
  Anti-hepatitis B virus effects of the traditional Chinese herb Artemisia capillaris and its active enynes. *Journal of Ethnopharmacology*, 224(June), 283–289.
- Geuenich, S., Goffinet, C., Venzke, S., Nolkemper, S., Baumann, I., Plinkert, P., ... Keppler, O. T. (2008). Aqueous extracts from peppermint, sage and lemon balm leaves display potent anti-HIV-1 activity by increasing the virion density. *Retrovirology*, *5*, 1–16.
- Glatthaar-Saalmüller, В., Fal, A. M., Schönknecht, K., Conrad, F., Sievers, H., & Saalmüller, A. (2015). Antiviral activity of an aqueous extract derived from Aloe arborescens Mill. against a broad panel of viruses causing infections of the upper respiratory tract. Phytomedicine, 22(10), 911-920.
- Grytdal, S. P., DeBess, E., Lee, L. E., Blythe, D., Ryan, P., Biggs, C., ... Hall, A. J. (2016).
  Incidence of norovirus and other viral pathogens that cause acute gastroenteritis (AGE) among kaiser permanente member populations in the United States, 2012-2013. *PLoS ONE*, 11(4), 2012–2013.
- Guo, S., Jiang, K., Wu, H., Yang, C., Yang, Y.,
  Yang, J., ... Deng, G. (2018). Magnoflorine ameliorates lipopolysaccharide-induced acute lung injury via suppressing NF-κB and MAPK activation. *Frontiers in Pharmacology*.



Yayıncı / Publisher Sağlık Bilimleri Üniversitesi University of Health Sciences

- Hand, J., Rose, E. B., Salinas, A., Lu, X., Sakthivel, S. K., Schneider, E., & Watson, J. T. (2018). Severe respiratory illness outbreak associated with human coronavirus NL63 in a long-term care facility. *Emerging Infectious Diseases*, 24(10), 1964–1966.
- Harvey, R. A. (2007). *Microbiology*. Lippincott Williams & Wilkins.
- Hu, X., Fu, Y., Lu, X., Zhang, Z., Zhang, W., Cao, Y., & Zhang, N. (2017). Protective effects of platycodin D on lipopolysaccharide-induced acute lung injury by activating LXRα-ABCA1 signaling pathway. *Frontiers in Immunology*.
- Hua-Yew Cheng, Yang, C.-M., Lin, T.-C., Lin, L.-T., Chiang, L.-C., & Lin, C.-C. (2011). Excoecarianin, isolated from Phyllanthus urinaria Linnea, inhibits herpes simplex virus type 2 infection through inactivation of viral particles. *Evidence-Based Complementary and Alternative Medicine*, 2011.
- Huang, F., Wang, B.-R., Wu, Y.-Q., Wang, F.-C., Zhang, J., & Wang, Y.-G. (2016). Oncolytic viruses against cancer stem cells: A promising approach for gastrointestinal cancer. *World Journal of Gastroenterology*, 22(35), 7999.
- Huang, Q., Zhang, S., Huang, R., Wei, L., Chen, Y., Lv, S., ... Lin, X. (2013). Isolation and identification of an anti-hepatitis B virus compound from Hydrocotyle sibthorpioides Lam. *Journal of Ethnopharmacology*, 150(2), 568–575.
- Hutchings, A. (1996). Zulu medicinal plants: An *inventory*. University of Natal press.
- Ji, S., Li, Z., Song, W., Wang, Y., Liang, W., Li, K., ... Ye, M. (2016). Bioactive Constituents of Glycyrrhiza uralensis (Licorice): Discovery of the Effective Components of a Traditional Herbal Medicine. *Journal of Natural Products*, 79(2), 281–292.
- Kapewangolo, P., Omolo, J. J., Fonteh, P., Kandawa-Schulz, M., & Meyer, D. (2017).Triterpenoids from ocimum labiatum activates latent HIV-1 expression in vitro:

Potential for use in adjuvant therapy. *Molecules*, 22(10), 1–14.

- A., Mohammadi-Kamalabadi, Karimi, M., Rafieian-Kopaei, М., Amjad, L.. & Salimzadeh, L. (2016). Determination of antioxidant activity, phenolic contents and antiviral potential of methanol extract of Euphorbia spinidens Bornm (Euphorbiaceae). Tropical Journal of Pharmaceutical Research, 15(4), 759–764.
- Karimi, A., Rafieian-Kopaei, M., Moradi, M. T., & Alidadi, S. (2017). Anti–Herpes Simplex Virus Type-1 Activity and Phenolic Content of Crude Ethanol Extract and Four Corresponding Fractions of Quercus brantii L Acorn. Journal of Evidence-Based Complementary and Alternative Medicine, 22(3), 455–461.
- Kim, D. eun, Jung, S., Ryu, H. W., Choi, M., Kang, M., Kang, H., ... Cho, S. (2018).
  Selective oncolytic effect in Epstein-Barr virus (EBV)-associated gastric carcinoma through efficient lytic induction by Euphorbia extracts. *Journal of Functional Foods*, 42(January 2017), 146–158.
- Kim, H., & Song, M.-J. (2011). Analysis and recordings of orally transmitted knowledge about medicinal plants in the southern mountainous region of Korea. *Journal of Ethnopharmacology*, 134(3), 676–696.
- Koenen, E. von. (2001). *Medicinal, poisonous, and edible plants in Namibia.* Klaus hess publishers.
- Kreis, W., Kaplan, M. H., Freeman, J., Sun, D.
  K., & Sarin, P. S. (1990). Inhibition of HIV replication by Hyssop officinalis extracts. *Antiviral Research*, 14(6), 323–337.
- Kudo, E., Taura, M., Matsuda, K., Shimamoto, M., Kariya, R., Goto, H., ... Okada, S. (2013). Inhibition of HIV-1 replication by a tricyclic coumarin GUT-70 in acutely and chronically infected cells. *Bioorganic & Medicinal Chemistry Letters*, 23(3), 606–609.
- Kuiken, C., Leitner, T., Foley, B., Hahn, B., Marx, P., McCutchan, F., ... Korber, B.

Ì

Journal of Integrative and Anatolian Medicine Bütünleyici ve Anadolu Tibbi Dergisi Yayıncı / Publisher Sağlık Bilimleri Üniversitesi University of Health Sciences

(2009). HIV Sequence Compendium 2009. *Theoretical Biology and Biophysics*, 1–436.

- Kulkosky, J., Culnan, D. M., Roman, J., Dornadula, G., Schnell, M., Boyd, M. R., & Pomerantz, R. J. (2001). Prostratin: Activation of latent HIV-1 expression suggests a potential inductive adjuvant therapy for HAART. *Blood*, *98*(10), 3006– 3015.
- Lago, J. H. G., Tezoto, J., Yazbek, P. B., Cassas, F., Santos, J. de F. L., & Rodrigues, E. (2016).
  Exudates used as medicine by the "caboclos river-dwellers" of the Unini River, AM, Brazil – Classification based in their chemical composition. *Brazilian Journal of Pharmacognosy*, 26(3), 379–384.
- Lam, S. K., & Ng, T. B. (2010). Acaconin, a chitinase-like antifungal protein with cytotoxic and anti-HIV-1 reverse transcriptase activities from Acacia confusa seeds. *Acta Biochimica Polonica*, *57*(3), 299–304.
- Lans, C. (2007). Comparison of plants used for skin and stomach problems in Trinidad and Tobago with Asian ethnomedicine. *Journal of Ethnobiology and Ethnomedicine*, *3*.
- Lau, K. M., Lee, K. M., Koon, C. M., Cheung, C. S. F., Lau, C. P., Ho, H. M., ... Fung, K. P. (2008). Immunomodulatory and anti-SARS activities of Houttuynia cordata. *Journal of Ethnopharmacology*, 118(1), 79–85.
- Lewis, W., Day, B. J., & Copeland, W. C. (2003). Mitochondrial toxicity of NRTI antiviral drugs: An integrated cellular perspective. *Nature Reviews Drug Discovery*, 2(10), 812– 822.
- Li, S., Long, C., Liu, F., Lee, S., Guo, Q., Li, R., & Liu, Y. (2006). Herbs for medicinal baths among the traditional Yao communities of China. *Journal of Ethnopharmacology*, *108*(1), 59–67.
- Li, S. Y., Chen, C., Zhang, H. Q., Guo, H. Y., Wang, H., Wang, L., ... Tan, X. (2005). Identification of natural compounds with antiviral activities against SARS-associated

coronavirus. Antiviral Research, 67(1), 18–23.

- Li, T., Liu, L., Wu, H., Chen, S., Zhu, Q., Gao, H., ... Peng, T. (2017). Anti-herpes simplex virus type 1 activity of Houttuynoid A, a flavonoid from Houttuynia cordata Thunb. *Antiviral Research*, 144, 273–280.
- Lin, C. W., Tsai, F. J., Tsai, C. H., Lai, C. C., Wan, L., Ho, T. Y., ... Chao, P. D. L. (2005). Anti-SARS coronavirus 3C-like protease effects of Isatis indigotica root and plantderived phenolic compounds. *Antiviral Research*, 68(1), 36–42.
- Ling, L. jun, Lu, Y., Zhang, Y. yi, Zhu, H. yan, Tu, P., Li, H., & Chen, D. feng. (2020). Flavonoids from Houttuynia cordata attenuate H1N1-induced acute lung injury in mice via inhibition of influenza virus and Toll-like receptor signalling. *Phytomedicine*, 67(December 2019).
- Liu, S. N., Hu, J., Tan, S. H., Wang, Q., Xu, J., Wang, Y., ... Gu, Q. (2017). Ent -Rosane diterpenoids from Euphorbia milii showing an Epstein-Barr virus lytic replication assay. *RSC Advances*, 7(74), 46938–46947.
- Liu, S., Wei, W., Shi, K., Cao, X., Zhou, M., & Liu, Z. (2014). In vitro and in vivo antihepatitis B virus activities of the lignan niranthin isolated from Phyllanthus niruri L. *Journal of Ethnopharmacology*, *155*(2), 1061–1067.
- Locarnini, S. (2004). Molecular virology of hepatitis B virus. In *Seminars in liver disease* (Vol. 24, pp. 3–10). Copyright© 2004 by Thieme Medical Publishers, Inc., 333 Seventh Avenue, New York.
- Lodish, H., Berk, A., Zipursky, S. L., Matsudaira,
  P., Baltimore, D., & Darnell, J. (2000).
  Viruses: Structure, function, and uses. In Molecular Cell Biology. 4th edition. WH Freeman.
- Logan, C. M., & Rice, M. K. (1987). *Medical and scientific abbreviations*. Lippincott.



Yayıncı / Publisher Sağlık Bilimleri Üniversitesi University of Health Sciences

- Lu, H. (2020). Drug treatment options for the 2019-new coronavirus (2019- nCoV). *BioScience Trends.*, *14*(1), 69–71.
- Lubbe, A., Seibert, I., Klimkait, T., & Van Der Kooy, F. (2012). Ethnopharmacology in overdrive: The remarkable anti-HIV activity of Artemisia annua. *Journal of Ethnopharmacology*, 141(3), 854–859.
- MACRAE, W. D., HUDSON, J. B., & TOWERS, G. H. N. (1988). STUDIES ON THE PHARMACOLOGICAL ACTIVITY OF AMAZONIAN EUPHORBIACEAE. Journal of Ethnopharmacology, 22, 143–172.
- Mhlongo, L. S., & Van Wyk, B.-E. (2019). Zulu medicinal ethnobotany: new records from the Amandawe area of KwaZulu-Natal, South Africa. *South African Journal of Botany*, *122*, 266–290.
- Modi, M., Dezzutti, C. S., Kulshreshtha, S., Rawat, A. K. S., Srivastava, S. K., Malhotra, S., ... Gupta, S. K. (2013). Extracts from Acacia catechu suppress HIV-1 replication by inhibiting the activities of the viral protease and Tat. *Virology Journal*, 10, 309.
- Mohan, G., Rao, N., & Pragada, P. M. (2014). Survey and documentation of some important medicinal applications of Mangrove plants of Andhra Pradesh, India. *The Journal of Ethnobiology and Traditional Medicine*, (March).
- Mosaddegh, M., Naghibi, F., Moazzeni, H., Pirani, A., & Esmaeili, S. (2012).
  Ethnobotanical survey of herbal remedies traditionally used in Kohghiluyeh va Boyer Ahmad province of Iran. *Journal of Ethnopharmacology*, 141(1), 80–95.
- Mukhtar, M., Arshad, M., Ahmad, M.,
  Pomerantz, R. J., Wigdahl, B., & Parveen, Z.
  (2008). Antiviral potentials of medicinal plants. *Virus Research*, 131(2), 111–120.
- Musarra-Pizzo, M., Ginestra, G., Smeriglio, A., Pennisi, R., Sciortino, M. T., & Mandalari, G. (2019). The Antimicrobial and Antiviral Activity of Polyphenols from Almond

(Prunus dulcis L.) Skin. *Nutrients*, 11(10), 1–11.

- Mustafa, B., Hajdari, A., Pulaj, B., Quave, C. L., & Pieroni, A. (2020). Medical and food ethnobotany among Albanians and Serbs living in the Shtërpcë/Štrpce area, South Kosovo. *Journal of Herbal Medicine*, 100344.
- Naithani, R., Huma, L. C., Holland, L. E., Shukla,
  D., McCormick, D. L., Mehta, R. G., &
  Moriarty, R. M. (2008). Antiviral activity of
  phytochemicals: a comprehensive review. *Mini Reviews in Medicinal Chemistry*, 8(11),
  1106–1133.
- Nomura, T., Fukushi, M., Oda, K., Higashiura, A., Irie, T., & Sakaguchi, T. (2019). Effects of Traditional Kampo Drugs and Their Constituent Crude Drugs on Influenza Virus Replication in Vitro: Suppression of Viral Protein Synthesis by Glycyrrhizae Radix. *Evidence-Based Complementary and Alternative Medicine*, 2019.
- Panyadee, P., Balslev, H., Wangpakapattanawong, P., & Inta, A. (2019).
  Medicinal plants in homegardens of four ethnic groups in Thailand. *Journal of Ethnopharmacology*, 239(April), 111927.
- Parada, M., Carrió, E., Bonet, M. À., & Vallès, J. (2009). Ethnobotany of the Alt Emporda region (Catalonia, Iberian Peninsula): plants used in human traditional medicine. *Journal* of Ethnopharmacology, 124(3), 609–618.
- Parker Jr, F., & Nye, R. N. (1925). Studies on filterable viruses: II. Cultivation of herpes virus. *The American Journal of Pathology*, *1*(3), 337.
- Parvez, M. K., Al-Dosari, M. S., Alam, P., Rehman, M. T., Alajmi, M. F., & Alqahtani, A. S. (2019). The anti-hepatitis B virus therapeutic potential of anthraquinones derived from Aloe vera. *Phytotherapy Research*, 33(11), 2960–2970.
- Pieroni, A., Quave, C. L., & Santoro, R. F. (2004). Folk pharmaceutical knowledge in the territory of the Dolomiti Lucane, inland

Yayıncı / Publisher Sağlık Bilimleri Üniversitesi University of Health Sciences

southern Italy. *Journal of Ethnopharmacology*, 95(2–3), 373–384.

- Porter, R. S., & Bode, R. F. (2017). A Review of the Antiviral Properties of Black Elder (Sambucus nigra L.) Products. *Phytotherapy Research*, 31(4), 533–554.
- Powell, M. K., Benková, K., Selinger, P., Dogo'i, M., Luňáčková, I. K., Koutníková, H., ... Heneberg, P. (2016). Opportunistic infections in HIV-infected patients differ strongly in frequencies and spectra between patients with low CD4+ cell counts examined postmortem and compensated patients examined antemortem irrespective of the HAART Era. *PLoS ONE*, 11(9), 1–19.
- Qi, W., Hua, L., & Gao, K. (2014). Chemical Constituents of the Plants from the Genus Phyllanthus. *Chemistry and Biodiversity*, *11*(2), 181–196.
- Rajasekaran, D., Palombo, E. A., Yeo, T. C., Ley, D. L. S., Tu, C. L., Malherbe, F., & Grollo, L. (2013). Identification of traditional medicinal plant extracts with novel anti-influenza activity. *PLoS ONE*, 8(11), 1–15.
- Reddy, V. L. N., Reddy, S. M., Ravikanth, V., Krishnaiah, P., Goud, T. V., Rao, T. P., ... Venkateswarlu, Y. (2005). A new bisandrographolide ether from Andrographis paniculata nees and evaluation of anti-HIV activity. *Natural Product Research*, 19(3), 223–230.
- Redeploying plant defences. (2020). Nature Plants, 6(3), 177.
- Reichling, J., Neuner, A., Sharaf, M., Harkenthal, M., & Schnitzler, P. (2009). Antiviral activity of Rhus aromatica (fragrant sumac) extract against two types of herpes simplex viruses in cell culture. *Pharmazie*, 64(8), 538–541.
- Ribeiro, R. V., Bieski, I. G. C., Balogun, S. O., & Martins, D. T. de O. (2017). Ethnobotanical study of medicinal plants used by Ribeirinhos in the North Araguaia microregion, Mato Grosso, Brazil. *Journal of Ethnopharmacology*, 205(December 2016), 69–102.

- of Rigat, M., Bonet, M. À., Garcia, S., Garnatje, T., & Vallès, J. (2007). Studies on pharmaceutical ethnobotany in the high river Ter valley (Pyrenees, Catalonia, Iberian py Peninsula). Journal of Ethnopharmacology, 113(2), 267–277.
  - Rivera, D., Allkin, R., Obón, C., Alcaraz, F., Verpoorte, R., & Heinrich, M. (2014). What is in a name? the need for accurate scientific nomenclature for plants. *Journal of Ethnopharmacology*, *152*(3), 393–402.
  - Roumy, V., Ruiz, L., Ruiz Macedo, J. C., Gutierrez-Choquevilca, A. L., Samaillie, J., Encinas, L. A., ... Hennebelle, T. (2020).
    Viral hepatitis in the Peruvian Amazon: Ethnomedical context and phytomedical resource. *Journal of Ethnopharmacology*, 255(December 2019), 112735.
  - Rullas, J., Bermejo, M., García-Pérez, J., Beltrán,
    M., González, N., Hezareh, M., ... Alcamí, J.
    (2004). Prostratin induces HIV activation and downregulates HIV receptors in peripheral blood lymphocytes. *Antiviral Therapy*, 9(4), 545–554.
  - Ryu, W.-S. (2016). *Molecular virology of human* pathogenic viruses. Academic Press.
  - Ryu, Y. B., Jeong, H. J., Kim, J. H., Kim, Y. M., Park, J. Y., Kim, D., ... Lee, W. S. (2010).
    Biflavonoids from Torreya nucifera displaying SARS-CoV 3CLpro inhibition. *Bioorganic and Medicinal Chemistry*, 18(22), 7940–7947.
  - Safa, O., Soltanipoor, M. A., Rastegar, S., Kazemi, M., Nourbakhsh Dehkordi, K., & Ghannadi, A. (2013). An ethnobotanical survey on hormozgan province, Iran. *Avicenna Journal of Phytomedicine*, 3(1), 64–81.
  - Salehi, B., Anil Kumar, N. V., Şener, B., Sharifi-Rad, M., Kılıç, M., Mahady, G. B., ... Sharifi-Rad, J. (2018). Medicinal plants used in the treatment of human immunodeficiency virus. *International Journal of Molecular Sciences*, 19(5).



- Salguero, C. P. (2003). A compendium of traditional Thai herbal medicine. In *A Thai Herbal-traditional Recipes for Health and Harmony*. Findhorn Press Scotland.
   Sheu, T. G., Deyde, V. M., Okomo-Adhiambo, M., Garten, R. J., Xu, X., Bright, R. A., ... Gubareva, L. V. (2008). Surveillance for neuraminidase inhibitor resistance among
- Sánchez-Machado, D. I., López-Cervantes, J., Sendón, R., & Sanches-Silva, A. (2017). Aloe vera: Ancient knowledge with new frontiers. *Trends in Food Science and Technology*, 61, 94–102.
- Savić, J., Mačukanović-Jocić, M., & Jarić, S. (2019). Medical ethnobotany on the Javor Mountain (Bosnia and Herzegovina). *European Journal of Integrative Medicine*, 27(October 2018), 52–64.
- Schaeffer, H. J., Gurwara, S., Vince, R., & Bittner, S. (1971). Novel substrate of adenosine deaminase. *Journal of Medicinal Chemistry*, 14(4), 367–369.
- Scherrer, A. M., Motti, R., & Weckerle, C. S. (2005). Traditional plant use in the areas of Monte Vesole and Ascea, Cilento National Park (Campania, Southern Italy). *Journal of Ethnopharmacology*, 97(1), 129–143.
- Schillie, S. F., Murphy, T. V, Sawyer, M., Ly, K., Hughes, E., Jiles, R., ... Ward, J. W. (2013).CDC guidance for evaluating health-care personnel for hepatitis B virus protection and for administering postexposure management.
- Schoeman, D., & Fielding, B. C. (2019). Coronavirus envelope protein: Current knowledge. *Virology Journal*, 16(1), 1–22.
- Shah, A., Bharati, K. A., Ahmad, J., & Sharma, M. P. (2015). New ethnomedicinal claims from Gujjar and Bakerwals tribes of Rajouri and Poonch districts of Jammu and Kashmir, India. *Journal of Ethnopharmacology*, *166*, 119–128.
- Shamsabadipour, S., Ghanadian, M., Saeedi, H., Reza Rahimnejad, M., Mohammadi-Kamalabadi, M., Ayatollahi, S. M., & Salimzadeh, L. (2013). Triterpenes and steroids from euphorbia denticulata lam. with anti-herpes symplex virus activity. *Iranian Journal of Pharmaceutical Research*, 12(4), 759–767.

- Sheu, T. G., Deyde, V. M., Okomo-Adhiambo, M., Garten, R. J., Xu, X., Bright, R. A., ... Gubareva, L. V. (2008). Surveillance for neuraminidase inhibitor resistance among human influenza A and B viruses circulating worldwide from 2004 to 2008. *Antimicrobial Agents and Chemotherapy*, 52(9), 3284– 3292.
- Sriwilaijaroen, N., Fukumoto, S., Kumagai, K., Hiramatsu, H., Odagiri, T., Tashiro, M., & Suzuki, Y. (2012). Antiviral effects of Psidium guajava Linn.(guava) tea on the growth of clinical isolated H1N1 viruses: Its role in viral hemagglutination and neuraminidase inhibition. *Antiviral Research*, 94(2), 139–146.
- Stiver, G. (2003). The treatment of influenza with antiviral drugs. *Cmaj*, *168*(1), 49–57.
- Suárez, M. E. (2019). Medicines in the forest: Ethnobotany of wild medicinal plants in the pharmacopeia of the Wichí people of Salta province (Argentina). *Journal of Ethnopharmacology*, 231, 525–544.
- Suroowan, S., Pynee, K. B., & Mahomoodally, M. F. (2019). A comprehensive review of ethnopharmacologically important medicinal plant species from Mauritius. *South African Journal of Botany*.
- Thakur, M., Asrani, R. K., Thakur, S., Sharma, P. K., Patil, R. D., Lal, B., & Parkash, O. (2016).
  Observations on traditional usage of ethnomedicinal plants in humans and animals of Kangra and Chamba districts of Himachal Pradesh in North-Western Himalaya, India. *Journal of Ethnopharmacology*, *191*(June 2016), 280–300.
- Thompson, C. J. S. (1928). The History and Lore of Cinchona. *British Medical Journal*, 1188– 1190.
- Treml, J., Gazdová, M., Šmejkal, K., Šudomová, M., Kubatka, P., & Hassan, S. T. S. (2020).
  Natural products-derived chemicals: Breaking barriers to novel anti-HSV drug development. *Viruses*, 12(2).



Yayıncı / Publisher Sağlık Bilimleri Üniversitesi University of Health Sciences

- TSUMURA & CO. (2016). About Kampo. Retrieved March 25, 2020, from https://www.tsumura.co.jp/english/kampo/
- Urdang, G. (1945). The legend of Cinchona. *The Scientific Monthly*, *61*(1), 17–20.
- Vlietinck, A. J., Van Hoof, L., Totté, J., Lasure,
  A., Berghe, D. Vanden, Rwangabo, P. C., &
  Mvukiyumwami, J. (1995). Screening of
  hundred Rwandese medicinal plants for
  antimicrobial and antiviral properties. *Journal of Ethnopharmacology*, 46(1), 31–47.
- Wagner, R. R., & Krug, R. M. (2020). Virus. In *Encyclopædia Britannica*. Encyclopædia Britannica, inc. Retrieved from https://www.britannica.com/science/virus
- Wang, B., Wei, Y., Zhao, X., Tian, X., Ning, J., Zhang, B., ... Wang, C. (2018). Unusual entatisane type diterpenoids with 2-oxopropyl skeleton from the roots of Euphorbia ebracteolata and their antiviral activity against human rhinovirus 3 and enterovirus 71. *Bioorganic Chemistry*, 81(August), 234– 240.
- Wang, M., Cao, R., Zhang, L., Yang, X., Liu, J., Xu, M., ... Xiao, G. (2020). Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. *Cell Research*, 30(3), 269–271.
- Waterson, A. P., & Wilkinson, L. (1978). An *introduction to the history of virology*.
  Cambridge University Press, Bentley House, 200 Euston Road, London NW1 2DB.
- Wei, W., Pan, Y., Chen, Y., Lin, C., Wei, T., & Zhao, S. (2005). Carboxylic acids from Phyllanthus urinaria. *Chemistry of Natural Compounds*, 41(1), 17–21.
- Weiss, R. A. (1993). How does HIV cause AIDS? *Science*, *260*(5112), 1273–1279.
- Weng, J. R., Lin, C. S., Lai, H. C., Lin, Y. P., Wang, C. Y., Tsai, Y. C., ... Lin, C. W. (2019). Antiviral activity of Sambucus FormosanaNakai ethanol extract and related phenolic acid constituents against human

coronavirus NL63. Virus Research, 273(September), 197767.

- Wermig-morgan, J. H. (2020). Elderberry is antibacterial, anti-viral and modulates the immune system :, (February).
- WHO. (2014). Smallpox. Retrieved March 25, 2020, from https://www.who.int/biologicals/vaccines/sm allpox/en/
- WHO. (2017). Herpes Simplex Virus. Retrieved March 25, 2020, from https://www.who.int/news-room/factsheets/detail/herpes-simplex-virus
- WHO. (2019a). Hepatitis B. Retrieved March 25, 2020, from https://www.who.int/newsroom/fact-sheets/detail/hepatitis-b
- WHO. (2019b). HIV/AIDS. Retrieved March 25, 2020, from https://www.who.int/newsroom/fact-sheets/detail/hiv-aids
- Willcox, M. L., Burton, S., Oyweka, R., Namyalo, R., Challand, S., & Lindsey, K. (2011). Evaluation and pharmacovigilance of projects promoting cultivation and local use of Artemisia annua for malaria. *Malaria Journal*, 10(April).
- Won, J. N., Lee, S. Y., Song, D. S., & Poo, H. Y. (2013). Antiviral activity of the plant extracts from Thuja orientalis, Aster spathulifolius, and Pinus thunbergii against influenza virus A/PR/8/34. Journal of Microbiology and Biotechnology, 23(1), 125–130.
- Wu, S., Patel, K. B., Booth, L. J., Metcalf, J. P., Lin, H. K., & Wu, W. (2010). Protective essential oil attenuates influenza virus infection: An in vitro study in MDCK cells. *BMC Complementary and Alternative Medicine*, 10.
- Wyles, D. L. (2013). Antiviral resistance and the future landscape of hepatitis C virus infection therapy. *Journal of Infectious Diseases*, 207(SUPPL.1), 33–39.
- Xi, S., & Gong, Y. (2017). Essentials of Chinese Materia Medica and Medical Formulas: New Century Traditional Chinese Medicine. Academic Press.



Yayıncı / Publisher Sağlık Bilimleri Üniversitesi University of Health Sciences

- Yang, C.-M., Cheng, H.-Y., Lin, T.-C., Chiang, L.-C., & Lin, C.-C. (2005). Acetone, ethanol and methanol extracts of Phyllanthus urinaria inhibit HSV-2 infection in vitro. *Antiviral Research*, 67(1), 24–30.
- Yang, C. M., Cheng, H. Y., Lin, T. C., Chiang, L. C., & Lin, C. C. (2007). The in vitro activity of geraniin and 1,3,4,6-tetra-O-galloyl-β-dglucose isolated from Phyllanthus urinaria against herpes simplex virus type 1 and type 2 infection. *Journal of Ethnopharmacology*, *110*(3), 555–558.
- Yaseen, G., Ahmad, M., Sultana, S., Suleiman Alharrasi, A., Hussain, J., Zafar, M., & Shafiq-Ur-Rehman. (2015). Ethnobotany of medicinal plants in the Thar Desert (Sindh) of Pakistan. *Journal of Ethnopharmacology*, 163, 43–59.
- Yu, M. S., Lee, J., Lee, J. M., Kim, Y., Chin, Y. W., Jee, J. G., ... Jeong, Y. J. (2012). Identification of myricetin and scutellarein as novel chemical inhibitors of the SARS coronavirus helicase, nsP13. *Bioorganic and Medicinal Chemistry Letters*, 22(12), 4049–4054.
- Zhao, Y., Geng, C. A., Sun, C. L., Ma, Y. B., Huang, X. Y., Cao, T. W., ... Chen, J. J. (2014). Polyacetylenes and anti-hepatitis B virus active constituents from Artemisia capillaris. *Fitoterapia*, 95, 187–193.
- Zheng, X. long, & Xing, F. wu. (2009). Ethnobotanical study on medicinal plants around Mt.Yinggeling, Hainan Island, China. *Journal of Ethnopharmacology*, 124(2), 197– 210.