

PAPER DETAILS

TITLE: On the Subsheaves of the Sheaf of the Fundamental Groups

AUTHORS: Murat ÖZDEMİR, Mustafa BAYRAKTAR

PAGES: 15-21

ORIGINAL PDF URL: <https://dergipark.org.tr/tr/download/article-file/252471>

ON THE SUBSHEAVES OF THE SHEAF OF THE FUNDAMENTAL GROUPS

Murat ÖZDEMİR ve Mustafa BAYRAKTAR

Atatürk University, Faculty of Arts and Sciences, Department of Mathematics
25240 Erzurum-Turkey

ABSTRACT

Let X be a connected and locally arcwise connected topological space and H be the sheaf of fundamental groups on X . In this paper, any two subsheaves of H is constructed and it is proven that if stalks of these subsheaves are conjugate subgroups for every $x \in X$, then subsheaves are homomorphic.

ÖZET

X bağlantılı ve lokal eğrisel bağlantılı bir topolojik uzay ve H , X üzerinde esas grupların demeti olsun. Bu makalede, H nin herhangi iki alt demeti teşkil edildi ve her $x \in X$ için bu alt demetlerin sapları eşlenik altgruplar ise alt demetlerin homomorf olduğu ispatlandı.

1. INTRODUCTION

Definition 1. A sheaf of groups on X is a pair (H, π) where

- i) H is a topological space.
- ii) $\pi : H \rightarrow X$ is a local homeomorphism onto X .
- iii) Each $H_x = \pi^{-1}(x)$, for $x \in X$, is a group (and is called the stalk of H at x).
- iv) The group operations are continuous.

Let X be a connected and locally arcwise connected topological space and H_x be the fundamental group of X based any $x \in X$, that is, $H_x =$

$\pi_1(X, x)$. Let $X = (X, c)$ be a pointed topological spaces, for an arbitrary fixed point $c \in X$. Let us denote by H the disjoint union of the fundamental groups obtained for each $x \in X$. i.e., $H = \bigvee_{x \in X} H_x$. Also H is a set over X and the mapping $\phi : H \rightarrow X$ defined by $\phi(\sigma_x) = ([\alpha]_x) = x$, for any $\sigma_x = [\alpha]_x \in H_x \subset H$, is onto.

We introduce a topology on H as follows : Let H_c be the fundamental group of X with respect to c , $x_0 \in X$ be an arbitrary fixed point, $W = W(x_0)$ be an arcwise connected open neighborhood of x_0 and $\sigma_c = [\alpha]_c \in H_c$ be any point of H . Let us define a mapping $s : W \rightarrow H$ such that $s(x) = [(\gamma^{-1}\alpha)\gamma]_x$ for every $x \in W$, where $\gamma \in [\gamma]$ is an arc with initial point c and terminal point x . $[\gamma]$ determines s mapping between H_c and H_x . Let us choose the homotopy class $[\gamma]$ arbitrarily fixed for each $x \in W$. Thus, $s = s(\sigma_c)$ and s a well-defined mapping from W to H such that $\phi \circ s = I_W$ [1]. Let us denote the totality of the mappings s defined on W by $\Gamma(W, H)$.

If B is a base for X , then $B^* = \{s(W) : W \in B, s \in \Gamma(W, H)\}$ is a base for H . In this topology, the mappings ϕ and s are continuous and ϕ is a locally topological mapping. Then (H, ϕ) is a sheaf over X . (H, ϕ) (or only H) is called "The Sheaf of the Fundamental Groups" over X [1]. For any open set $W \subset X$, an element s of $\Gamma(W, H)$ is called a section of the sheaf H over W . The set $\Gamma(W, H)$ is a group with the pointwise operation of multiplication. Thus, H is a sheaf of the groups over X [2]. Furthermore, the $H_x = \pi_1(X, x)$ is called the stalk of the sheaf H for any $x \in X$.

2. CHARACTERISTIC FEATURES OF H

i) Let $W \subset X$ be an open set. Then, any section over W can be extended to a global section over X [3]. Furthermore,

$$\Gamma(W, H) = \{s|_W : s \in \Gamma(X, H)\} = \Gamma(X, H)|_W$$

ii) Any two stalks of H are isomorphic with each other [3].

iii) Let $W_1, W_2 \subset X$ be any two open sets, $s_1 \in \Gamma(W_1, H)$ and $s_2 \in \Gamma(W_2, H)$. If $s_1(x_0) = s_2(x_0)$ for any point $x_0 \in W_1 \cap W_2$, then $s_1 = s_2$ over the whole $W_1 \cap W_2$ [5].

iv) Let $W \subset X$ be an open set and $s_1, s_2 \in \Gamma(W, H)$. If $s_1(x_0) = s_2(x_0)$ for any point $x_0 \in W$, then $s_1 = s_2$ over the whole W [5].

v) To each point $\sigma_x \in H_x \subset H$, there is uniquely corresponds a section $s \in \Gamma(W, H)$ such that $s(x) = \sigma_x$. Hence $H_x \cong \Gamma(W, H)$. In particular, $H_x \cong \Gamma(X, H)$ [2].

vi) Let $x \in X$ be any point and $W = W(x)$ be any open set. Then, $\pi^{-1}(W) = \bigcup_{i \in I} s_i(W)$, $s_i \in \Gamma(W, H)$ and $\pi|_{s_i(W)} : s_i(W) \rightarrow W$ is a topological mapping for every $i \in I$. Hence, $W = W(x)$ is evenly covered by π . Then, π is a cover projection and (H, π) is a covering space of X . Moreover (H, π) is regular, because the group T of cover transformations of H is isomorphic to the group H_x , that is, T is transitive on H_x [4].

Definition 2. Let (H, π) be the sheaf of fundamental groups over X and $H' \subset H$ be an open set. Then H' is called a subsheaf of groups, if:

i) $\pi(H') = X$

ii) For each point $x \in X$, the stalk H'_x is subgroup of H_x .

Let $H' \subset H$ be a subsheaf of groups and $W \subset X$ be an open set. Then, the set $\Gamma(W, H') \subset \Gamma(W, H)$ is a subgroup. In particular, if we take $W = X$, then $\Gamma(X, H') \subset \Gamma(X, H)$. Conversely, let us suppose that $\Gamma(X, H)$ is the group of global sections of H over X and $D \subset \Gamma(X, H)$ be a subgroup. Then, the set $\{s_i(x) : s_i \in D\}$ is a subgroup of H_x over X for each $x \in X$. Let us denote $\{s_i(x) : s_i \in D\}$ by H'_x . Then $H' = \bigcup_{x \in X} H'_x$ is a set over X with the natural projection $\pi' = \pi|_{H'}$ and $D = \Gamma(W, H')$. One can show that (H', π') is a subsheaf of groups [1].

If $H', H'' \subset H$ be any two subsheaves of groups, then H'_x and H''_x are two subgroups of H_x for each $x \in X$ by definition 2. Moreover,

subsheaves H' and H'' have a relative topology. If $s \in \Gamma(W_1, H)$ for every $x \in X$ and $W_1 = W_1(x) \subset X$, then whenever $H' \subset H$ and $H'' \subset H$ are open subsets, the sets $s(W_1) \cap H' = s'(W)$ and $s(W_1) \cap H'' = s''(W)$ are open in H' and H'' , respectively. Therefore the sets

$$B^*_1 = \{s'(W) : W = W(x)\}$$

and

$$B^*_2 = \{s''(W) : W = W(x)\}$$

are base for a topology on H' and H'' , respectively.

Definition 3. Let $(H_1, \pi_1), (H_2, \pi_2)$ be any two sheaf on X and $\phi : H_1 \rightarrow H_2$ be a mapping.

i) The mapping $\phi : H_1 \rightarrow H_2$ is called stalk preserving, if $\pi_2 \circ \phi = \pi_1$ (therefore, $\phi(H_1)_x \subset (H_2)_x$ for all $x \in X$)

ii) Let $\phi : H_1 \rightarrow H_2$ be a stalk preserving and continuous mapping. Then the mapping ϕ is called a sheaf morphism between the sheaves H_1 and H_2 .

iii) Let $\phi : H_1 \rightarrow H_2$ be a sheaf morphism. If ϕ is a homomorphism on each stalk, then it is called a sheaf homomorphism between the sheaves H_1 and H_2 .


iv) Let $\phi : H_1 \rightarrow H_2$ be a sheaf homomorphism. If ϕ is a homeomorphism, then it is called a sheaf isomorphism between the sheaves H_1 and H_2 .

Theorem 4. Let X be a connected and locally arcwise connected topological space, (H, π) be the sheaf of the fundamental groups over X , (H', π_1) and (H'', π_2) be any two subsheaves of H . If H'_x and H''_x are conjugate subgroups of H_x with an element $\sigma = [\delta]_x$ for every $x \in X$, i.e., $H''_x = [\delta]_x H'_x [\delta]^{-1}_x$

then, the sheaves H' and H'' are homomorphic, where $\pi_1 = \pi|_{H'}$ and $\pi_2 = \pi|_{H''}$.

Proof. Let $x \in X$ be an arbitrarily fixed point and β_1, β_2 be any two closed path based at x . If $\beta_1 \sim \beta_2$, then $(\delta\beta_1)\delta^{-1} \sim (\delta\beta_2)\delta^{-1}$. Thus correspondence

$[\beta]_x \rightarrow [(\delta\beta)\delta^{-1}]_x$ is well-defined. Since the point $x \in X$ is arbitrarily fixed, the above correspondence gives us a map $\varphi : H' \rightarrow H''$ such that $\varphi([\beta]_x) = ([\delta]_x[\beta]_x) [\delta]^{-1}_x$ for every $[\beta] \in H'$.



i) φ **preserves the stalks**. In fact, for arbitrarily fixed point $x \in X$ and any $[\beta]_x \in H'$

$$\begin{aligned}
 \pi_2 \circ \varphi([\beta]_x) &= \pi_2(\varphi([\beta]_x)) \\
 &= \pi_2([(\delta]_x[\beta]_x)[\delta]^{-1}_x]) \\
 &= \pi_2([\delta\beta]_x[\delta]^{-1}_x) \\
 &= \pi_2([\delta\beta)\delta^{-1}]_x) \\
 &= \pi_2([\rho]_x) \\
 &= x \\
 &= \pi_1([\beta]_x).
 \end{aligned}$$

Since the point $x \in X$ is arbitrarily fixed, we obtain $\pi_2 \circ \varphi = \pi_1$.

i) φ is **continuous**. Let us show that if $U_2 \subset H''$ is any open set, then $\varphi^{-1}(U_2) = U_1 \subset H'$ is an open set. Without loss of generality, we assume that $U_2 = s''(W)$, where $W \subset X$ is an open set and $s'' \in \Gamma(W, H'')$. Thus, $\pi_2(U_2) = \pi_2(s''(W)) = W$. Now let $\sigma_2 = [\rho]_x \in U_2$ be an element. Then, there exists at least one element $\sigma_1 = [\beta]_x \in U_1 = \varphi^{-1}(U_2)$ such that $\varphi(\sigma_1) = \sigma_2$. Since $\pi_1(\sigma_1) = \pi_1([\beta]_x) = x$, there is a section $s' \in \Gamma(W, H')$ such that $s'(x) = [\beta]_x = \sigma_1$ and $s'(W) \subset H'$ is an open. Also $s'(W) \subset U_1$. It is easily seen that $U_1 = \bigcup_{i \in I} s'_i(W_i)$. Therefore, $U_1 \subset H'$ is an open set, that is, φ is a sheaf morphism.

iii) φ a **sheaf homomorphism**. For every $x \in X$ the map $\varphi|_{(H')_x} : (H')_x \rightarrow (H'')_x$ is homomorphism. In fact, if β_1, β_2 are two arcs at $x \in X$ then $(\delta\beta_1)\delta$, $(\delta\beta_2)\delta$ are two arcs at $x \in X$. Hence,

$$\begin{aligned}\varphi([\beta_1]_x) \varphi([\beta_2]_x) &= (([\delta][\beta_1][\delta]^{-1}) ([\delta][\beta_2][\delta]^{-1})) \\ &= ([\delta][\beta_1]) ([\beta_2][\delta]^{-1}) \\ &= [\delta]([\beta_1] [\beta_2])[\delta]^{-1} \\ &= [\delta][\beta_1\beta_2][\delta]^{-1} \\ &= \varphi([\beta_1\beta_2]_x).\end{aligned}$$

Theorem 5. Let X be a connected and locally arcwise connected topological space, (H, \cdot) be the sheaf of the fundamental groups over X , (H', \cdot_1) and (H'', \cdot_2) be any two subsheaves of H . If subsheaves H' and H'' are homomorphic, then the map $\varphi_* : \Gamma(W, H') \rightarrow \Gamma(W, H'')$ defined by $\varphi_*(s') = \varphi \circ s'$ is a group homomorphism for any $W \subset X$.

Proof. Let s'_1 ve $s'_2 \in \Gamma(W, H')$. For every $x \in W$,

$$\begin{aligned}(\varphi \circ (s'_1.s'_2))(x) &= \varphi((s'_1.s'_2)(x)) \\ &= \varphi(s'_1(x).s'_2(x)) \\ &= \varphi([\beta_1]_x[\beta_2]_x) \\ &= \varphi([\beta_1\beta_2]_x) \\ &= ([\delta][\beta_1\beta_2][\delta]^{-1}) \\ &= ([\delta]([\beta_1][\beta_2]))[\delta]^{-1} \\ &= ([\delta]([\beta_1][\delta]^{-1}[\delta][\beta_2]))[\delta]^{-1} \\ &= (([\delta][\beta_1][\delta]^{-1})([\delta][\beta_2][\delta]^{-1})) \\ &= \varphi([\beta_1]_x) \varphi([\beta_2]_x) \\ &= \varphi(s'_1(x)) \varphi(s'_2(x)) \\ &= (\varphi \circ s'_1)(x) (\varphi \circ s'_2)(x).\end{aligned}$$

Thus, $\varphi_*(s'_1.s'_2) = \varphi_*(s'_1) \varphi_*(s'_2)$.

REFERENCES

- [1] Balcı, S., "On the Restricted Sheaf", Comm. Fac. Sci. Univ. Ankara, Ser. A1. Vol. 37., 41-51 (1988)
- [2] Balcı, S., "On the Sheaf of the Homology Groups of the Complex Manifold", Jour. Inst. Math. Comp. Sci. Math. Ser., India., Vol. 1, No. 2., 127-133 (1988)
- [3] Saunders, M.L. and Leke, Moerdijk., "Sheaves in Geometry and Logic", Springer-Verlag, New York, 1992.
- [4] Uluçay, C., "On Homology Covering Spaces and Sheaf Associated to the homology group", Comm. Fac. Sci. Univ. Ankara. Ser. A1: Math., Vol. 33., 22-28 (1984)
- [5] Verschoren, A., "Relative Invariants of Sheaves", Marcel Dekker, Inc., New York, 1987.