PAPER DETAILS

TITLE: Süt Ineklerinin Beslenmesinde Nisastanin Önemi ve Düsük Nisastali Rasyonlarla Besleme

Stratejileri

AUTHORS: M USTA, P SAÇAKLI

PAGES: 0-0

ORIGINAL PDF URL: https://dergipark.org.tr/tr/download/article-file/66228



ERCİYES ÜNİVERSİTESİ VETERİNER FAKÜLTESİ DERGİSİ

Journal of Faculty of Veterinary Medicine, Erciyes University

Araştırma Makalesi / Research Article 10(2), 101-105, 2013

Investigation of Bovine Viral Diarrhoea Virus in Dairy Cattle Premises Where Aborts Occur^{*}

Metin GÜRÇAY¹, Mustafa İSSİ², Yusuf GÜL²

¹ Republic of Turkey, Ministry of Food Agriculture and Livestock, Veterinary Control and Research Institute, Elazığ-TURKEY ² University of Firat, Faculty of Veterinary Medicine, Department of Internal Disease, Elazığ-TURKEY

Summary: In this study, the presence of Bovine Viral Diarrhoea Virus (BVDV) was investigated by serologic and virologic methods in dairy cattle enterprises where aborts occur. Study materials include 602 blood samples taken from aborted cows and 241 tissue samples taken from waste fetus samples (abortion/stillborn fetus) admitted to the Control Institute of Veterinary Medicine (Elazığ) from dairy cattle enterprises where abort cases were seen in Elazığ, Malatya and Bingöl between September 2011 and April 2012. The lung, liver, spleen and brain tissue samples taken from aborted fetus and blood serum samples were tested by ELISA. With regard to BVDV antibodies, the serum blood samples were founded to be positive in 70.76% (426/602) in total. This ratio was determined as 76.11% (137/180), 56.25% (18/32), 69.48% (271/390) in Elazığ, Malatya and Bingöl provinces, respectively. The prevalence of BVDV antigen in the tissue samples of aborted fetuses was detected as 28.63% (69/241) in total, 25.75% (17/66) in Elazığ, 36.36% (4/11) in Malatya and 29.26% (48/164) in Bingöl.

As a result, in aborted dairy cows in Elazığ, Malatya and Bingöl, antigens and antibodies have been reported to highseroprevalance. It was noted that in order to reduce economic losses due to this virus, there should be emphasis on preventive measures.

Key Words: Abortion, BVDV, dairy cattle, ELISA

Abort Olgularının Görüldüğü Süt Sığırı İşletmelerinde Bovine Viral Diarrhoea Virusun Araştırılması

Özet: Bu çalışmada, yavru atma olaylarının görüldüğü süt sığırı işletmelerinde Bovine Viral Diarrhoea Virus (BVDV)'un varlığı serolojik ve virolojik olarak araştırıldı. Çalışma materyalini, Eylül 2011 ile Nisan 2012 ayları arasında Elazığ, Malatya ve Bingöl illerinde yavru atma olaylarının görüldüğü süt sığırı işletmelerinden Elazığ Veteriner Kontrol Enstitüsüne gönderilen 602 adet inek kan serumu ile 241 adet atık numunesinden (abort/ölü doğmuş fötüsden) alınan doku örnekleri oluşturdu. Fötüse ait akciğer, karaciğer, dalak ve beyin doku örneklerinden hazırlanan doku ekstraktları ile kan serumu örnekleri oluşturdu. Fötüse ait akciğer, karaciğer, dalak ve beyin doku örneklerinden hazırlanan doku ekstraktları ile kan serumu örnekleri ELISA testi ile incelendi. Kan serumu örneklerinde BVDV antikorları yönünden genel toplamda %70.76 (426/602) belirlenirken Elazığ ilinde %76.11 (137/180), Malatya ilinde %56.25 (18/32) ve Bingöl ilinde %69.48 (271/390) oranında pozitiflik tespit edildi. Atık numune örneklerinin ise toplamda ve illerde sırasıyla %28.63 (69/241), %25.75 (17/66), %36.36 (4/11) ve %29.26 (48/164) oranında BVDV antijenleri yönünden pozitif olduğu belirlendi. Sonuç olarak Elazığ, Malatya ve Bingöl bölgesinde atık yapmış süt ineklerinde BVDV antijen ve antikorlarının yüksek seroprevalansa sahip olduğu ve bu virusa bağılı ekonomik kayıpların önlenmesi için koruyucu tedbirlere önem verilmesi gerektiği düşünülmektedir.

Anahtar Kelimeler: BVDV, ELİSA, süt sığırı, yavru atma

Introduction

Included in pestivirus genus of Flaviridiae family, Bovine Viral Diarrhoea Virus (BVDV) infections are observed worldwide as well as in our country and cause significant economic losses (1, 18, 20). Virus with genotype of BVDV-1 and BVDV-2 has two

Geliş Tarihi/Submission Date : 03.01.2013 Kabul Tarihi/Accepted Date : 13.05.2013 biotypes according to their cytopathic effects on cell cultures namely; cytopathogenic (cp-BVDV) and noncytopathogenic (ncp-BVDV) biotypes (16, 18, 20).

In cattle with BVDV, bovine viral diarrhea, acute or chronic mucosal disease, congenital anomalies or persistent infected calves (virus emitting), abortion and calves with low chance of survival due to transplacental infections have been reported in various clinical manifestations (1, 18).

In this study, we intend to analyze abortions caused by BVDV enfections with serological and

^{*} This study was presented as a poster at the Xth National Veterinary Microbiology Congress (with International participation), 24-27 September 2012, Kuşadası/Aydın.

Investigation of bovine viral diarrhoea virus ...

virological aspects, in dairy cow enterprises in Elazığ, Malatya and Bingöl.

Material and Methods

Study materials include tissue samples taken from 241 waste fetus samples (abortion/stillborn fetus) seen in the events of abortion and sent to the Control Institute of Veterinary Medicine (Elazığ) for etiological diagnosis at small family-type dairy cow enterprises and 602 units of blood samples taken from clinically healthy aborted cows in these enterprises between September 2011 and April 2012 in Elazığ, Malatya and Bingöl (Table 1). Anamnesis revealed that none of these cows received protective vaccination for BVDV and calves born had no chance of living between 6 and 9 months of pregnancy.

BVDV Antigen Test from Tissues of Aborted Fetuses

In order to research BVDV antigen, brain, liver, lung and spleen tissue samples taken from aborted fetus were stirred with 2 ml PBS diluents (1/10). Also MagNaLyser (Roche, Mannheim, Germany) were homogenized in beaded tubes for 3000 xg and 3 minutes. Homogeneta Eppendorf tubes were centrifuged for 10 minutes and 3000 xg and solid materials and beads were removed. Supernatant that remained solution was stored at -80 °C until used as inoculums. The presence of antigens in inoculums was investigated with commercial BVDV Ag ELISA kit (BVDV Ag Serum Plus HerdChek IDEXX Laboratories Westbrook, Maine 04 092 USA). Results were analyzed at 450 nm absorbance filter and as sample-negative control were >0.300 nm, BVDV antigen was evaluated as positive (7, 14, 24).

Table 1. Distribution by province of the number of samples obtained from waste fetus samples taken from dairy cow enterprises and blood serum samples

City	Fetus tiss	sue samples	Blood serum samples		
	n	%	n	%	
Elazığ	66	27.38	180	29.90	
Malatya	11	4.56	32	5.31	
Bingöl	164	68.04	390	64.78	
TOTAL	241		602		

Table 2. Results of BVDV antigen and antibody screening test of dairy cows included in study

City _	BVDV antigen ELISA results of fetus tissue samples				BVDV antibody ELISA results of blood serum samples					
	n	-	%	+	%	n	-	%	+	%
Elazığ	66	49	74.24	17	25.75	180	43	23.88	137	76.11
Malatya	11	7	63.63	4	36.36	32	14	43.75	18	56.25
Bingöl	164	116	70.73	48	29.26	390	119	30.51	271	69.48
TOTAL	241	172	71.36	69	28.63	602	176	29.23	426	70.76

Obtaining of Blood Sera and BVDV Antibody Test

For BVDV antibody search, blood taken from *v. jugularis* of cows was centrifuged at 3000 xg for 10 minutes. Serum remained above was transferred into sterile tubes and stored at -20 °C until use. Presence of antigens in blood was investigated with commercial BVDV total antibody ELISA test kit (IDEXX Laboratories Westbrook, Maine 04092 USA) (8).

Statistic Analysis

For analyzing the difference between the prevalence of the diseases amongst the provinces, Chi-Square test form SPSS software program (Version 13.0) was used (10).

Results

Results of BVDV antigen and antibody screening tests of fetal tissue and blood serum samples of aborted dairy cows are shown at Table 2. It was determined that the difference between the provinces regarding BVDV antigens of the fetal tissues was not significant (P>0.05) whereas that between the same regions in relation to BVDV antibodies was significant (P<0.05) among Elazığ-Malatya and Malatya-Bingöl (Table 3).

Discussion

In postnatal infections due to BVDV commonly seen in ruminants, fertility disorders can be seen. Also in transparental infections, abortions depending on stage of fetal development, embryonic death and resorption of fetus, congenital anomalies, birth of weak calves and newborn calves with appearance healthy but seropositive immune tolerance and persistent infection have been reported. Reports show that calves with persistent infections play an important role at BVDV contamination. Although they seem healthy, they spread virus with their secretions and excretion and this factor plays an important role in scattering of the disease to other animals (1, 18). In addition, reports (1, 18, 20) indicate that deaths due to gastrointestinal and respiratory tract diseases cause large economic losses.

According to Ok (16) abortions caused by BVDV infections can be seen any time at pregnancy and cases of abortion are observed in animals at a variety of periods of pregnancy.

BVDV infections are common worldwide (18, 20) and there have been studies reporting the presence of BVDV infections in our country (5, 11, 17, 23, 25) and in our region (9). Bovine Viral Diarrhoea Virus infections have been reported to have very different seroprevalence (2-6, 11-13, 15, 17, 19, 21-23, 25). Seropositivity has been reported as 58-93% in South Vietnam (6), 5% Switzerland (19), 21% in Spain (13), 35% in North Portugal (15), 84.4% in Britain (2), 32% in Sweden (3), 38% in South Korea (12), 14% in Mexico (21), 94% in Peru (22), 79.2% in Crotia (4) and in our country 86% in Aydın vicinity (23), 81.62% in Northeast Anatolia region (25), 90.63% in Konya region (11) and 82.1%, 39.49%, 83.4% and 78.6% with a total of 64.2% in Ankara, Konya, Muğla ve Samsun respectively (5). Also, in Afyonkarahisar, seropositivity has been found to range between 77.7% and 100% in organized flocks and 57.1% in family-type farms (23).

In this study, in terms of BVDV antibodies in blood sera, results were found to be positive in Elazığ as 76.11% (137/180), in Malatya 56.25% (18/32) and in Bingöl 69.48% (271/390), according to waste

Table 3. The evaluation significance of the diseases between the provinces

	Fetus tiss	ue samples	Blood serum samples		
	χ^2 value	Statistical significant	χ^2 value	Statistical significant	
Elazığ-Malatya	0.134	P>0.05	4.489	P<0.05	
Elazığ-Bingöl	0.189	P>0.05	2.34	P>0.05	
Malatya-Bingöl	0.329	P>0.05	20.59	P<0.001	

Investigation of bovine viral diarrhoea virus ...

samples, seropositivity for BVDV antigens was found as 25.75% (17/66), 36.36% (4/11) and 29.26% (48/164) in Elazığ, Malatya and Bingöl, respectively.

For the determining presence of virus at fetal tissues methods such as virus isolation, RT-PCR and ELISA are used (7, 8, 14, 24). In a comparative study (7), diagnosis of BVDV in blood and fetal fluids were determined with RT-PCR and Ag ELISA, results of both tests are reported as similar. Same researchers stated that Ag ELISA a valuable test method for the diagnosis of BVDV at routine laboratories owing to its good performance, low price, and it can even be used on autolyzed fetuses. In this study, BVDV antigen and antibodies in blood serum and fetal tissue samples were screened with ELISA test kits.

Due to lack of treatment options, in order to prevent BVDV infections, first of all animals with persistent infection should be removed from herd and vaccination programs should be set (1, 16, 18, 20).

As a result, in aborted dairy cows in Elazığ, Malatya and Bingöl, antigens and antibodies have been reported to high-seroprevalance. It was noted that in order to reduce economic losses due to this virus, there should be emphasis on preventive measures.

References

- Batmaz H. Sığırların İç Hastalıkları. Semptomdan Tanıya Tanıdan Tedaviye. Genişletilmiş İkinci Baskı. Bursa: F Özsan Mat San ve Tic Ltd Şti, 2010; pp. 308-12.
- Beaudeau F, Belloc C, Seegers H, Assie S, Sellal E, Joly A. Evaluation of a blocking ELISA forthe detection of bovine viral diarrhoea virus (BVDV) antibodies in serum and milk. Vet Microbiol 2001; 80: 329-37.
- Björkman C, Alenius S, Manuelsson U, Uggla A. Neosporacaninum and bovine viral diarrhoea virus infections in Swedish dairy cows in relationtoabortion. Vet J 2000; 159: 201-6.
- Buik-Rudan N, Cvetnic S, Madic J, Rudan D. Prevalence of antibodies to IBR and BVD viruses in dairy cows withre productive disorders. Theriogenology 1999; 51: 875-81.
- Burgu İ, Alkan F, Yeşilbağ K. The prevalence of persistent BVD virus infection in cattle in Turkey. Ankara Üniv Vet Fak Derg 1999; 46: 169-77.

- Duong MC, Alenius S, Huong LTT, Björkman C. Prevalance of *Neosporacaninum* and bovine viral diarrhoea virus in dairy cows in Southern Vietnam. Vet J 2008; 175(3): 390-4.
- Garcia-Perez AL, Miguijon E, Barandika JF, Aduriz G, Povedano I, Juste RA, Hurtado A. Detection of border disease virus in fetuses, stillbirths, and newborn lambs from natural and experimental infections. J Vet Diagn Invest 2009; 21(3): 331-7.
- Hilbe M, Stalder H, Peterhans E, Haessig M, Nussbaumer M, Egli C, Schelp C, Zlinszky K, Ehrensperger F. Comparison of five diagnostic methods for detecting bovine viral diarrhea virus infection in calves. J Vet Diagn Invest 2007; 19(1): 28-34.
- İssi M, Gülaçtı İ, Kızıl Ö, Karapınar T, Bulut H, Gül Y. The mucosal disease cases observed in our clinic and confirmed with reverse transcription – polymerase chain reaction (RT - PCR). Fırat Üniv Sağlık Bil Derg 2006; 20 (3): 253-8.
- Kalaycı Ş. SPSS Uygulamalı Çok Değişkenli İstatistik Teknikleri. İkinci Baskı, Ankara: Asil Yayın, 2006.
- 11. Kayacan G. Konya ve Çevresinde Bulunan Süt Sığırcılığı İşletmelerindeki Hayvanlara Ait Kan ve Süt Serumlarında Bovine Viral Diarrhoea Virus (BVDV)'una Karşı Oluşan Antikorların ELİSA ile Araştırılması. Yüksek Lisans Tezi. Selçuk Üniv. Sağlık Bilimleri Enstitüsü, Konya-Türkiye, 2008.
- Lee DH, Park SW, Choi EW, Lee CW. Investigation of the prevalence of bovine viral diarrhoea virus in dairy cows in South Korea. Vet Rec 2008; 162: 211-3.
- Mainar-Jaime RC, Berzal-Herranz B, Arias P, Rojo-Vazquez FA. Epidemiological pattern and risk factors associated with bovine viral diarrhoea virus (BVDV) infection in a nonvaccinated dairy cattle population from the Asturias region of Spain. Prev Vet Med 2001; 52: 63-73.
- 14. Nettleton PF, Gilray JA, Russo P, Dlissi E. Border disease of sheep and goats. Vet Res 1998; 29(3-4): 327-40.
- Niza-Ribeiro J, Pereira A, Souza J, Maderia H, Barbosa A, Afonso C. Estimated BVDV prevalence, -control and- vaccine use in dairy herds in Northern Portugal. Prev Vet Med 2005; 72: 81-5.

Infovet, 2010; pp. 72-80.

- Özel M. Yeni Doğan Buzağılarda ve Annelerinde Infectious Bovine Rhinotracheitis (IBR) ve Bovine Viral Diarrhoea (BVD) Enfeksiyonlarının Serolojik Olarak Araştırılması. Yüksek Lisans Tezi. Afyon Kocatepe Üniv. Sağlık Bilimleri Enstitüsü, Afyon-Türkiye, 2008.
- Radostits OM, Gay CC, Hinchcliff KW, Constable PD. Veterinary Medicine. Tenth Edition. Edinburgh, London, New York, Oxford, Philadelphia, St Louis, Sydney, Toronto: Saunders Elsevier, 2008; pp. 1248-77.
- Siegwart N, Hilbe M, Hassig M, Braun U. Increased risk of BVDV infection of calves from pregnant dams on communal Alpine pastures in Switzerland. Vet J 2006; 172: 386-8.
- Smith BP. Large Animal Internal Medicine. Fourth Edition. St Louis, Missouri: Mosby Elsevier, 2009; pp. 791-8.
- 21. Solis-Calderon JJ, Segura-Correa VM, Segura-Correa JC. Bovine viral diarrhoea virus in beef cattle herds of Yucatan, Mexico: seroprevalence and risk factors. Prev Vet Med 2005; 72: 253-62.
- Stahl K, Rivera H, Vagsholm I, Moreno-Lopez J. Bulk milk of testing for antibody seroprevalences to BVDV and BHV-1 in a rural region of Peru. Prev Vet Med 2002; 56: 193-202.
- Tan TM, Karaoğlu T, Erol N, Yıldırım Y. Serological and virological investigations of bovine viral diarrhoea virus (BVDV) infection in dairy cattle herds in Aydın province. Turk J Vet Anim Sci 2006; 30: 299-4.
- 24. Yan L, Zhang S, Pace L, Wilson F, Wan H, Z h a n g M. C o m b i n a t i o n o f reverse transcription real-time polymerase chain reaction and antigen capture enzyme-linked immunosorbent assay for the detection of animals persistently infected with bovine viral diarrhea virus. J Vet Diagn Invest 2011; 23(1): 16-25.

 Yıldırım Y, Burgu İ. The seroprevalence of bluetongue (BT), IBR, PI-3, EBL and BVD infections in cattle in northeastern Anatolia. Ankara Üniv Vet Fak Derg 2005; 52: 113-7.

Correspondence Address :

Doç. Dr. Mustafa İSSİ Fırat Üniversitesi Veteriner Fakültesi, İç Hastalıkları Anabilim Dalı, 23119, Elazığ, TÜRKİYE Tel: +90424 2370000 / 3874 E-posta: mustafaissi@hotmail.com