

PAPER DETAILS

TITLE: Secondary Metabolites from Bioactive Methanolic Extract of *Verbascum pycnostachyum* Boiss. & Helder Flowers

AUTHORS: I İrem TATLI, Wolfgang SCHUHLY, Zeliha S AKDEMİR

PAGES: 23-32

ORIGINAL PDF URL: <https://dergipark.org.tr/tr/download/article-file/842136>

Secondary Metabolites from Bioactive Methanolic Extract of *Verbascum pycnostachyum* Boiss. & Helder Flowers

Received : 02.03.2007

Revised : 09.07.2007

Accepted : 12.07.2007

İ. İrem Tatlı^{*◦}, Wolfgang Schuhly^{}, Zeliha S. Akdemir^{***}**

Introduction

The genus *Verbascum*, which is known as 'mullein', is represented by 228 species, 185 of which are endemic to Turkey¹. The leaves, flowers and whole aerial parts of *Verbascum* L. species have been used to treat respiratory problems, eczema and other types of inflammatory skin conditions in traditional Turkish medicine. They have also been widely utilized as a folk medicine to have a soothing and anti-inflammatory effect on the urinary tract. Additionally, various species are commonly used to treat hemorrhoids, rheumatic pain, superficial fungal infections, wounds and diarrhea. They are traditionally consumed as a tea to relieve abdominal pains²⁻⁴.

The iridoid and phenylethanoid glycosides are widely distributed in the genus *Verbascum*⁵. Although the taxonomic and morphological aspects of the genus *Verbascum* appear more or less complex, the frequent occurrence of the iridoid and phenylethanoid glycosides in the Scrophulariaceae has been used in chemotaxonomic studies^{6, 7}. Iridoids display an interesting spectrum of biological activity such as anti-inflammatory⁸. Likewise, phenylethanoid glycosides are known to possess antioxidant activity⁹.

* Hacettepe University, Faculty of Pharmacy, Department of Pharmaceutical Botany, Sıhhiye, 06100 Ankara, Turkey

** University of Graz, Institute of Pharmaceutical Sciences, Department of Pharmacognosy Karl-Franzens-University, Universitätsplatz 4, A-8010 Graz, Austria

*** Hacettepe University, Faculty of Pharmacy, Department of Pharmacognosy, Sıhhiye, 06100 Ankara, Turkey

◦ Corresponding author. I. I. Tatlı. Tel: 90 312 3051089, Fax: 90 312 3114777, E-mail: itatli@hacettepe.edu.tr

In order to evaluate folkloric utilization, both antinociceptive and anti-inflammatory activities of endemic *Verbascum* species, *V. pycnostachyum* Boiss. & Heldr. was investigated in our previous studies. Antinociceptive activity was investigated via *p*-benzoquinone-induced writhing test, while the anti-inflammatory activity was studied using carrageenan-induced hind paw edema, PGE₁-induced hind paw edema and 12-*O*-tetradecanoyl-13-acetate (TPA)-induced mouse ear edema models in mice. The methanolic extract of the flowers of *V. pycnostachyum* displayed significant antinociceptive and anti-inflammatory activity at 200 mg/kg dose, per os, without inducing any apparent acute toxicity as well as gastric damage¹⁰.

As a part of our continuing search for bioactive agents from *Verbascum* species, we here have report the results of the isolation and structure elucidation of iridoid glycosides, aucubin¹, ajugol², ajugoside³, harpagoside⁴ and a phenylethanoid glycoside, verbascoside⁵ from the bioactive methanolic extract of *Verbascum pycnostachyum* Boiss. & Helder flowers, which is an endemic species distributed in South Anatolia¹.

Materials and Methods

2.1. General Experimental Procedures

The UV spectra (λ_{\max}) were recorded on a Hitachi HP 8452 A spectrophotometer. The IR spectra (ν_{\max}) were determined on ATI Mattson Genesis Series FTIR spectrophotometer. The ¹H and ¹³C NMR spectra were obtained on Bruker Avance DRX 500 and 300 spectrometer operating at 500 and 300 MHz for ¹H NMR and at 125 and 75 MHz for ¹³C NMR spectra. The chemical shift values are reported as parts per million (ppm) relative to tetramethylsilane (TMS), and the coupling constants are in hertz (Hz, in parentheses). LC-ESIMS FT data were obtained using a Bruker BioApex FT-MS instrument in the ESI mode. Reverse-phase material (C-18, Sepra-lyte 40 μ m) was used for vacuum liquid chromatography (VLC). Medium pressure liquid chromatography (MPLC) separations were performed on a Labomatic glass column packed with LiChroprep RP-18 (Merck), using a Lewa M5 peristaltic pump. Si gel (230-400 mesh) (Merck) and Sephadex LH-20 were used for column chromatography (CC). Pre-coated silica gel 60 F₂₅₄ aluminum sheets (Merck) were used for thin-layer chromatography (TLC) with developing solvent-system, CHCl₃-MeOH-H₂O (61:32:7).

Plates were examined by UV fluorescence and sprayed with 1% vanillin in concentrated H_2SO_4 , followed by heating at 105°C for 1-2 mins.

2.2. Plant Material



Verbascum pycnostachyum Boiss.& Heldr. (Scrophulariaceae) was collected from Mut to Karaman, 1300 m, in June 2000. A voucher specimen was deposited in the Herbarium of the Pharmacognosy Department, Faculty of Pharmacy, Hacettepe University, Ankara, Turkey (HUEF 00182).

2.3. Extraction and Isolation


The air-dried and powdered flowers of *Verbascum pycnostachyum* (320.46 g) were extracted twice with MeOH (2x2000 ml) at 40°C . After evaporation of the combined extract in vacuo, 48.71 g MeOH extract was obtained. The isolation of compounds was guided on TLC autographic assay using 0.2 % DPPH solution in MeOH to search for potential antioxidant molecules. The crude extract (48.71 g) was fractionated by vacuum-liquid chromatography over reverse-phase material (VLC, 350 g), eluting with H_2O and gradient MeOH- H_2O mixtures (5-30 %) to yield compounds 1 (326.9 mg), 2 (317.1 mg) and fraction A. Fraction A (1.3 g) was subjected to vacuum liquid chromatography (VLC) using reversed-phase material (Separylite 40 μm , 175 g), employing MeOH/ H_2O mixtures (0-50 %) to give compound 5 (152.4 mg) and fraction A1. Fraction A1 (278.5 mg) was carried out on C_{18} -MPLC using gradient H_2O -MeOH mixtures (10-100%) to give fractions A1a-d. Fraction A1a (53.6 mg) was chromatographed on a Si gel column (8 g) eluted with CHCl_3 -MeOH mixtures (90:10, 85:15, 80:20, 70:30) and CHCl_3 -MeOH- H_2O mixtures (80:20:2) to yield compound 3 (8.5 mg) and fraction A1aI. Fraction A1aI (6.7 mg) was further purified on a Sephadex LH-20 (10 g) column using MeOH to give compound 4 (3.3 mg).

Results and Discussion

Compounds 1-5 were isolated from the methanolic extract of the flowers of *Verbascum pycnostachyum* by a combination of vacuum liquid chromatography (VLC) and open column chromatographic methods, with the following results (Fig.).

Aucubin (**1**)

	R ₁	R ₂
Ajugol (2)	H	OH
Ajugoside (3)	H	OCOCH ₃
Harpagoside (4)	OH	trans-cinnamoyl

Verbascoside (**5**)**Figure 1**

Isolated compounds from *Verbascum pycnostachyum*

Aucubin (1): UV (MeOH, λ_{\max} , nm): 202. IR (KBr, ν_{\max} , cm⁻¹): 3630 (OH), 1665 (C=C), 1545, 1360 (aromatic ring). Positive ion LC-ESIMS m/z 368 ((M+Na)⁺, calc. for C₂₁H₃₂O₁₃). ¹H NMR (500 MHz, DMSO-*d*₆) and ¹³C NMR (125 MHz, DMSO-*d*₆) data were superimposable with those reported in the literature¹¹.

Ajugol (2): UV (MeOH, λ_{\max} , nm): 220. IR (KBr, ν_{\max} , cm^{-1}): 3470 (OH), 1655 (C=C). Positive ion- LC-ESIMS m/z 370 ($(M+Na)^+$, calc. for $C_{15}H_{24}O_9$). ^1H (500 MHz, $\text{DMSO}-d_6$) and ^{13}C (125 MHz, $\text{DMSO}-d_6$) NMR data (Table) were superimposable with those reported in the literature¹².

TABLE I
 ^1H - and ^{13}C -NMR (500 and 125 MHz, $\text{DMSO}-d_6$) data of compounds **2**, **3**.

	2			3		
	δ_{C}	δ_{H}	J (Hz)	δ_{C}	δ_{H}	J (Hz)
Aglycone						
1	92.7	5.45 s	-	92.1	5.63 s	-
3	139.4	6.15 d	5.1	139.7	6.20 d	6.4
4	104.9	4.85 †	-	103.4	4.70 d	6.4
5	40.3	2.72 m	-	39.7	2.62 m	-
6	77.2	3.90 †	-	74.5	3.86 d	7.6
7a	49.0	1.79 dd	4.5/13.4	47.3	2.00 dd	5.5/15.0
7b		2.04 dd	5.6/13.4		2.10 d	15.0
8	78.5	-	-	87.8	-	-
9	50.8	2.54 d	9.4	48.0	2.66 m	-
10	24.2	1.31 s	-	22.3	1.45 s	-
OCOCH_3	-			170.0	-	-
OCOCH_3	-			22.0	1.94 s	-
β -Glucose						
1'	98.4	4.64 d	7.9	97.8	4.46 d	7.8
2'	73.8	3.15- 3.40 †	-	73.0	2.96 t	8.9
3'	76.8	3.15- 3.40 †	-	76.6	3.14 t	8.9
4'	70.7	3.19 t	8.7	70.0	3.05 d	9.0
5'	77.0	3.15- 3.40 †	-	76.8	3.13 m	-
6'a	61.9	3.66 dd	4.8/11.6	61.1	3.46 dd	6.0/12.0
6'b		3.89 †	-		3.69 d	12.0

† Signal patterns are unclear due to overlapping

Ajugoside (3): UV (MeOH, λ_{\max} , nm): 224. IR (KBr, ν_{\max} , cm^{-1}): 3450 (OH), 1705 (C=O), 1650 (C=C). Positive ion- LC-ESIMS m/z 412 ($(\text{M}+\text{Na})^+$, calc. for $\text{C}_{17}\text{H}_{26}\text{O}_{10}$). ^1H (500 MHz, $\text{DMSO}-d_6$) and ^{13}C (125 MHz, $\text{DMSO}-d_6$) NMR data (Table) were superimposable with those reported in the literature^{13, 14}.

Harpagoside (4): UV (MeOH, λ_{\max} , nm): 228. IR (KBr, ν_{\max} , cm^{-1}): 3600 (OH), 1705 (C=O), 1637 (C=C), 1604, 1363 (aromatic ring). Positive ion LC-ESIMS m/z 517 ($(\text{M}+\text{Na})^+$, calc. for $\text{C}_{24}\text{H}_{30}\text{O}_{11}$). ^1H NMR (500 MHz, $\text{DMSO}-d_6$) and ^{13}C NMR (125 MHz, $\text{DMSO}-d_6$) data were superimposable with those reported in the literature⁹.

Verbascoside (= acteoside, $(\beta\text{-(3,4-dihydroxyphenyl)-ethyl})\text{-(3'-O-}\alpha\text{-L-rhamnopyranosyl)-(4'-O-caffeoyl)-}\beta\text{-D-glucopyranoside}$) (5): UV (MeOH, λ_{\max} , nm): 212, 332. IR (KBr, ν_{\max} , cm^{-1}): 3689 (OH), 1708 (C=O), 1634 (C=C), 1604, 1515, 1385 (aromatic ring). Positive ion-LC-ESIMS m/z 647 ($(\text{M}+\text{Na})^+$, calc. for $\text{C}_{29}\text{H}_{36}\text{O}_{15}$). ^1H NMR (300 MHz, $\text{DMSO}-d_6$) and ^{13}C NMR (75 MHz, $\text{DMSO}-d_6$) data superimposable with those reported in the literature⁹.

Compound **1** was obtained as an amorphous powder. Its structure was identified as aucubin¹⁵ by comparing its ^1H and ^{13}C NMR data with previously published data and by direct comparison with an authentic sample on a TLC plate¹¹.

Compound **2** was isolated as a yellow amorphous powder with the molecular formula $\text{C}_{15}\text{H}_{24}\text{O}_9$ (LC-ESIMS m/z 370.9 ($\text{M}+\text{Na})^+$). Its UV spectrum suggested the presence of an iridoid enolether system (220 nm) and in its IR spectra absorption bands were typical for a hydroxyl group (3416 cm^{-1}) and a double bond (1656 cm^{-1}). The ^1H and ^{13}C NMR spectra of **2** (see Table) are superimposable with those of ajugol¹². Based on this evidence, compound **2** was identified as ajugol.

Compound **3** proved to have the molecular formula $\text{C}_{17}\text{H}_{26}\text{O}_{10}$, as seen from the positive-ion ESIMS (m/z 412 ($\text{M}+\text{Na})^+$) combined with ^1H and ^{13}C NMR data (see Table). The UV and IR data of compound **3** showed that **3** consist of a non-conjugated enol-ether system. The ^1H NMR signals at δ_{H} 6.20 (d , $J = 6.4\text{ Hz}$), 4.70 (d , $J = 6.4\text{ Hz}$) were attributed to H-3 and H-4, respectively, whose chemical shift values and multiplicities indicated that C-5 was non-substituted. This assumption was also supported by the H-9 signal (δ_{H} 2.66, m). On the other hand, the multiplet signal at

δ_{H} 3.86 was attributed to an oxymethine proton at C-6 (δ_{C} 74.5), which was coupled to H₂-7 (δ_{H} 2.00, *dd*, $J = 5.5/15.0$ and 2.10, *d*, $J = 15.0$ Hz) methylene protons. In the ^1H NMR spectrum of **3**, δ_{C} 170.0 and 22.0 signal patterns implied the presence of an acethyl group. Thus, the location of the acethyl group was ascertained from downfield acetylation shifts (ca. 9.3 ppm) observed for C-8 (δ_{C} 87.8) resonance comparing with that of ajugol (δ_{C} 78.5)¹². Accordingly, the structure of **3** was determined to be ajugoside^{13, 14}.

Compound **4** was obtained as amorphous powder whose UV spectra indicated its non-conjugated enol-ether functional group. Its IR spectra showed absorption bands typical of conjugated carbonyl groups. The molecular formula of compound **4** was determined by LC-ESIMS, which exhibited a pseudomolecular ion at m/z 517 ($\text{M}+\text{Na}$)⁺, and ^1H and ^{13}C NMR data as $\text{C}_{24}\text{H}_{30}\text{O}_{11}$. The ^1H NMR spectrum of **4** revealed the resonances of two olefinic protons, observed as an AX system, at δ_{H} 6.47 and 7.53 (*d*, $J_{\text{AX}} = 16.0$ Hz) and 5 aromatic protons at δ_{H} 7.34 (1H), 7.35 (2H) and 7.62 (2H), consistent with the presence of a *trans*-cinnamoyl moiety. The chemical shift values of both C-8 and H₃-10 indicated that the acyl group was attached at C-8. From the above findings and comparison with the published data, compound **4** was considered identical to harpagoside⁹.

Compound **5** was also obtained as an amorphous powder. Its structure was identified as verbascoside⁹ by comparing its ^1H and ^{13}C NMR data with previously published data and by direct comparison with the authentic sample on a TLC plate.

Conclusion

Concerning the iridoid and phenylethanoid glycosides of the genus *Verbascum*, the isolation of iridoid glucosides, aucubin¹, ajugol², harpagoside⁴ and a phenylethanoid glycoside, verbascoside⁵ from several other *Verbascum* species has been reported previously⁵. It is well known that these compounds are common iridoid and phenylethanoid glycosides and taxonomic markers in the genus *Verbascum* and family Scrophulariaceae. To the best of our knowledge, ajugoside³ has been isolated from *Verbascum* species for the first time. Additionally, this is the first report on the isolation and characterization of all these compounds from *Verbascum pycnostachyum* as well as a *Verbascum* species from Group K of the ge-

nus (1), although several sterols such as β -sitosterol and stigmasterol were isolated from *V. pycnostachyum* in previous studies¹⁵. Our continuing studies will be of assistance in clarifying the chemotaxonomic classification of the genus *Verbascum*.

Results of our previous study had clearly demonstrated that the methanolic extract of the flowers of *Verbascum pycnostachyum* possess significant antinociceptive and anti-inflammatory activities which support the traditional utilization in Turkey¹⁰. The isolated compounds, aucubin¹ was also found to possess significant antinociceptive and anti-inflammatory activities, per os without inducing any apparent acute toxicity or gastric damage⁸. Harpagoside⁴ and verbascoside⁵, exhibited a dose-dependent inhibition of bioautographic and spectrophotometric DPPH activities^{9, 16}.

In connection with the role of aucubin as well as the roles of harpagoside and verbascoside which were identified as free radical scavengers of *V. pycnostachyum*, it seems that they could be synergistic with each other in the methanolic extract. In order to correlate the obtained data in the field, further examinations in different assays can be evaluated.

Summary

Secondary Metabolites from Bioactive Methanolic Extract of *Verbascum pycnostachyum* Boiss. & Helder Flowers

Four iridoid glucosides, aucubin¹, ajugol², ajugoside³, harpagoside⁴, and a phenylethanoid glycoside, verbascoside⁵ were isolated from the flowers of bioactive methanolic extract of *Verbascum pycnostachyum* Boiss & Helder. The structures of the compounds were determined from spectral methods (UV, IR, 1D NMR and Mass Spec.). Ajugoside³ is encountered for the first time from *Verbascum* species.

Keywords: Scrophulariaceae, *Verbascum pycnostachyum* Boiss & Helder, iridoid glucosides, phenylethanoid glycoside.

Özet

***Verbascum pycnostachyum* Boiss. & Helder Çiçeklerinin Biyoaktif Metanol Ektresinin Sekonder Metabolitleri**

Verbascum pycnostachyum Boiss & Helder'in çiçekli kısımlarının biyoaktif metanol ekstresinden, dört iridoit glukoziti, okubin¹, aju-

gol², ajugozit³, harpagozit⁴ ve bir feniletanoit glikoziti, verbaskozit⁵ izole edilmiştir. Bileşiklerin yapıları spektral yöntemlerle (UV, IR, 1D NMR ve Mass Spekt.) tespit edilmiştir. Ajugozit³ ilk defa *Verbascum* türlerinden elde edilmiştir.

Anahtar kelimeler: Scrophulariaceae, *Verbascum pycnostachyum* Boiss & Helder, iridoit glukozitleri, feniletanoit glikoziti.

Acknowledgements

The authors are grateful to Prof. Dr. Hayri Duman (Gazi University, Faculty of Science, Etiler, Ankara, Turkey) for the authentication of the plant specimen.

REFERENCES

1. Huber-Morath, A. *Verbascum*, in "Flora of Turkey and the East Aegean Islands", Davis, P. H. (Ed.), Edinburgh, University Press, (1978), Vol. 6, 461.
2. Tuzlaci, E., Erol, M. K.: Turkish folk medicinal plants. Part II: Egirdir (Isparta). *Fitoterapia*, 70, 593 (1999)
3. Sezik, E., Yesilada, E., Honda, G., Takaishi, Y., Takeda, Y., Tanaka, T.: Traditional medicine in Turkey X. Folk medicine in Central Anatolia. *J. Ethnopharmacol.*, 75, 95 (2001)
4. Turker, A., Camper, N. D.: Biological activity of common mullein, a medicinal plant. *J. Ethnopharmacol.*, 82, 117 (2002)
5. Tatli, I. I., Akdemir, Z. S.: *Chemical constituents of Verbascum L. species. J. Pharm. Sci. (FABAD)*, 29 (2), 93 (2004)
6. Boros, C. A., Stermitz, F. R.: Iridoids-An Updated Review 1. *J. Nat. Prod.*, 53 (5), 1055 (1990)
7. Jimenez, C., Riguera, R.: Phenylethanoid Glycosides in Plants: Structure and Biological activity. *Natural Product Reports*, 591 (1994)
8. Kupeli, E., Tatli, I. I., Akdemir, Z. S., Yesilada, E.: Bioassay-guided isolation of anti-inflammatory and antinociceptive glycoterpenoids from the flowers of *Verbascum lasianthum* Boiss ex Benth. *J. Ethnopharmacol.*, 110, 444 (2007)
9. Akdemir, Z. S., Tatli, I. I., Bedir, E., Khan, I. A.: Iridoid and phenylethanoid glycosides from *Verbascum lasianthum*. *Turk. J. Chem.*, 28, 227 (2004)
10. Tatli, I. I., Kupeli, E., Yesilada, E., Akdemir, Z. S.: Assessment of antinociceptive and anti-inflammatory activity on seven endemic *Verbascum L.* species growing in Turkey. *J. Ethnopharmacol.*, (2007) (*submitted*).
11. Akdemir, Z. S., Tatli, I. I., Bedir, E., Khan, I. A.: Acylated iridoid glycosides from *Verbascum lasianthum*. *Turk. J. Chem.*, 28, 101 (2004)
12. Tatli, I. I., Akdemir, Z. S., Bedir, E., Khan, I. A.: Saponin, iridoid, phenylethanoid and monoterpene glycosides from *Verbascum pterocalycinum* var. *mutense*. *Turk. J. Chem.*, 28, 111 (2004)
13. Guiso, M., Marinibe, R., Agostini, A.: Ajugoside and ajugol-Structure and configuration. *Gazzetta Chimica Italiana*, 104 (1-2), 25 (1974)

14. Davini, E., Iavarone, C., Trogolo, C.: Revised stereochemistry of C(6), hydroxyl group in ajugol and ajugoside. *Gazzetta Chimica Italiana*, 112 (1-2), 57 (1982)
15. Erlacin, S., Gozler, B.: Sterols and sterol glycosides of *Verbascum pycnostachyum* Boiss. et Heldr. *Pharmazie*, 37, 149 (1982)
16. Tatli, I. I., Takamatsu, S., Khan, I. A., Akdemir, Z. S.: Screening for Free Radical Scavenging and Cell-Aggregation Inhibitory Activities on Secondary Metabolites from Turkish *Verbascum* species. *Z. Naturforsch. C*, (2007) (*in press*)