

PAPER DETAILS

TITLE: An Anomaly of Pecten in Mesobuthus turcicus Kovarík et al., 2022 (Scorpiones: Buthidae)

AUTHORS: Ersen Aydın YAGMUR,Özgün SIPAHIOGLU,Ömer YILMAZ,Mehmet Sait KILIÇ

PAGES: 116-118

ORIGINAL PDF URL: <https://dergipark.org.tr/tr/download/article-file/2334864>

An Anomaly of Pecten in *Mesobuthus turcicus* Kovařík et al., 2022 (Scorpiones: Buthidae)

Ersen Aydın YAĞMUR^{*1}, Özgün SIPAHIOĞLU², Ömer YILMAZ³, Mehmet Sait KILIÇ⁴

¹Manisa Celal Bayar University, Alaşehir Vocational School, Alaşehir, Manisa, 45600, TÜRKİYE

²Göztepe Neighborhood, Ege Street, 3/15, Kadıköy, İstanbul, TÜRKİYE

³Mimar Sinan Anatolian High School, Bostancı Neighborhood, Koçmustafa Street, No. 3, Şahinbey, Gaziantep, TÜRKİYE

⁴Sarsalkım Secondary School, No. 285, Sarısalkım Neighborhood, Street No. 121001, Şahinbey, Gaziantep, TÜRKİYE

ORCID ID: Ersen Aydın YAĞMUR: <https://orcid.org/0000-0002-0396-3975>; Özgün SIPAHIOĞLU: <https://orcid.org/0000-0001-7160-8775>;

Ömer YILMAZ: <https://orcid.org/0000-0002-6214-6510>; Mehmet Sait KILIÇ: <https://orcid.org/0000-0003-2858-037X>

Received: 27.03.2022

Accepted: 16.04.2022

Published online: 20.06.2022

Issue published: 30.06.2022

Abstract: An anomaly was recorded in the pecten shape of the scorpion species *Mesobuthus turcicus* Kovařík et al., 2022 that was recently described from Türkiye. In this case, it is observed that marginal and median lamellae and fulcræ are fused in the right pecten. For this reason, the pecten is bent counter clockwise. Different from other pectinal anomalies, the size of teeth is developed normally but teeth number is fewer. This teratological anomaly on pecten is described and illustrated. This is the first report of pecten anomaly based on marginal and median lamellae.

Keywords: Teratology, Türkiye, first report, scorpion.

Mesobuthus turcicus Kovařík et al., 2022'ta bir pektin anomalisi (Scorpiones: Buthidae)

Öz: Yakın zamanda Türkiye'den tanımlanan *Mesobuthus turcicus* Kovařík et al., 2022 akrep türünün pekten şeklinde bir anomali kaydedilmiştir. Bu vakada sağ pektende marjinal ve medyan lameller ve fulkralar kaynaştığı gözlenmiştir. Bu sebeple pekten saat yönünün tersine bükülmüştür. Diğer pektin anomalilerinden farklı olarak diş boyutları normal gelişmiştir ancak diş sayısı daha azdır. Pecten üzerindeki bu teratolojik anomali tanımlanmış ve resimlendirilmiştir. Bu, marjinal ve medyan lamellerden kaynaklanan ilk pekten anomalisinin raporudur.

Anahtar kelimeler: Teratoloji, Türkiye, ilk rapor, akrep.

Anomalies on scorpion body occurring during embryonic development cause teratologic disorders (Yağmur et al., 2021). Anomalies may be deformations (malformations) on body parts of the scorpion body. Some parts may be absent; in addition, duplications, division or fusion may be seen on some scorpion parts. Duplications cases were reported by Berland (1913), Karataş & Kürtüllü (2006), and Teruel & Baldazo-Monsivaiz (2015) on prosoma, pedipalp, and pectinal organs, respectively. Metasoma, vesicle, and aculeus duplications are most common duplication types reported by many scholars (Alqahtani & Badry, 2021; Berland, 1913; Briseño, 1963; Campos, 1918; Galvis & Flórez-D., 2016; Lourenço & Hypolite, 2010; Sadine, 2021; Salabi et al., 2021; Seiter & Teruel, 2014; Sergeant, 1946; Shulov & Amitai, 1955; Sissom & Shelley, 1995; Vachon, 1952, 1953, 1972; Williams, 1971). Pedipalp fusion was reported by Cao & Solórzano (1991) and pedipalp malformation cases were reported by Mattoni (2005), Graham (2006), and Jahanifard et al. (2008).

Fusion and division cases on carapace and tergites were reported by Armas (1976), Teruel (2003) and Mattoni (2005); pedipalp fusion by Cao & Solórzano (1991); leg malformation by Armas (1977); pectinal malformation by Ayrey (2011) and Šarić & Tomić (2020), and pedipalp malformation by Mattoni (2005), Graham (2006), and Jahanifard et al. (2008). In addition, Jahanifard et al. (2008) reported a vesicle malformation case and David (2012) reported a leg absence. Teruel (2003) compiled pedipalp,

tergite, and cheliceran anomalies. Yağmur et al. (2021) reported a malformation on chelicera.

Only a few anomaly cases reported on scorpion pectines. Teruel & Baldazo-Monsivaiz (2015) reported a pectinal duplication based on gynandromorphism but pectines were well developed. Ayrey (2011) and Šarić & Tomić (2020) reported malformation on pectines. Ayrey (2011) reported that a pectinal anomaly on a female *Vaejovis lapidicola* on both two pectines. Pectines of the specimen are quite shriveled and shorter than the normal pectines. The abnormal pectines have significantly shorter anterior lamellae than the normal pectines. On the abnormal pectines the tooth area is abnormal, large majority of the teeth are fused into a single irregularly shaped sclerite but only the most distal tooth developed. Šarić & Tomić (2020) reported in the scorpion *Euscorpis* sp. that some pecten teeth are fused or malformed. They also reported that some of the fulcræ are also malformed.

Mesobuthus turcicus was described by Kovařík et al. (2022) just recently from Konya (Karapınar District), Türkiye in two localities. This species is known in these two localities. The adult female of *M. turcicus* (Figs. 1A-B) was collected from Meke Salt Lake, Karapınar, Konya Province, Türkiye (37°41'39"N 33°38'33"E, 1033 m, 09.07.2021, leg. E. A. Yağmur & Ö. Sipahioğlu, AZMM/Sco-2021:18). This specimen was collected with type specimens of *M. turcicus*. It is preserved in 96% ethanol and deposited in AZMM (Alaşehir Zoological

*Corresponding author: ersen.yagmur@gmail.com

Museum, Manisa Celal Bayar University, Alaşehir, Manisa, Türkiye). Identification of the specimen was done according to Kovařík et al. (2022). Terminology for pectines follows Hjelle (1990).

The examined specimen has an abnormal right pecten whereas left pecten is normal (Figs. 2A-B). The pectines are comb-like sensory organs which are present only in scorpions. The pectines consist of three marginal lamellae and a variable number of median lamellae, fulcra, and pectinal teeth (Hjelle, 1990). An area called the peg sensilla is always located at the anteroventral edge of each pectinal tooth (Hjelle, 1990). Besides, the pectines generally are with various macrosetae and microsetae (Hjelle, 1990). The normal pecten has 3 marginal lamellae, 7 median lamellae, 21 fulcra and 21 pectinal teeth. Besides there are numerous macrosetae and microsetae on marginal lamellae and median lamellae. All marginal and median lamellae and fulcrae are fused (except one fulcrum exist on proximal of the pecten) but size of teeth is developed normally. Therefore, the pecten is bent counter clockwise; besides, the teeth number is fewer. The abnormal pecten has 11 teeth whereas normal pecten has 21 teeth. Numerous setae present on normal pecten. Only one seta was detected on fulcrum and six setae on the rest of the abnormal pecten (Figs. 3A-B).

A few cases of pectinal anomaly have been reported up to now. Teruel & Baldazo-Monsivaiz (2015) reported a pectinal duplication because of gynandromorphism. There are two couple pectens in this case that normally developed. The cases of Ayrey (2011) and Šarić & Tomić (2020) are simply about fusion of teeth. The teeth sizes and shapes are normal and they are not fused but fewer in number in this case. Pectens are not bent and some teeth are fused in the cases by Ayrey (2011) and Šarić & Tomić (2020) that is different from our case. This is the first report of the anomaly based on undeveloped marginal lamellae and median lamellae.




Figure 1. Female *Mesobuthus turcicus*. A. Dorsal view. B. Ventral view. Scale bar: 10 mm.

Ethics committee approval: Ethics committee approval is not required for this study.

Conflict of interest: The authors declare that there is no conflict of interest.




Figure 2. Carapace and mesosoma of ventral view of *Mesobuthus turcicus*. A. Under white light. B. Under UV light.




Figure 3. ventral view of coxosternal area of *Mesobuthus turcicus*. A. Under white light. B. Under UV light.

Author Contributions: Conception – Ö.Y.; Design – M.S.K.; Supervision – E.A.Y.; Fund – M.S.K.; Materials – E.A.Y., Ö.S.; Data Collection or Processing – E.A.Y.; Analysis Interpretation – E.A.Y.; Literature Review – E.A.Y.; Writing – E.A.Y., Ö.S.; Critical Review – M.S.K., Ö.Y.

References

- Alqahtani, A.R., & Badry, A. (2021). A rare telson anomaly in *Parabuthus liosoma* (Ehrenberg, 1828) (Scorpiones: Buthidae). *Euscorpius*, 336, 1-4. <http://doi.org/10.5281/zenodo.5742109>
- Armas, L.F. (1976). Escorpiones del archipiélago Cubana. Familia Diplocentridae (Arachnida: Scorpionida). *Poeyana*, 147, 1-35.

- Armas, L.F. (1977). Anomalías en algunos Buthidae (Scorpionida) de Cuba y Brasil. *Poeyana*, 176, 1-6.
- Ayrey, R.F. (2011). An anomaly of pectinal organs in *Vaejovis lapidicola* (Scorpiones: Vaejovidae). *Euscorpius*, 130, 1-6. <https://dx.doi.org/10.18590/euscorpius.2011.vol2011.iss130.1>
- Berland, L. (1913). Note sur un Scorpion muni de deux queues. *Bulletin de la Société entomologique de France*, 18, 251-252.
- Briseno, C. (1963). Presencia de un ejemplar de alacran de la especie *Centruroides noxius*, con dos colas. *Revista del Instituto de Salubridad y Enfermedades Tropicales (México)*, 23(3/4), 185-186.
- Campos, F. (1918). Algunos casos teratologicos observados en los Arthropodos. *Annals of the Entomological Society of America*, 11, 97-98.
- Cao, J., & Solórzano, L. (1991). Escorpión con pedipalpo anómalo. *Resúmenes II Simposio de Zoología, La Habana*: p. 48.
- Galvis, W., & Flórez-D, E. (2016). A new telson teratology in the scorpion *Opisthacanthus* Peters, 1861 (Scorpiones: Hormuridae). *Arachnology*, 17(3), 157-158. <https://doi.org/10.13156/arac.2006.17.3.157>
- David, D. (2012). A seven-legged scorpion: the first teratological leg absence found in *Scorpio maurus fuscus* (Scorpiones: Scorpionidae). *Euscorpius*, 151, 1-4. <https://dx.doi.org/10.18590/euscorpius.2012.vol2012.iss151.1>
- Graham, M.R. (2006). Malformed pedipalp finger dentition of the scorpion *Superstitionia donensis* (Scorpiones: Superstitioniidae). *Euscorpius*, 42, 1-4. <https://dx.doi.org/10.18590/euscorpius.2006.vol2006.iss42.1>
- Hjelle, J.T. (1990). Anatomy and morphology. Pp. 9-63 in: Polis, G.A. (ed.), *Biology of Scorpions*. Stanford, CA: Stanford University Press.
- Jahanifard, E., Navidpour S., & Mashipour, B. (2008). Pedipalps and venom vesicle anomalies in two families of scorpions (Scorpiones: Hemiscorpidae, Buthidae) from Iran. *Pakistan Journal of Biological Sciences*, 11(2), 309-311. <https://dx.doi.org/10.3923/pjbs.2008.309.311>
- Karatas, A., & Kürtüllü, M. (2006). Duplication of pedipalp segments in the scorpion *Androctonus crassicauda* (Olivier, 1807) (Scorpiones: Buthidae). *Euscorpius*, 43, 1-4. <https://dx.doi.org/10.18590/euscorpius.2006.vol2006.iss43.1>
- Kovářík, F., Fet, V., Gantenbein, B., Graham, M.R., Yağmur, E.A., Štáhlavský, F., Poverennyi, N.M., & Novruzov, N.E. (2022). A revision of the genus *Mesobuthus* Vachon, 1950, with a description of 14 new species (Scorpiones: Buthidae). *Euscorpius*, 348, 1-189.
- Lourenço, W.R., & Hypolite, F. (2010). A new case of duplication of the metasoma and telson in the scorpion *Euscorpius flavicaudis* (DeGeer, 1778) (Euscorpidae). *Euscorpius*, 102, 1-2. <https://dx.doi.org/10.18590/euscorpius.2010.vol2010.iss102.1>
- Mattoni, C.I. (2005). Tergal and sexual anomalies in bothriurid scorpions (Scorpiones, Bothriuridae). *The Journal of Arachnology*, 33(2), 622-628. <https://doi.org/10.1636/04-125.1>
- Sadine, S.E. (2021). A remarkable bifid aculeus in *Androctonus amoreuxi* (Audouin, 1826) from central Algeria (Scorpiones: Buthidae). *Revista ibérica de aracnología*, 38, 191-192.
- Salabi, F., Jafari, H. & Forouzan, A. (2021). Report of a Rare Anomaly in the Metasoma of *Hottentotta zagrosensis* (Scorpiones: Buthidae). *Iranian Journal of Science and Technology, Transactions A: Science*, 45(2), 405-408. <https://doi.org/10.1007/S40995-020-01057-1>
- Šarić, M., & Tomić, J. (2016). The first record of malformed pectines in genus *Euscorpius* (Scorpiones: Euscorpidae). *Euscorpius*, 221, 1-10. <https://dx.doi.org/10.18590/euscorpius.2016.vol2016.iss221.1>
- Seiter, M., & Teruel, R. (2014). Two new cases of metasomal duplication in scorpions, with notes on their reproductive biology (Scorpiones: Buthidae). *Revista Ibérica de Aracnología*, 24, 127-129.
- Sergent, E. (1946). Anomalies chez les scorpions. *Archives de l'Institut Pasteur d'Algérie*, 24(1), 80-82.
- Shulov, A., & Amitai, P. (1955). A scorpion *Leiurus quinquestratus* H. et E. with two stings. *Bulletin of the Research Council, Israel*, 5B(2), 193-194.
- Sissom, W.D., & M. Shelley, R. (1995). Report on a rare developmental anomaly in the scorpion, *Centruroides vittatus* (Buthidae). *The Journal of Arachnology*, 23, 199-201.
- Teruel, R. (2003). Nuevos casos de anomalías morfológicas en escorpiones (Scorpiones: Bothriuridae, Euscorpidae, Hemiscorpidae, Ischnuridae, Iuridae, Buthidae, Chactidae, Chaerilidae, Diplocentridae, Scorpionidae). *Revista Ibérica de Aracnología*, 7, 235-238.
- Teruel, R., & Baldazo-Monsivaiz, J.G. (2015). Hermaphroditism, gynandromorphism, and four pectines: an extreme case of developmental anomaly in scorpions (Scorpiones: Vaejovidae). *Euscorpius*, 197, 1-7. <https://dx.doi.org/10.18590/euscorpius.2015.vol2015.iss197.1>
- Vachon, M. (1952). Etudes sur les Scorpions. Institut Pasteur d'Algérie, Algér, 482 pp.
- Vachon, M. (1953). The biology of scorpions. *Endeavour*, 12, 80-87.
- Vachon, M. (1972). Remarques sur les scorpions appartenant au genre *Isometrus* H. et E. (Buthidae). A propos de l'espèce *Isometrus maculatus* (Geer) habitant l'Île de Pâques. *Cahiers du Pacifique*, 16, 169-180.
- Williams, S.F. (1971). Developmental anomalies in the scorpion *Centruroides sculpturatus*. *Pan-Pacific Entomologist*, 47 (1), 76-77.
- Yağmur, E.A., Kılıç, M.S., & Yılmaz, Ö. (2021). An anomaly of chelicera in *Scorpio kruglovi* Birula, 1910 (Scorpiones: Scorpionidae). *Euscorpius*, 335, 1-4. <http://doi.org/10.5281/zenodo.5742101>