

PAPER DETAILS

TITLE: ANALYSIS OF SOCIODEMOGRAPHIC CHARACTERISTICS OF EARLY AND ADVANCED STAGE CHRONIC KIDNEY DISEASE PATIENTS

AUTHORS: Ustun YILMAZ,Selin KIR,Ayca INCI,Nilgun AKSOY,Gulay DEMIRTAS,Semih GUL,Ummuhan MADEN

PAGES: 424-429

ORIGINAL PDF URL: <https://dergipark.org.tr/tr/download/article-file/1919476>

Analysis of Sociodemographic Characteristics of Early and Advanced Stage Chronic Kidney Disease Patients

Erken ve İleri Evre Kronik Böbrek Hastalarının Sosyodemografik Özelliklerinin Analizi

Ustun YILMAZ¹, Selin KIR², Ayca INCI¹, Nilgun AKSOY³, Gulay DEMIRTAS¹, Semih GUL¹, Ummuhan MADEN¹

¹ University of Health Sciences Antalya Training and Research Hospital, Nephrology, Antalya, Türkiye

² University of Health Sciences Antalya Training and Research Hospital, Internal Medicine, Antalya, Türkiye

³ Akdeniz University Faculty of Nursing, Surgical Nursing, Antalya, Türkiye

Yazışma Adresi
Correspondence Address

Ustun YILMAZ
University of Health Sciences
Antalya Training and Research
Hospital, Nephrology,
Antalya, Türkiye
ustunfy@gmail.com

Geliş tarihi / Received : Tem 28, 2020
Kabul tarihi / Accepted : Ara 11, 2020
Elektronik yayın tarihi : Eyl 01, 2021
Online published

Bu makalede yapılacak atıf:
Cite this article as:
Yilmaz U, Kir S, Inci A, Aksoy N,
Demirtas G, Gul S, Maden U.
Analysis of Sociodemographic
Characteristics of Early and
Advanced Stage Chronic Kidney
Disease Patients

Akd Tıp D / 2021; 7(3):424-429

Ustun YILMAZ
ORCID ID: 0000-0003-0237-0278
Selin KIR
ORCID ID: 0000-0001-7792-0882
Ayca INCI
ORCID ID: 0000-0002-7894-8913
Nilgun AKSOY
ORCID ID: 0000-0001-9038-3348
Gulay DEMIRTAS
ORCID ID: 0000-0001-5273-1452
Semih GUL
ORCID ID: 0000-0003-3494-3931
Ummuhan MADEN
ORCID ID: 0000-0002-9195-6918

ABSTRACT

Objective

It was aimed to investigate sociodemographic features of chronic kidney disease patients, analyze difference of these features in early and advanced stage patients, find correctable factors that may change course of disease and to find new solutions in treatment and follow-up.

Material and Methods

Predialysis 345 patients diagnosed with chronic kidney disease were retrospectively analyzed. Patients were divided into early and advanced stages. Patients with Glomerular Filtration Rate 30 ml / min were defined as early stage, <30 ml / min as advanced stage. Sociodemographic and disease related features and laboratory parameters of patients were analyzed.

Results

In the study, 243 (70.4%) patients were early stage and 102 (29.6%) patients were advanced stage chronic kidney patients. Most common etiologies are; In early stage, chronic glomerulonephritis (30.5%), diabetes mellitus (16.9%) and genetic diseases (16.0%), while at advanced stage, diabetes mellitus (34.3%), hypertension (16.7%) and unknown causes (15.7%) were. Alcohol use was higher and regular drug use was lower in early stage patients compared to advanced stage patients. In advanced stage patients, habit of exercising regularly was found to be low compared to early stage patients.

Conclusion

It was found that there were correctable factors that could change course of disease. These factors differed in early and advanced stages. It is necessary to be aware of these factors for slower progressive of disease and take precautions for correctable factors from early stages of disease.

Keywords

Chronic kidney disease, Correctable factors, Sociodemographic feature

ÖZ

Amaç

Kronik böbrek hastalarının sosyodemografik özelliklerinin araştırılması, erken ve ileri evre hastalarda bu özelliklerin farklılığının incelenmesi, hastalığın seyrini değiştirebilecek düzeltilebilir faktörlerin saptanması, tedavi ve takipte yeni çözümler bulunması amaçlanmıştır.

Gereç ve Yöntemler

Kronik böbrek hastalığı tanısı almış, prediyaliz 345 hasta retrospektif olarak incelenmiştir. Hastalar erken ve ileri evrelere ayrılmıştır. Glomerüler Filtrasyon Hızı ≥ 30 ml / dk olan hastalar erken, <30 ml / dk olan hastalar ileri evre olarak tanımlandı. Hastaların sosyodemografik ve hastalıkla ilişkili özellikleri ile laboratuvar parametreleri analiz edilmiştir.

Bulgular

Çalışmada 243 (%70,4) hasta erken evre (Evre 1-2-3), 102 (%29,6) hasta ileri evre (Evre 4-5) kronik böbrek hastalığı olarak saptandı. En sık görülen etyolojiler; erken evrede kronik glomerulonefrit (%30,5), diabetes mellitus (%16,9) ve genetik hastalıklar (%16,0) iken ileri evrede diabetes mellitus (%34,3), hipertansiyon (%16,7) ve nedeni bilinmeyen (%15,7) sebepler idi. Erken evre hastalarda ileri evre hastalara kıyasla alkol kullanımının daha yüksek, düzenli ilaç kullanımının daha düşük olduğu saptandı. İleri evre hastalarda ise düzenli egzersiz yapma alışkanlığı erken evre hastalara göre düşük bulundu.

Sonuç

Hastalığın seyrini değiştirebilecek düzeltilebilir faktörlerin olduğu, erken ve ileri evrelerde bunların farklılık gösterebildiği görülmüştür. Daha yavaş progresyonlu hastalık seyri için bu faktörlerin farkında olunması ve hastalığın erken evrelerinden itibaren takip edilmesi önemlidir.

Anahtar kelimeler

Kronik Böbrek Hastalığı, Düzeltilebilir faktörler, Sosyodemografik özellikler

Ethical approval: This study was approved by the Ethics Committee of the Scientific Research Ethics Committee of Health Sciences University Antalya Training and Research Hospital for the study (Decision number: 10/22, decision date: 03/07/2020).

INTRODUCTION

Chronic kidney disease (CKD) is an important public health problem that has become almost epidemic in our country and in the world (1,2,3). CKD is a progressive disease and has an important place in terms of both morbidity and mortality for patients. It is a disease that places a great burden on the health budget as well as being common. The most common etiology is diabetes mellitus (DM), followed by hypertension (HT). Apart from these, glomerulonephritis, kidney stones, genetic kidney diseases and systemic diseases with kidney involve-

ment take place in etiology (4,5,6). In the course of CKD; serious complications develop due to many reasons such as problems related to uremic toxins, endocrinological disorders, metabolic disorders and anemia, and patients quality of life, social life and productivity are severely damaged.

While CKD and its complications harm patients in many respects, low socioeconomic factors, bad habits, and negative living conditions have also been found to facilitate the progression of the disease (7,8). Therefore, it is important to examine the sociodemographic characteristics of the patients. In our study, it was aimed to analyze the sociodemographic characteristics, laboratory parameters and etiologies of early and advanced CKD patients followed in our clinic and to identify correctable factors that may change the course of the disease.

MATERIALS AND METHODS

Patients

This study was conducted by retrospectively examining the files of 345 patients diagnosed with CKD who were followed up at the Nephrology clinic at the Health Sciences University Antalya Training and Research Hospital. Glomerular filtration rates (GFR) of patients; It was calculated using the formula 'Chronic kidney disease epidemiology collaboration' (CKD-EPI) (9). According to this calculation; Stage 1-2-3 patients with GFR ≥ 30 ml / min are defined as early stage, and Stage 4-5 patients with GFR <30 ml / min are defined as advanced stage. Being diagnosed with CKD for at least 3 months, no history of kidney transplant and being over 18 years old were determined as the inclusion criteria for the study. Those excluded from these parameters were not included in the study. Approval was obtained from the ethics committee of Health Sciences University Antalya Training and Research Hospital for the study (Decision number: 10/22, decision date: 03/07/2020). The study was conducted in accordance with the ethical standards defined in the 1964 Helsinki declaration.

Statistical analysis

SPSS (Statistical Package for Social Science) for Windows 23.0 was used for statistical analysis of the data obtained in the study. Descriptive analyzes were given with frequency distribution, mean and standard deviation. Initially, the normal distribution of data was tested using the Kolmogorov-Smirnov test. As a result of the tests performed, the properties of normally distributed data were analyzed by parametric tests, and those that did not, were analyzed by non-parametric tests. Difference of two independent averages of the groups was made by using "student t test". In cases where parametric test assumptions were not provided, non-parametric alternative of this test, "Mann-Whitney U" was used. Pearson and Spearman correlation analysis was performed to determine the relationship between dependent quantitative variables. Statistical significance was accepted for $P < 0.05$ values.

RESULTS

The study was performed in 345 patients with CKD who were at the pre-dialysis period at the Health Sciences University Antalya Training and Research Hospital. In our study, 243 (70.4%) patients were found to be in the early stage (Stage 1-2-3), and 102 (29.6%) patients were in the advanced stage (Stage 4-5). Sociodemographic characteristics of early and advanced stage patients are shown in Table I.

Table I. Distribution and comparison of socio-demographic and physical characteristics of patients by groups

	Stage 1-2-3 (n=243)	Stage 4-5 (n=102)		p
Age (Mean ± SD)	49.06±14.52	59.53±13.95	Z=-6.028	0.000*
Body mass index (Mean ± SD)	28.23±5.34	27.90±4.99	Z=-0.280	0.779*
Gender (%)				
Female	115(%47.33)	38(%37.25)	$\chi^2 = 2.95$	0.086**
Male	128(%52.67)	64(%62.75)		
Education status			$\chi^2 = 8.28$	0.082**
Illiterate	17 (%6.99)	14(%13.73)		
Elementary school	110(%45.27)	53(%51.96)		
Junior high school graduate	27(%11.11)	11(%10.78)		
High school graduate	57(%23.46)	14(%13.73)		
Graduated from a university	32(%13.17)	10(%9.80)		
Marital status			$\chi^2 = 0.00$	0.986**
Married	188(%77.37)	79(%77.45)		
Single	55(%22.63)	23(%22.55)		
Home partner status			$\chi^2 = 7.348$	0.119**
Lives with partner	68(%27.98)	31 (%30.39)		
Lives with partner and children	120(%49.38)	49 (%48.04)		
Lives alone	18(%7.41)	7 (%6.86)		
Lives with children	12(%4.94)	11 (%10.78)		
Lives with parents	25(%10.29)	4 (%3.92)		
Settlement unit				
City	154 (%63.37)	56(%54.90)	$\chi^2 = 2.451$	0.294**
District	60 (%24.69)	33(%32.35)		
Village	29 (%11.93)	13(%12.75)		
Home type				
Apartment	162(%66.67)	64(%62.75)	$\chi^2 = 0.489$	0.484**
Detached house + other	81(%33.33)	38(%37.25)		
Home ownership status				
Rent	76(%31.28)	29(%28.43)	$\chi^2 = 1.640$	0.440**
Belongs to the family	28(%11.52)	8(%7.84)		
Own home	139(%57.20)	65(%63.73)		
Working status				
Bodily work	72 (%29.6)	13 (%12.7)	$\chi^2 = 13.998$	0.003**
Retired	51 (%21.0)	43 (%42.2)		
Not working	100 (%41.2)	36 (%35.3)		
Desk job	20 (%8.2)	10 (%9.8)		
Income level status				
Under minimum wage	57(%23.46)	32(%31.37)	$\chi^2 = 3.272$	0.195**
Minimum wage	118(%48.56)	49(%48.04)		
Above minimum wage	68(%27.98)	21(%20.59)		
Smoking status				
Smoking	67(%27.57)	19(%18.62)	$\chi^2 = 5.637$	0.060**
Never smoked	133(%54.73)	55(%53.92)		
Quit smoking	43(%17.70)	28(%27.45)		
Alcohol using status				
Yes	39(%16.05)	8(%7.84)	$\chi^2 = 4.112$	0.043
No	204(%83.95)	94(%92.16)		
Surgical history				
Yes	131(%53.91)	64(%62.75)	$\chi^2 = 2.282$	0.131**
No	112(%46.09)	38(%37.25)		
Duration of kidney disease follow-up				
0-6 months	115(%47.33)	27(%26.47)	$\chi^2 = 19.971$	0.000**
6-12 months	15(%6.17)	8(%7.84)		
1-6 years	56(%23.05)	46(%45.10)		
6 years and over	57(%23.46)	21(%20.59)		
Use of a walking aid device				
Yes	11(%4.53)	13(%12.75)	$\chi^2 = 7.497$	0.006**
No	232(%95.47)	89(%87.25)		
Regular exercise status				
Yes	108(%44.44)	29(%28.43)	$\chi^2 = 7.695$	0.006**
No	135(%55.56)	73(%71.57)		
Family history of kidney disease				
Yes	90(%37.04)	28(%27.45)	$\chi^2 = 2.934$	0.087**
No	153(%62.96)	74(%72.55)		
Regular drug use status				
Yes	214(%88.07)	99(%97.06)	$\chi^2 = 6.905$	0.009**
No	29(%11.93)	3(%2.94)		

*Mann-Whitney U test , ** Chi-square test

The analysis showed that the average age of advanced stage CKD patients was higher ($Z = -6.028$; $p = .000$). Gender and body mass index (BMI) parameters were not statistically different in early and advanced stage CKD patients ($p > 0.05$). No statistically significant difference was found between the stages in terms of education status, marital status, home partner status, settlement unit, home type, home ownership status, income level status, smoking status, surgical history, family history of kidney disease ($p > 0.05$). While the rate of physical work was high in the early stage patients, the rate of retirement was found to be significantly higher in the advanced stage patients ($\chi^2 = 13.998$; $p = 0.003$). Alcohol use was higher ($\chi^2 = 4.112$; $p = 0.043$) and regular drug use was lower ($\chi^2 = 6.905$; $p = 0.009$) in the early stage compared to the advanced stage. In advanced stage patients, the habit of exercising regularly was lower compared to the early stage ($\chi^2 = 7.695$; $p = 0.006$), while use of a walking aid device ($\chi^2 = 7.497$; $p = 0.006$) was higher. Duration of kidney disease follow-up was determined to be concentrated between 0-6 months in the early stage and 1-6 years in the advanced stage ($\chi^2 = 19.971$; $p = 0.000$).

Table II. Distribution of kidney disease causes by disease stage

Kidney Disease Cause	Stage 1-2-3 (n=243)	Stage 4-5 (n=102)
Chronic glomerulonephritis	74(%30.5)	8(%7.8)
Congenital structural disorders	2(%0.8)	-
Stone or obstruction	8(%3.3)	6(%5.9)
Hypertension	26(%10.7)	17(%16.7)
Genetic diseases	39(%16.0)	9(%8.8)
Diabetes mellitus	41(%16.9)	35(%34.3)
Systemic diseases	4(%1.6)	4(%3.9)
Urinary tract infection	1(%0.4)	-
Other reasons	13(%5.3)	7(%6.9)
Reason unknown	35(%14.4)	16(%15.7)

The distribution of causes CKD by early and advanced groups is shown in Table II. In etiology, chronic glomerulonephritis (30.5%), DM (16.9%) and genetic diseases (16.0%) were the most common causes of CKD in the early stage, while DM (34.3%), HT (16.7%) and unknown cause (15.7%) were the most common causes in advanced stage.

When early and advanced stage CKD groups were compared in terms of laboratory parameters; creatinine ($Z = -14.147$; $P = 0.000$), phosphorus ($Z = -5.808$; $P = 0.000$), uric acid ($Z = -2.117$; $P = 0.034$), parathormone ($Z = -9.927$; $P = 0.000$), ferritin ($Z = -4.761$; $P = 0.000$) and spot urine protein ($Z = -3.667$; $P = 0.000$) averages were found to be significantly higher in the advanced stage. Calcium ($Z = -2.137$; $P = 0.033$), GFR ($Z = -14.481$; $P = 0.000$) and hemoglobin ($F = 0.078$; $P = 0.000$) averages were found at a higher level in the early stage. It was found that there was no significant difference between the groups in terms of the average of albumin ($Z = -1.333$; $P = 0.183$) level (Table III).

Table III. Comparison of the laboratory parameters of the patients according to the stages

Laboratory Parameters	Stage 1-2-3 (n=243) (Mean ± SD)	Stage 4-5 (n=102) (Mean ± SD)	
Creatinine (mg / dl)	1.34±0.43	2.93±0.80	Z=-14.147 P=0.000*
GFR (ml / min / 1.73m2)	62.53±26.69	21.85±6.61	Z=-14.481 P=0.000*
Calcium (mg / dl)	9.76±5.15	9.27±0.70	Z=-2.137 P=0.033*
Phosphorus (mg / dl)	3.67±2.07	4.09±0.83	Z=-5.808 P=0.000*
Albumin (g / dl)	4.30±2.46	4.13±0.47	Z=-1.333 P=0.183*
Uric acid (mg / dl)	6.72±4.62	6.97±1.69	Z=-2.117 P=0.034*
Hemoglobin (g / dl)	13.08±1.89	11.90±1.78	F=0.078 P=0.000**
Parathormon (pg / ml)	72.16±61.85	188.32±124.92	Z=-9.927 P=0.000*
Ferritin (mcg / l)	69.55±76.38	108.99±103.99	Z=-4.761 P=0.000*
Spot urine protein (mg / g)	1.98±3.28	2.52±2.97	Z=-3.667 P=0.000*

*Mann-Whitney U testi

**Independent Samples T Test

When correlation analysis between CKD disease stage and laboratory parameters is made; creatinine ($r = 0.763$; $p < 0.001$), phosphorus ($r = 0.313$; $p < 0.001$), uric acid ($r = 0.114$; $p = 0.034$), parathormone ($r = 0.535$; $p < 0.001$), ferritin ($r = 0.257$; $p < 0.001$) and spot urine protein ($r = 0.198$; $p < 0.001$) statistically significant positive correlation was found. Mean levels of hemoglobin ($r = -0.279$; $p < 0.001$), GFR ($R = -0.781$; $p < 0.001$) and calcium ($R = -0.115$; $p < 0.001$) were statistically significant negative correlations between the disease stage.

DISCUSSION

It is a disease that should be taken seriously in terms of the burden of CKD costs on the countries and the increasing frequency of them. Therefore, it seems important to take precautions according to the results that may change the course of the disease as a result of examining the life styles, social status, habits, kidney disease causes, and biochemical disorders related to the disease. In our study, we examined the sociodemographic features and laboratory parameters of our patients diagnosed with predialysis CKD, which were followed up in our clinic, for early and advanced stages.

In our study, we found that the average age of patients with advanced stage was higher. This finding is in line with the information that advancing age is a risk factor for chronic kidney diseases (10,11).

It has been reported that weight gain increases the risk of CKD, even if it falls within the normal BMI limits (12). In addition, in patients who have not started renal replacement therapy, it was determined that oral intake began to deteriorate when GFR decreased below 50 ml/min, impaired renal function increased and provided the basis for malnutrition (13). In our study, no difference was found between BMI according to disease stages.

In the CREDIT study conducted in our country, female gender was found to be a risk factor for CKD (3). In another study,

male gender (70.7%) was found in the majority (14). In our study, no difference was found in terms of gender in both early and advanced stages.

Increasing the level of education; Adopting a lifestyle that will prevent the development of many systemic diseases known to be included in the CKD etiology is an important factor in terms of having disease awareness and compliance with treatment. In some studies supporting this situation, low level of education has been reported to be a risk factor for CKD (10). In our study, no difference in education level was found between early and advanced stage CKD patients.

In a study conducted by Inci et al., It was stated that parameters such as the patient's home partner status, settlement unit, home type, home ownership status, income level did not differ between the disease stages (15). Similarly, in our study, no difference was found between these parameters and disease stages.

CKD is a disease that leads to significant loss of labor and associated decline in income. In our study, no difference was observed in the income levels of patients according to the stages. The reason for this may be that the income level was offset due to the high rate of physical work of early stage patients and the high number of retirements in advanced stage patients.

In a study, smoking and alcohol use have been reported to accelerate CKD development and disease progression (16). In our study, we found that while the rate of those who smoke in the early stages is high, the percentage of those who quit smoking in the advanced stage is high, but there is no statistical difference between the stages. We found that alcohol use was significantly higher in the early stages. This may be associated with a decrease in alcohol use behavior due to low awareness of the disease at the early stage or due to increased anxiety with disease progression.

In our study, it was found that after the diagnosis of CKD, the duration of kidney disease follow-up was concentrated in 0-6 months in the early stage patients, and in 1-6 years in the advanced stage patients. Long follow-up in advanced stage patients is an expected finding.

In our study, use of a walking aid device was significantly more frequent in advanced stage patients. This need is thought to occur due to the advanced complications of CKD disease and additional medical problems brought about by the increase in age.

It has been reported that a moderate exercise is recommended in patients with CKD, since it reduces bone and joint diseases and cardiovascular risk caused by age (17). In our study, regular exercise was found to be higher in the early stage, while the rate of regular exercise decreased as the disease stage progressed.

Family history of kidney disease was similarly detected in early and advanced stage patients. This is an expected finding. When studies on regular drug use were examined; It has been reported that the level of importance that individuals attach to their illness and not knowing the drug benefits due to lack of education affect adherence to treatment (18,19). In our study, it was found that regular drug use was higher in advanced

CKD patients. The reason for this situation may be due to factors such as those with early stage disease not being able to grasp the importance of the disease or not knowing the benefit of the drug for the disease.

Among the causes leading to chronic kidney failure, DM, HT and glomerulonephritis was reported as the first three causes (4-6). When the causes leading to kidney disease were examined in our study; the three most common etiologies; while chronic glomerulonephritis was 30.5%, DM 16.9% and genetic diseases were 16.0% in early stage patients, DM was 34.3%, HT 16.7% and unknown reason 15.7% in advanced stage patients. Glomerulonephritis and genetic-induced kidney diseases may have caused an early and advanced etiological factor difference due to their early follow-up in our clinic.

In our study, it was observed that the level of creatinine, phosphorus, parathormone, uric acid, ferritin and spot urine proeinuria was higher in advanced stage patients and positively correlated with the disease stage. GFR, calcium, hemoglobin levels were found to be low in advanced stage and negatively correlated with disease stage. Although albumin levels were found to be relatively lower in advanced stage, there was no statistically difference between stages. These findings were found to be compatible with the literature (1,20).

Due to the fact that some patients received a diagnosis in an external center and then began to follow up to our clinic, the inability to reach all of the GFR values at the time of diagnosis, the fact that it is single-center and cross-sectional, can be shown among the limitations of our study.

CONCLUSIONS

In our study, the three most common etiologies in CKD are; while chronic glomerulonephritis, DM and genetic diseases were detected in early stage patients, DM, HT and unknown causes were found in advanced stage patients. It was found that the average age of patients with advanced stage, the use of a walking aid device was higher and the regular exercise rate was lower in these patients. In early stage patients, it was observed that alcohol use was high and regular drug use was lower. These findings indicate that there are correctable causes that can cause this disease to progress. These factors differed in early and advanced stages. It is necessary to be aware of these factors for slower progressive of the disease and take precautions for correctable factors from early stages of disease. Our study and new studies to be conducted in a multi-centered manner with more patients will illuminate this issue more and will guide the follow-up of this disease.

Competing interests: The authors declare that they have no competing interest.

Financial Disclosure: There are no financial supports.

Ethical approval: This study was approved by the Ethics Committee of the Scientific Research Ethics Committee of Health Sciences University Antalya Training and Research Hospital for the study (Decision number: 10/22, decision date: 03/07/2020).

REFERENCES

1. Levey AS, Coresh J. Chronic kidney disease. *Lancet* 2012; 14;379(9811):165-80.
2. Süleymanlar G. Kronik Böbrek Hastalığı ve Yetmezliği: Tanımı, Evreleri ve Epidemiyolojisi. *Türkiye Klin J Int Med Sci* 2007; 3(38).
3. Süleymanlar G, Utaş C, Arinsoy T, Ateş K, Altun B, Altıparmak MR, Ecder T, Yılmaz ME, Çamsarı T, Başçı A, Odabas AR, Serdengeçti K. A population-based survey of Chronic Renal Disease In Turkey--the CREDIT study. *Nephrol Dial Transplant* 2011; 26(6):1862-71.
4. National Kidney Foundation. K/DOQI Clinical Practice Guidelines for Chronic Kidney Disease: Evaluation, Classification and Stratification. *Am J Kidney Dis* 2002; 39(2 Suppl 1):S1-266.
5. Hart PD, Bakris GL. Hypertensive nephropathy: prevention and treatment recommendations. *Expert Opin Pharmacother* 2010; 11(16):2675-86.
6. Jha V, Garcia-Garcia G, Iseki K, Li Z, Naicker S, Plattner B, Saran R, Yee-Moon Wang A, Yang CW. Chronic kidney disease: global dimension and perspectives. *Lancet*. 2013; 382(9888):260-72.
7. Naicker S. End-stage renal disease in sub-Saharan and South Africa. *Kidney Int Suppl* 2003; (83):S119-22.
8. Zeng X, Liu J, Tao S, Hong HG, Li Y, Fu P. Associations between socioeconomic status and chronic kidney disease: a meta-analysis. *J Epidemiol Community Health* 2018; 72(4):270-279.
9. Levey AS, Stevens LA, Schmid CH, Zhang YL, Castro AF 3rd, Feldman HI, Kusek JW, Eggers P, Van Lente F, Greene T, Coresh J. A new equation to estimate glomerular filtration rate. *Ann Intern Med* 2009; 150(9):604-12.
10. Bartmańska M, Więcek A. Chronic kidney disease and the aging population. *G Ital Nefrol* 2016; 33 Suppl 66:33.S66.11.
11. Levey AS, Coresh J, Balk E, Kausz AT, Levin A, Steffes MW, Hogg RJ, Perrone RD, Lau J, Eknoyan G. National Kidney Foundation practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Ann Intern Med* 2003; 139(2):137-47.
12. Ryu S, Chang Y, Woo HY, Kim SG, Kim DI, Kim WS, Suh BS, Choi NK, Lee JT. Changes in body weight predict CKD in healthy men. *J Am Soc Nephrol* 2008; 19(9):1798-805.
13. Mehrotra R, Kopple JD. Nutritional management of maintenance dialysis patients: why aren't we doing better? *Annu Rev Nutr* 2001; 21: 343-79.
14. Ladi-Akinyemi TW, Ajayi I. Risk factors for chronic kidney disease among patients at oabisi onabanjo university teaching hospital in sagamu, Nigeria: A retrospective cohort study. *Malawi Med J* 2017; 29(2):166-170.
15. İnci A, Çoban M, Sarıkaya M, Maden Ü. Prediyaliz Kronik Böbrek Hastalarında Kesitsel Bir Çalışma: Yaşam Kalitesi ve Etkileyen Faktörler. *Kocatepe Tıp Dergisi* 2017; 18:130-135
16. Shankar A, Klein R, Klein BE. The association among

- smoking, heavy drinking, and chronic kidney disease. *Am J Epidemiol* 2006; 164(3):263-71.
17. Taş D, Akyol A. Egzersiz ve Kronik Böbrek Yetmezliği. *Türk Nefroloji, Diyaliz ve Transplantasyon Hemşireleri Derneği Nefroloji Hemşireliği Dergisi* 2017;1 (12).
 18. Kamran A, Sadeghieh Ahari S, Biria M, Malepour A, Heydari H. Determinants of patient's adherence to hyper tension medications: application of health belief model among rural patients. *Ann Med Health Sci Res* 2014;4(6):922-7.
 19. Yakar B, Demir M. Kronik Böbrek Yetmezliği Olan Hastalarda Antihipertansif İlaç Tedavisine Uyum Oranları ve Etkileyen Faktörlerin Araştırılması. *Dicle Tıp Dergisi / Dicle Med J* 2019; 46 (4) : 685 - 696.
 20. Akpolat T, Utaş C, Süleymanlar G. *Nefroloji El Kitabı*, 4.baskı. İstanbul: Nobel Tıp Kitabevi, 2007 , sayfa 283-323.