PAPER DETAILS

TITLE: Draft Genome Sequence of Staphylococcus felis HARRANVET Strain

AUTHORS: Özkan ASLANTAS, Oktay KESKIN, Kaan BÜYÜKALTAY, Ayfer GÜLLÜ YÜCETEPE

PAGES: 78-81

ORIGINAL PDF URL: https://dergipark.org.tr/tr/download/article-file/1594698



Draft Genome Sequence of Staphylococcus felis HARRANVET Strain

Özkan Aslantaş¹, Oktay Keskin², Kaan Büyükaltay³, Ayfer Güllü Yücetepe²

¹Department of Microbiology, Faculty of Veterinary Medicine, Hatay Mustafa Kemal University, Hatay/TURKEY ²Department of Microbiology, Faculty of Veterinary Medicine, Harran University, Şanlıurfa/TURKEY ³Bionformatics Institute, Middle East Technical University, Ankara/TURKEY

Key Words: cat Staphylococcus felis whole-genome sequencing

Anahtar Kelimeler: kedi *Staphylococcus felis* tüm genom dizileme

 Received
 : 22.02.2021

 Accepted
 : 07.04.2021

 Published Online
 : 31.08.2021

 Article Code
 : 884932

Correspondence: Ö. ASLANTAŞ (aslantas@mku.edu.tr)

ORCID:	
Ö. ASLANTAŞ	: 0000-0003-0407-8633
O. KESKİN	: 0000-0002-5977-7872
K. BÜYÜKALTAY	: 0000-0001-6074-8940
A. GÜLLÜ-YÜCETEPE	: 0000-0002-9842-3305

ABSTRACT

Staphylococcus felis is one of staphylococci residing on skin of cats, and has been increasingly reported as a potential facultative pathogen. A recently isolated *Staphylococcus felis*(named as *S. felis* HARRANVET strain) from necrotizing fasciitis case of a cat was characterized using whole-genome sequencing (WGS), and assembled genome was *in silico* screened for putative virulence and antimicrobial resistance genes. The isolate was pan-susceptible to all antimicrobials tested, indicated absence of resistance genes. A limited number of virulence genes associated with adhesion was found. For the first time, in Turkey, the draft genome of *Staphylococcus felis* was obtained and made publicly accessible.

Staphylococcus felis HARRANVET Suşunun Taslak Genom Dizisi

ÖΖ

Staphylococcus felis, kedilerin derisine kolonize olan stafilokok türlerinden biridir ve potansiyel bir fakültatif patojen olarak giderek daha fazla rapor edilmektedir. Yakın zamanda bir kedinin nekrotizan fasiit vakasından izole edilen *Staphylococcus felis* (*S. felis* HARRANVET suşu olarak adlandırılan) tüm genom dizileme (WGS) ile karakterize edildi ve birleştirilmiş genom virülans ile antimikrobiyal direnç genleri yönünden *in silico* olarak analiz edildi. İzolat incelenen tüm antimikrobiyallere karşı duyarlı bulundu. Adhezyon ile ilişkili sınırlı sayıda virülans geni tespit edildi. Türkiye'de ilk kez, shotgun dizileme teknolojisi ile *Staphylococcus felis*'in genom sekans taslağı elde edildi ve erişime açık hale getirildi.

INTRODUCTION

Although *Staphylococcus* spp. are known as part of natural microbiota of the skin, mucous membranes, alimentary and urogenital tracts of a wide range of mammals, they can also lead to a variety of clinical manifestations in humans and animals (2, 3, 12). *Staphylococcus felis* (*S. felis*) is one of commensal pathogen, regardless of health status of cats (9, 12). *S. felis* infections were associated with urinary tract infections, pyoderma, postoperative wound infections and otitis (11, 12). While the ability of *S. felis* isolates to produce urease is accepted an indication of potential urinary tract pathogen (11), their ability to form biofilm and produce proteolytic enzyme synthesis is accepted an indication of opportunistic skin pathogen (9).

Limited number of *S. felis* draft genome and complete genome are available in databases to date. Also, there is no study characterizing *S. felis* in Turkey. Therefore, we sequenced the complete genome of *S. felis* HARRANVET strain and investigated phenotypic and genotypic antimicrobial resistance profile.

MATERIAL and METHODS

Isolation and Identification of S. felis

The isolate was recovered from necrotizing fasciitis on a cat's head in Microbiology Laboratory, Faculty of Veterinary Medicine, Harran University. The isolate was preliminarily identified using phenotypic tests (Gram staining, catalase, co-agulase testing) (6) and confirmed by MALDI-TOF MS and BLASTn analysis of 16S rRNA sequencing using universal primers (27F – 1492R). The tube coagulase and slide agglutination tests were conducted using feline and rabbit plasma to determine the ability of the isolate to form clot formation (8).

Antimicrobial Susceptibility Testing

Antimicrobial susceptibility of the isolate was determined using disc diffusion method in accordance with Clinical Laboratory Standards Institute (CLSI) criteria (6), and the following discs were used: oxacillin (1 μ g), penicillin (10 U), amoxicillin-clavulanic acid (30 μ g), trimethoprim-sulfamethoxazole (25 μ g), ciprofloxacin (5 μ g), clindamycin (2 μ g), gentamicin (10 μ g), chloramphenicol (30 μ g), rifampicin (5 μ g), tetracycline

Bu makaleye atıfta bulunmak için/ To cite this article:

Aslantaş Ö, Keskin O, Büyükaltay K, Yücetepe AG, Draft Genome Sequence of Staphylococcus felis HARRANVET Strain. MAE Vet Fak Derg. 2021; 6 (2):78-81.

(30 µg), linezolid (30 µg), erythromycin (15), tigecycline (15 µg), vancomycin (30 µg), amikacin (30 µg), fusidic acid (10 µg), kanamycin (30 µg), streptomycin (10 µg), quinopristin-dalphopristin (15 µg), ampicillin (10 µg) and tobramycin (10 µg). *S. aureus* ATCC 29213 was used as a positive control for coagulase and clumping factor tests and antimicrobial susceptibility testing as well.

Coagulase Testing

The tube coagulase test was conducted using both feline and rabbit plasma to determine the ability of the isolate to form clot formation. The isolate was accepted coagulasepositive if an immobile clot formed at the bottom of the tube after 4 hours incubation at 37°C or after overnight incubation at room temperature [6]. In addition, the slide agglutination test was used to assess the presence of clumping factor using rabbit and feline plasma. *S. aureus* ATCC 29213 was used as a positive control for coagulase and clumping factor test.

Whole-genome sequencing

The genome of *S. felis* HARRANVET strain was sequenced using the Illumina Novaseq platform with pair-end reads (PE 150) following the manufacturer's instructions (Illumina, San Diego, USA). Preprocessed reads were assembled using SPAdes Genome Assembler version 3.14.1 (1). The genome annotation was carried out with the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) (http://www.ncbi.nlm. nih.gov/books/NBK174280/). The assembled genome was deposited at NCBI under accession no. JAEDAQ000000000. The sequenced reads were also assembled to *S. felis* ATTC 49168 using Bowtie2 alignment tool (10).

Screening of antimicrobial resistance and virulence genes

The assembled genome was screened *in silico* for acquired antimicrobial resistance and virulence genes, using openaccess bioinformatics website, the Center for Genomic Epidemiology (CGE) (https://cge.cbs.dtu.dk//services/). ResFinder v4.1 (15) and VirulenceFinder v2.0 (7) were used to identify acquired antimicrobial resistance genes and virulence genes, respectively. Additional virulence factors described for staphylococci, but not included in VirulenceFinder, were also



Figure 1. Maximum-likelihood phylogenetic tree based on 31 S. felis genomes present in PATRIC databases. Geographic origin of the isolates is highlighted in different colours. GenBank accession numbers of each genome are given in parentheses.

investigated using BLASTn tool of NCBI.

Phylogenetic analysis

Whole-genome sequences of 30 *S. felis* isolates of cats and one isolate with unknown origin from different countries retrieved from the PATRIC *S. felis* database for phylogenetic analysis (4). GenBank accession numbers of genomes used in the phylogenetic comparison was seen in Figure 1. The tree was generated with maximum likelihood estimation methods. The evolutionary relationship between the *S. felis* strains was inferred by the aligned core-genomes using the program RAxML in PATRIC

RESULTS

S. felis isolate

The isolate was identified as *S. felis* by MALDI-TOF MS and BLASTn analysis of 16S rRNA sequencing. The isolate was named *S. felis* HARRANVET strain.

Antimicrobial resistance

The isolate was found to be susceptible to all tested antimicrobials.

Virulence factors

In silico analysis of the assembled genome revealed absence of resistance genes from ResFinder database. The isolate did not harbor any virulence gene homologues deposited in the VirulenceFinder database. However, the isolate harbored putative virulence gene homologues for fibronectin/fibrinogen-binding protein (fnbB) and autolysin/adhesion (atl).

Genomic analysis

The sequenced reads were assembled into a draft genome consisting 338 contigs, with the total length of 2,618,484 bp. The average G + C content was 34.86147. The assembled genome encoded 2,513 proteins and 54 transfer RNAs genes based on the annotation done at NCBI Prokaryotic Genome Annotation Pipeline (PGAP). On average, 88.73% of the sequence reads mapped to the 2,479,423 bp reference genome, *S. felis* ATCC 49168.

Phylogenetic analysis

Phylogenetic analysis based on 31 *S. felis* genomes present in the PATRIC database revealed two main clusters and several subclusters (Figure 1). The phylogenetic tree indicated that the *S. felis* HARRANVET strain was closely related with the isolates from Australian isolates.

DISCUSSION

The skin and mucous membranes of cats are colonized with staphylococcus species, with a dominance of coagulase-negative staphylococci (CoNS). Different interactions between staphylococci and the host may lead to development of symbiotic, commensal and opportunistic infections. Under favourable circumstances, opportunistic infections can occur and damage host tissues and organs (3).

Staphylococci can produce several virulence factors playing important role in its pathogenesis. The isolate did not show coagulase activity, which is one of the virulence factors produced by some Staphylococcus species. In a comprehensive study, Worthing et al. (14) detected coagulase activity only in an isolate (1/27) with cat plasma, while they found the rest of the isolates being negative with both feline and rabbit plasma. In terms of other putative virulence factors investigated, the isolate harboured only fibronectin/fibrinogen-binding protein (*fnbB*) and autolysin (atl) gene homologues, which are implicated in adhesion to host cells. The *fnb*B is a cell-bound protein that allows staphylococci to bind fibronectin, fibrinogen and elastin found on host cells, and is one of microbial cell surface protein receptors designated MSCRAMMs (5). The atl is a surface expressed hydrolase with diverse functions such as binding host matrices, involvement in biofilm formation and releasing cytoplasmic proteins from staphylococcal cell wall (13). Despite being searched for other virulence factors, other putative virulence genes were not determined in contrast to findings of Worthing et al (14), who detected genes responsible for both adhesion and toxin production. This drawback could be overcomed by using long-read sequence technology in future studies to elucidate other putative virulence determinants.

The isolate was susceptible to all the antimicrobials tested in the study. In contrast to other CoNS isolated from cats, it has been reported that *S. felis* strains had low level phenotypic antimicrobial resistance (11, 12, 14). Possible reasons for low or no antimicrobial resistance observed among clinical *S. felis* isolates are explained in two ways: (i) fitness-cost of resistance can lead to reduced growth rate, competitive ability and virulence, compared with susceptible strains, (ii) the *S. felis* genome could have low plasticity to acquire a maintain mobile genetic elements (MGEs) conferring resistance (13).

As it can be seen in phylogenetic analysis (Figure), *S. felis* HARRANVET strain was placed in a cluster, closer to Australian isolates. The phylogenetic tree also showed high genetic diversity indicating host-adapted variations in the isolates.

In conclusion, for the first time, the draft genome of *S. felis* HARRANVET strain was obtained and made publicly available.

DECLARATIONS

Ethics Approval

This study does not present any ethical concerns.

Conflict of Interest

The authors declare that they have no competing interests.

Author Contribution

Idea, concept and design: O Aslantaş, O Keskin, K Büyükaltay

Data collection and analysis: O Aslantaş, O Keskin, K Büyükaltay, A Güllü Yücetepe

Drafting of the manuscript: O Aslantaş

Critical review: O Aslantaş, O Keskin, K Büyükaltay, A Güllü Yücetepe

Data Availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Financial Support

This research received no grant from any funding agency/sector

REFERENCES

1. Becker K, Heilmann C, Peters G. Coagulase-negative staphylococci. Clin Microbiol Rev, 2014; 27(4): 870-926.

2. Qekwana DN, Sebola D, Oguttu JW, Odoi A. Antimicrobial resistance patterns of Staphylococcus species isolated from cats presented at the veterinary academic hospital in South Africa. BMC Vet Res, 2017; 13:286.

3. Bierowiec K, Korzeniowska-Kowal A, Wzorek A, Rypuła K, Gamian A. Prevalence of *Staphylococcus* species colonization in healthy and sick cats. Biomed Res Int, 2019; 2019:4360525.

4. Kwaszewska A, Lisiecki P, Szemraj M, Szewczyk EM. Animal Staphylococcus felis with the potential to infect human skin. Med Dosw Mikrobiol, 2015; 67:69-78.

5. Litster A, Moss SM, Honnery M, Rees B, Trott DJ. Prevalence of bacterial species in cats with clinical signs of lower urinary tract disease: Recognition of *Staphylococcus felis* as a possible feline urinary tract pathogen. Vet Microbiol, 2007; 121(1-2):182–188.

6. Koneman EW, Allen SD, Janda WM, Schreckenberger PC, Winn WC. The gram-positive cocci. in, color atlas and textbook of diagnostic microbiology. 5th ed., Lippincott Raven Publishers, Philadelphia; 1997, p. 539-565.

7. CLSI. Performance Standards for Antimicrobial Susceptibility Testing; Twenty-Second Informational Supplement, CLSI document M100-S22, Wayne, PA: Clinical and Laboratory Standart Institute, 2012.

8. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. J Comput Biol, 2012; 19:455-477.

9. Langmead B, Salzberg S: Fast gapped-read alignment with Bowtie 2. Nature Methods, 2012; 9:357-359.

10. Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, et al. Identification of acquired antimicrobial resistance genes. J Antimicrob Chemother, 2012; 67:2640–2644.

11. Joensen KG, Scheutz F, Lund O, Hasman H, Kaas RS, Nielsen EM, et al. Real-time whole-genome sequencing for routine typing, surveillance, and outbreak detection of vero-toxigenic Escherichia coli. J Clin Microbiol, 2014; 52:1501–1510.

12. Brettin T, Davis JJ, Disz T, EdwardsRA, GerdesS, OlsenGJ, et al. RASTtk: a modular and extensible implementation of the RAST algorithm for building custom annotation pipelines and annotating batches of genomes. Sci Rep, 2015; 5:8365.

13. Worthing K, Pang S, Trott DJ, Abraham S, Coombs GW, Jordan D, et al. Characterisation of *Staphylococcus felis* isolated from cats using whole genome sequencing. Vet Microbiol, 2018; 222:98-104.

14. Burke FM, McCormack N, Rindi S, Speziale P, Foster JF. Fibronectin-binding protein B variation in *Staphylococcus aureus*. BMC Microbiol, 2010; 10:160.

15. Porayath C, Suresh MK, Biswas R, Nair BG, Mishra N, Pal S. Autolysin mediated adherence of *Staphylococcus aureus* with Fibronectin, Gelatin and Heparin. Int J Biol Macromol. 2018; 110:179-184.