

## PAPER DETAILS

TITLE: DARBOUX VECTOR FIELD OF THE THIRD ORDER MANNHEIM PARTNER CURVE IN

AUTHORS: Seyda KILIÇOĞLU, Süleyman SENYURT

PAGES: 131-134

ORIGINAL PDF URL: <https://dergipark.org.tr/tr/download/article-file/329409>

## DARBOUX VECTOR FIELD OF THE THIRD ORDER MANNHEIM PARTNER CURVE IN $E^3$

Şeyda KILIÇOĞLU<sup>1</sup>, Süleyman ŞENYURT<sup>2\*</sup>

<sup>1</sup>Baskent University, Primary Mathematics Teaching Program Ankara, Turkey

<sup>2</sup>Ordu University, Faculty of Arts and Sciences Department of Geometry, Ordu, Turkey

(Received June 24,2016; Accepted January 25,2017)

### Abstract

In this study we worked on the modified Darboux vector on Mannheim partner curve  $\alpha_3$  which is called the *third order Mannheim partner* of Mannheim curve  $\alpha$ . Further we give the offset property of the *third order Mannheim partner*  $\alpha_3$  based on Frenet apparatus of Mannheim curve  $\alpha$ .

**Mathematics Subject Classification (2010):** 53A04, 53A05.

**Keywords:** Mannheim curves, Frenet apparatus, third order Mannheim curve.

### Özet

Bu çalışmada  $\alpha$  Mannheim eğrisinin  $\alpha_3$  Mannheim partner eğrisine ait modified Darboux vektörünün Mannheim eğrisinin Frenet aparatlarına bağlı ifadesi verildi. Daha sonra  $\alpha_3$  eğrisinin Mannheim eğrisi olma şartı verildi.

**Mathematics Subject Classification (2010):** 53A04, 53A05.

**Keywords:** Mannheim eğrileri, Frenet aparatları, modified Darboux vektörü.

\*senyutsuleyman@hotmail.com

## 1 Introduction and Preliminaries

$\alpha : I \rightarrow E^3$  be the  $C^2$ -class differentiable unit speed and the quantities  $\{T, N, B, \kappa, \tau\}$  are collectively Frenet-Serret apparatus of the curve  $\alpha$ . Darboux vector can be expressed as (Gray 1997),

$$D(s) = \tau(s)T(s) + \kappa(s)B(s). \quad (1.1)$$

Let a vector field be

$$\tilde{D}(s) = \frac{\tau}{\kappa}(s)T(s) + B(s) \quad (1.2)$$

along  $\alpha(s)$  under the condition that  $\kappa(s) \neq 0$  and it is called the modified Darboux vector field of  $\alpha$  (Izumiya et al. 2003). A curve is called a Mannheim curve if and only if  $\frac{\kappa}{(\kappa^2 + \tau^2)}$  is a

nonzero constant,  $\kappa$  is the curvature and  $\tau$  is the torsion. Recently, a new definition of the associated curves was given by Liu and Wang (Liu & Wang 2008) Mannheim curve was redefined by Liu and Wang as ; if the principal normal vector of first curve and binormal vector of second curve are linearly dependent, then first curve is called Mannheim curve, and the second curve is called Mannheim partner curve, see in (Liu & Wang 2008) . In (Orbay & Kasap 2005) Mannheim offsets of ruled surfaces are defined and characterized. The quantities  $\{T_3, N_3, B_3, \kappa_3, \tau_3\}$  are collectively Frenet-Serret apparatus of the curve  $\alpha_3$ . We called as  $\alpha_3$  is a *third order Mannheim partner* of the curve  $\alpha$ , which has the following parameterizations, *third order Mannheim partner*  $\alpha_3$  can be written as

$$\alpha_3(s) = \alpha + (\lambda_1 \sin \theta + \lambda_2 \sin \theta_1 \cos \theta)T - (\lambda N - \lambda_2 \cos \theta_1)N + (\lambda_1 \cos \theta - \lambda_2 \sin \theta_1 \sin \theta)B \quad (1.3)$$

Also  $|\lambda + \lambda_1 + \lambda_2|$  is the distance between the arclengthed curves  $\alpha$  and  $\alpha_3$ . Since we have

$$d(\alpha(s), \alpha_3(s)) = \|\alpha_3(s) - \alpha(s)\| = \|(\lambda + \lambda_1 + \lambda_2)N(s)\| = |\lambda + \lambda_1 + \lambda_2|.$$

The Frenet apparatus of *third order Mannheim partner*  $\alpha_3$  of a Mannheim curve  $\alpha$ , based on the Frenet apparatus of Mannheim curve  $\alpha$  are

$$\begin{cases} T_3 = (\cos \theta_2 \cos \theta_1 \cos \theta - \sin \theta_2 \sin \theta)T - \cos \theta_2 \sin \theta_1 N - (\sin \theta_2 \cos \theta + \cos \theta_2 \cos \theta_1 \sin \theta)B \\ N_3 = (\sin \theta_2 \cos \theta_1 \cos \theta + \cos \theta_2 \sin \theta)T - \sin \theta_2 \sin \theta_1 N + (\cos \theta_2 \cos \theta - \sin \theta_2 \cos \theta_1 \sin \theta)B \\ B_3 = \sin \theta_1 \cos \theta T + \cos \theta_1 N - \sin \theta_1 \sin \theta B \end{cases} \quad (1.4)$$

$$\kappa_3 = \frac{-\dot{\theta}_2}{\cos \theta \cos \theta_1 \cos \theta_2} \quad \text{and} \quad \tau_3 = \frac{\lambda_1 \kappa \dot{\theta}_1}{\lambda_2 \lambda \tau \dot{\theta} \cos \theta_1} \quad (1.5)$$

are the first and second curvatures of the *third order Mannheim partner*  $\alpha_3$ , respectively ] (Kılıçoğlu & Şenyurt 2017).

## 2 Darboux vector field of the third order Mannheim partner curve in $E^3$

**Theorem 2.1** *The modified Darboux vector of third order Mannheim partner  $\alpha_3$  of a Mannheim curve  $\alpha$ , based on the Frenet apparatus of Mannheim curve  $\alpha$  is*

$$\tilde{D}_3(s) = \left[ \frac{-\lambda_1 \kappa \dot{\theta}_1 \cos \theta \cos \theta_2}{\lambda \lambda_2 \tau \dot{\theta} \dot{\theta}_2} (\cos \theta_2 \cos \theta_1 \cos \theta - \sin \theta_2 \sin \theta) + \sin \theta_1 \cos \theta \right] T$$

$$+ \left[ \frac{\lambda_1 \kappa \dot{\theta}_1 \cos \theta \cos \theta_2}{\lambda \lambda_2 \tau \dot{\theta}_2} \cos \theta_2 \sin \theta_1 - \cos \theta_1 \right] N$$

$$+ \left[ \frac{\lambda_1 \kappa \dot{\theta}_1 \cos \theta \cos \theta_2}{\lambda \lambda_2 \tau \dot{\theta}_2} (\sin \theta_2 \cos \theta + \cos \theta_2 \cos \theta_1 \sin \theta) + \sin \theta_1 \sin \theta \right] B$$

*Proof.* Since  $\tilde{D}_3(s) = \frac{\tau}{\kappa_3} T_3(s) + B_3(s)$ , and we have the proof as in the following way

$$\tilde{D}_3(s) = \frac{\tau}{\kappa_3} T_3(s) + B_3(s),$$

$$= \left[ \frac{-\lambda_1 \kappa \dot{\theta}_1 \cos \theta \cos \theta_2}{\lambda \lambda_2 \tau \dot{\theta}_2} (\cos \theta_2 \cos \theta_1 \cos \theta - \sin \theta_2 \sin \theta) + \sin \theta_1 \cos \theta \right] T$$

$$- \left[ \frac{-\lambda_1 \kappa \dot{\theta}_1 \cos \theta \cos \theta_2}{\lambda \lambda_2 \tau \dot{\theta}_2} \cos \theta_2 \sin \theta_1 + \cos \theta_1 \right] N$$

$$- \left[ \frac{-\lambda_1 \kappa \dot{\theta}_1 \cos \theta \cos \theta_2}{\lambda \lambda_2 \tau \dot{\theta}_2} (\sin \theta_2 \cos \theta + \cos \theta_2 \cos \theta_1 \sin \theta) - \sin \theta_1 \sin \theta \right] B$$

$$\tilde{D}_3(s) = \left[ \frac{-\lambda_1 \kappa \dot{\theta}_1 \cos \theta \cos \theta_2}{\lambda \lambda_2 \tau \dot{\theta}_2} (\cos \theta_2 \cos \theta_1 \cos \theta - \sin \theta_2 \sin \theta) + \sin \theta_1 \cos \theta \right] T$$

$$+ \left[ \frac{\lambda_1 \kappa \dot{\theta}_1 \cos \theta \cos \theta_2}{\lambda \lambda_2 \tau \dot{\theta}_2} \cos \theta_2 \sin \theta_1 - \cos \theta_1 \right] N$$

$$+ \left[ \frac{\lambda_1 \kappa \dot{\theta}_1 \cos \theta \cos \theta_2}{\lambda \lambda_2 \tau \dot{\theta}_2} (\sin \theta_2 \cos \theta + \cos \theta_2 \cos \theta_1 \sin \theta) + \sin \theta_1 \sin \theta \right] B$$

where we use dot to denote the derivative with respect to the arclength parameter of the curve  $\alpha$ .

**Theorem 2.2** *The offset property of the third order Mannheim partner  $\alpha_3$  based on the frenet apparatus Mannheim curve  $\alpha$ , can be given if and only if the curvature  $\kappa$  and the torsion  $\tau$  of  $\alpha$  satisfy the following equation*

$$\lambda_3 = \frac{\lambda^2(\lambda_2)^2 \tau^2 \dot{\theta}_2 \dot{\theta}_2 \cos \theta \cos \theta_1 \cos \theta_2}{\lambda^2(\lambda_2)^2 \tau^2 \dot{\theta}_2^2 + (\lambda_1 \kappa \dot{\theta}_1)^2 \cos^2 \theta \cos^2 \theta_2} = constant.$$

*Proof.* The offset property of the *third order Mannheim partner*  $\alpha_3$  is

$$\lambda_3 = \frac{-\kappa_3}{\kappa_3^2 + \tau_3^2} = constant.$$

Hence

$$\begin{aligned} \lambda_3 &= \frac{-\kappa_3}{\kappa_3^2 + \tau_3^2} = \frac{-\frac{\dot{\theta}_2}{\cos \theta \cos \theta_1 \cos \theta_2}}{\left( \frac{-\dot{\theta}_2}{\cos \theta \cos \theta_1 \cos \theta_2} \right)^2 + \left( \frac{\lambda_1 \kappa \dot{\theta}_1}{\lambda \lambda_2 \tau \dot{\theta} \cos \theta_1} \right)^2} \\ &= \frac{\frac{\dot{\theta}_2}{\cos \theta \cos \theta_1 \cos \theta_2}}{\frac{\lambda^2(\lambda_2)^2 \dot{\theta}_2^2 \tau^2 + (\dot{\theta}_1 \lambda_1 \kappa)^2 \cos^2 \theta \cos^2 \theta_2}{\lambda^2(\lambda_2)^2 \dot{\theta}_2^2 \cos^2 \theta \cos^2 \theta_1 \cos^2 \theta_2 \tau^2}} \\ &= \frac{\lambda^2(\lambda_2)^2 \tau^2 \dot{\theta}_2 \dot{\theta}_2 \cos \theta \cos \theta_1 \cos \theta_2}{\lambda^2(\lambda_2)^2 \tau^2 \dot{\theta}_2^2 + (\lambda_1 \kappa \dot{\theta}_1)^2 \cos^2 \theta \cos^2 \theta_2}. \end{aligned}$$

## References

- Gray, A. *Modern Differential Geometry of Curves and Surfaces with Mathematica*, 2nd ed. Boca Raton, FL: CRC Press, 1997.
- Izumiya, S., Takeuchi, N., Special curves and Ruled surfaces . Beitr age zur Algebra und Geometrie Contributions to Algebra and Geometry, 44 (1) (2003), 203-212.
- Kılıçoğlu S., Şenyurt S., On the second order Mannheim partner curve in  $E^3$ . International J.Math. Combin., 1 (2017), 71-77.
- Liu H. and Wang F., Mannheim partner curves in 3-space, Journal of Geometry, 88 (1-2) (2008), 120-126.
- Orbay K. and Kasap E., On mannheim partner curves, International Journal of Physical Sciences, 4 (5) (2009), 261-264.